
12 Mental health management

CONTENTS

12.1 Profile of mental health management	12.2
12.2 Framework of performance indicators	12.11
12.3 Key performance indicator results	12.14
12.4 Future directions in performance reporting	12.53
12.5 Definitions of key terms	12.53
12.6 List of attachment tables	12.58
12.7 References	12.62

Attachment tables

Attachment tables are identified in references throughout this chapter by a '12A' prefix (for example, table 12A.1). A full list of attachment tables is provided at the end of this chapter, and the attachment tables are available from the website at www.pc.gov.au/rogs/2016.

This chapter reports on the Australian, State and Territory governments' management of mental health and mental illnesses through a variety of service types and delivery settings. The chapter focuses on State and Territory governments' specialised mental health services and specific mental health services provided by General Practitioners (GPs), psychiatrists, psychologists and other allied health professionals under the Medicare Benefits Schedule (MBS).

Improvements to the reporting of mental health management in this edition include:

- additional data disaggregations for MBS subsidised services for the 'New client index' indicator
- new data on the duration of seclusion events reported as contextual information for the 'Rate of seclusion' indicator
- a new measure for the 'Social and economic inclusion of people with mental illness' indicator on the social participation of people with mental illness.

All abbreviations used in this Report are available in a complete list in volume A: Approach to performance reporting.

12.1 Profile of mental health management

Health management is the ongoing process beginning with initial client contact and including all actions relating to the client: assessment/evaluation; education of the person, family or carer(s); diagnosis; and treatment. Problems associated with adherence to treatment and liaison with, or referral to, other agencies are also included. Measuring performance in the management of a health problem involves measuring the performance of service providers in specific settings, and the overall management of diseases, illnesses and injuries across the spectrum of services, including prevention, early detection and treatment programs. The appropriate mix of services and of service delivery mechanisms is also important.

Mental health relates to an individual's ability to negotiate the daily challenges and social interactions of life without experiencing undue emotional or behavioural incapacity (DHAC and AIHW 1999). The World Health Organization describes positive mental health as:

... a state of well-being in which the individual realizes his or her own abilities, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his or her community (WHO 2001).

Mental health is identified by governments as one of the national health priority areas. The national health priority areas represent a large proportion of the total burden of disease and injury in Australia and mental illnesses makes a significant contribution to this total burden (Begg et al. 2007). The total burden comprises the number of 'years' lost due to fatal events (years of life lost due to premature death) and non-fatal events (years of 'healthy' life lost due to disability). Mental illness is the leading cause of 'healthy' life years lost due to disability (Begg et al. 2007).

Mental illness is a term that describes a diverse range of behavioural and psychological conditions. These conditions can affect an individual's mental health, functioning and quality of life. Each mental illness is unique in its incidence across the lifespan, causal factors and treatments. The most common mental illnesses are anxiety, affective (mood) and substance use disorders. Mental illness also includes low prevalence conditions such as schizophrenia, bipolar disorder and other psychoses, and severe personality disorder (DoHA 2010). While of lower prevalence, these conditions can severely affect people's ability to function in their daily lives (Morgan et al. 2011).

Mental health management is offered across a spectrum of government and non-government service providers that include promotion, prevention, treatment, management and rehabilitation services. Psychiatric hospitals, general hospitals with psychiatric units, community mental health facilities, psychiatrists, clinical psychologists, psychotherapists, mental health clinicians in private practice and counsellors all provide specialised mental health care. Mental health care is also provided in non-specialised settings — for example, GPs, public hospital emergency departments and outpatient departments, and public hospital general wards (as distinct from specialised psychiatric

units). Some people with a mental illness are cared for in residential aged care services. Mental health is also the subject of programs designed to improve public health.

This chapter focuses on the performance of State and Territory governments' specialised mental health services that treat the mostly low prevalence, but severe, mental illnesses and also on the specific mental health services provided by GPs, psychiatrists, psychologists and other allied health professionals under the MBS. It also reports on the interaction and integration arrangements between hospital and community-based services and on the broad social, economic and physical health outcomes of people with a mental illness.

Roles and responsibilities

National mental health policy context

In 1991, Australian Health Ministers signed the *Mental Health Statement of Rights and Responsibilities*. This Statement seeks to ensure that consumers, carers, advocates, service providers and the community are aware of their rights and responsibilities and can be confident in exercising them (Australian Health Ministers 1991). During 2011-12, the Statement was updated to align with the *National Mental Health Policy 2008* and Australia's international obligations with respect to the *United Nations Convention on the Rights of Persons with Disabilities* and the *United Nations Convention on the Rights of the Child*.

The Statement underpins the National Mental Health Strategy (NMHS) endorsed by Australian, State and Territory governments in 1992, but has been reaffirmed by health ministers a number of times since then (Department of Health 2014). The NMHS was established to guide the reform agenda for mental health in Australia across the whole of government. The NMHS consists of the National Mental Health Policy that describes its broad aims and objectives and the National Mental Health Plan that outlines the approach to implementing these aims and objectives. The National Mental Health Policy was revised in 2008 and the Fourth National Mental Health Plan was released in November 2009. The Fourth Plan (2009–2014) has now expired and a Fifth Plan is currently being developed.

Service roles and responsibilities

State and Territory governments are responsible for the funding, delivery and/or management of specialised mental health services including inpatient/admitted care in hospitals, community-based ambulatory care and community-based residential care. Some of these services are provided by non-government organisations (NGOs), for example governments can fund private entities to provide admitted patient hospital care. State and Territory governments also fund not-for-profit, NGOs to provide a range of support services for people with psychiatric disability arising from their mental illness.

The Australian Government is responsible for the oversight and funding of a range of mental health services and programs that are primarily provided or delivered by private practitioners or NGOs. These services and programs include MBS subsidised services provided by GPs (under both general and specific mental health items), private psychiatrists and allied mental health professionals, Pharmaceutical Benefits Scheme (PBS) funded mental health-related medications and other programs designed to prevent suicide or increase the level of social support and community-based care for people with a mental illness and their carers. The Australian Government also funds State and Territory governments for health services, most recently through the approaches specified in the National Health Reform Agreement (NHRA), but the mental health component of this funding is not separately identified for this Report.

Funding


Real government recurrent expenditure of around \$7.7 billion was allocated to mental health services in 2013-14 (table 12A.4). State and Territory governments made the largest contribution (\$4.8 billion, or 62.4 per cent, which includes Australian Government funding under the NHRA), with Australian Government expenditure of \$2.9 billion or 37.6 per cent of total government recurrent expenditure on mental health services (table 12A.4).

Real recurrent government expenditure per person on mental health services increased from \$255.44 in 2005-06 to \$330.28 in 2013-14 (figure 12.1). The average annual growth rate for Australian Government real expenditure over this period was 5.7 per cent, which was slightly higher than the 4.7 per cent for State and Territory governments (table 12A.4).

Expenditure on MBS subsidised services was the largest component of Australian Government expenditure on mental health services in 2013-14 (\$971.0 million or 33.5 per cent) (table 12A.1). This comprised MBS payments for psychologists and other allied health professionals (15.0 per cent), consultant psychiatrists (11.1 per cent) and GP services (7.5 per cent) (table 12A.1). Another significant area of Australian Government expenditure on mental health services in 2013-14 was expenditure under the PBS for mental-health related medications (\$735.0 million) (table 12A.1).

Nationally, expenditure on admitted patient services is the largest component of State and Territory governments' expenditure on specialised mental health services (\$2.1 billion or 43.4 per cent in 2013-14), followed by expenditure on community-based ambulatory services (\$1.9 billion or 38.3 per cent) (figure 12.2). State and Territory governments' expenditure on specialised mental health services, by source of funds and depreciation (which is excluded from reporting) are in tables 12A.3 and 12A.5 respectively.


Figure 12.1 **Real recurrent governments' expenditure on mental health services, by funding source (2013-14 dollars)^a**



^a See table 12A.4 for detailed footnotes and caveats.

Source: Department of Health (unpublished); Australian Institute of Health and Welfare (AIHW) (unpublished) Mental Health Establishments (MHE) National Minimum Data Set (NMDS); table 12A.4.

Figure 12.2 **Recurrent expenditure on State and Territory governments' specialised mental health services, by service category, 2013-14^{a, b}**



^a Queensland does not have any in-scope community residential services. ^b See table 12A.6 for detailed footnotes and caveats.

Source: AIHW (unpublished) MHE NMDS; table 12A.6.

Size and scope of sector


Prevalence of mental illness and psychological distress

According to the National Survey of Mental Health and Wellbeing (SMHWB), in 2007, 1 in 5 people aged 16–85 years (20.0 per cent, or approximately 3.2 million adults) met the criteria for diagnosis of a lifetime mental disorder/illness and had symptoms in the previous 12 months, and a further 25.5 per cent of people in this age group had experienced a mental disorder at some point in their life (table 12A.76).

A proxy measure of the overall mental health and wellbeing of the population is the Kessler 10 (K10) psychological distress scale. Very high levels of psychological distress may signify a need for professional help and provide an estimate of the need for mental health services (ABS 2012). Higher levels of psychological distress are reported for:

- females compared to males (figure 12.3)
- people with disability compared to those without (table 12A.9)
- people in lower socioeconomic areas compared to those in higher areas (table 12A.9)
- Aboriginal and Torres Strait Islander Australians compared to non-Indigenous Australians (table 12A.15).

Figure 12.3 **Adults with very high levels of psychological distress, by gender, 2011-12^a**



^a See table 12A.8 for detailed footnotes and caveats.

Source: ABS (unpublished) *Australian Health Survey (AHS) 2011–13* (2011-12 National Health Survey (NHS) component), Cat. no. 4364.0; table 12A.8.

There is also a strong association between a high/very high K10 score and a current diagnosis of anxiety and affective disorders, and people with a mental illness are overrepresented in the population who had high/very high levels of psychological distress (ABS 2012 and table 12A.7).

Tables 12A.8–16 contain additional data on high/very high levels of psychological distress.

Mental health services — overview

There are a range of Australian, State or Territory governments' provided or funded services that are specifically designed to meet the needs of people with mental health issues; the key services are:

- MBS subsidised mental health specific services that are partially or fully funded under Medicare on a fee for service basis and are provided by GPs, psychiatrists, psychologists or other allied health professionals under specific mental health items.
- Admitted patient care in public hospitals — specialised services provided to inpatients in stand-alone psychiatric hospitals or psychiatric units in general acute hospitals.
- Community-based public mental health services, comprising:
 - ambulatory care services provided by outpatient clinics (hospital and clinic based), mobile assessment and treatment teams, day programs and other services dedicated to assessment, treatment, rehabilitation and care
 - residential services that provide beds in the community, staffed onsite by mental health professionals
- Not-for-profit, NGO services, funded by the Australian, State and Territory governments to provide community-based support for people with psychiatric disability, including accommodation, outreach to people living in their own homes, residential rehabilitation units, recreational programs, self-help and mutual support groups, carer respite services and system-wide advocacy (DoHA 2010).

There are also other health services provided and/or funded by governments that make a significant contribution to the mental health treatment of people with a mental illness, but are not specialised or specific mental health services. Tables 12A.30–32 provide information on these non-specialised services provided in hospitals.

MBS subsidised mental health services


GPs are often the first type of service accessed by people seeking help when suffering from a mental illness (AIHW 2014). GPs can diagnose, manage and treat mental illnesses and they also refer patients to more specialised service providers such as psychiatrists and psychologists.

In 2013-14, an estimated 12.8 per cent of GP encounters included management of mental health-related problems (equivalent to an estimated 17.1 million encounters) (table 12A.18). Data on GP mental-health related encounters by patient demographics are in table 12A.19.

A GP can manage more than one problem at a single encounter, hence the number of mental health problems can be greater than the number of mental-health related encounters. In 2013-14, 13.7 mental health-related problems were managed per 100 encounters (table 12A.20). Depression was the most frequently reported mental health-related problem managed (4.3 per 100 GP encounters), representing around one third of all mental health-related problems managed (table 12A.20).

In 2013-14, GPs provided 2.7 million MBS subsidised specific mental health items. A further 6.4 million MBS subsidised mental health services were provided by psychiatrists (2.2 million), psychologists (3.9 million) and allied health professionals (0.3 million) (table 12A.17). Service usage rates varied across states and territories (figure 12.4).

Figure 12.4 **MBS subsidised mental-health related services, by provider type, 2013-14^a**



^a See table 12A.17 for detailed footnotes and caveats.


Source: AIHW (2015) *Mental Health Services in Australia* (available at <http://mhsa.aihw.gov.au/home/>); table 12A.17.

State and Territory governments' specialised mental health services

State and Territory governments' specialised mental health services (covering the three service types of: admitted patient, community-based ambulatory and community-based residential) tend to treat people with the lower prevalence, but severe, mental illnesses. The

proportion of the total Australian population treated in these public services remained below 2 per cent between 2007-08 and 2013-14 (table 12A.42 and figure 12.5).

Figure 12.5 **Population using State and Territory governments' specialised mental health services^{a, b}**




^a Victorian 2011-12 and 2012-13 data are not available. ^b See table 12A.42 for detailed footnotes and caveats.

Source: AIHW (unpublished), derived from data provided by State and Territory governments; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period); table 12A.42.

Across states and territories, the mix of admitted patient and community-based services and care types can differ. As the unit of activity varies across these three service types (table 12A.25), service mix differences can be partly understood by considering items which have comparable measurement across service types such as expenditure (figure 12.2), numbers of full time equivalent (FTE) direct care staff (figure 12.6), accrued mental health patient days (figure 12.7) and mental health beds (figure 12.8).

Additional data are also available on the most common principal diagnosis for admitted patients (tables 12A.22 and 12A.33) and community-based ambulatory contacts by age group (table 12A.24).


Figure 12.6 **FTE health professional direct care staff, by service type, 2013-14^{a, b}**



^a Queensland does not have any in scope residential services. ^b See table 12A.28 for detailed footnotes and caveats.

Source: AIHW (unpublished) MHE NMDS; table 12A.28.


Figure 12.7 **Accrued mental health patient days, by service type, 2013-14^{a, b, c}**



^a Queensland does not have any in-scope residential services. ^b The ACT and the NT do not have non-acute hospital units. ^c See table 12A.21 for detailed footnotes and caveats.

Source: AIHW (unpublished) MHE NMDS; table 12A.21.

Figure 12.8 **Mental health beds in public hospitals and community-based residential units, 2013-14^{a, b, c}**



^a Queensland does not have any in-scope residential services. ^b Tasmania, the ACT and the NT do not have public psychiatric hospitals. ^c See table 12A.26 for detailed footnotes and caveats.

Source: AIHW (unpublished) MHE NMDS; table 12A.26.

Government funded not-for-profit, NGO services

There are limited data available on the size and scope of the mental health services provided by the Australian, State and Territory governments' funded not-for-profit, NGO sector. The targeted community care (Mental Health) program is one exception. In 2013-14, there were 157 670 participants in the program across three service types: 18 539 for Personal Helpers and Mentors (PHaMs), 98 664 for Family Mental Health Support Services and 40 467 for Mental Health Respite: Carer Support (table 12A.29).

12.2 Framework of performance indicators

The framework of performance indicators for mental health services draws on governments' broad objectives as expressed in the *National Mental Health Policy 2008* (box 12.1).

Box 12.1 **Broad objectives and policy directions of National Mental Health Policy**

The *National Mental Health Policy 2008* has an emphasis on whole-of-government mental health reform and commits the Australian, State and Territory governments to the continual improvement of Australia's mental health system. The key broad objectives are to:

- promote the mental health and well-being of the Australian community and, where possible, prevent the development of mental health problems and mental illness
- reduce the impact of mental health problems and mental illness, including the effects of stigma on individuals, families and the community
- promote recovery from mental health problems and mental illness
- assure the rights of people with mental health problems and mental illness, and enable them to participate meaningfully in society.


The key policy directions are summarised as follows:

- Rights and responsibilities of people with mental health problems and mental illness will be acknowledged and respected.
- Mental health promotion will support destigmatisation and assist people to be emotionally resilient, cope with negative experiences and participate in the community.
- The proportion of people with mental health problems, mental illness and people at risk of suicide will be reduced.
- Emerging mental health problems or mental illnesses will receive early intervention to minimise the severity and duration of the condition and to reduce its broader impacts.
- People will receive timely access to high quality, coordinated care appropriate to their conditions and circumstances.
- People with mental health problems and mental illness will enjoy full social, political and economic participation in their communities.
- The crucial role of carers will be acknowledged and respected and they will be provided with appropriate support to enable them to fulfil their role.
- The mental health workforce will be appropriately trained and adequate in size and distribution to meet the need for care.
- Across all sectors, mental health services should be monitored and evaluated to ensure they are of high quality and achieving positive outcomes.
- Research and evaluation efforts will generate new knowledge about mental health problems and mental illness that can reduce the impact of these conditions.

The performance indicator framework provides information on equity, efficiency and effectiveness, and distinguishes the outputs and outcomes of mental health management services (figure 12.9). It covers a number of service delivery types (MBS subsidised, admitted patient and community-based services) and includes outcome indicators of system-wide performance. The performance indicator framework shows which data are complete and comparable in the 2016 Report. Chapter 1 discusses data comparability from a Report-wide perspective (see chapter 1, section 1.6).

In addition to section 12.1, the Report’s statistical context chapter contains data that may assist in interpreting the performance indicators presented in this chapter. These data cover a range of demographic and geographic characteristics (chapter 2).

Figure 12.9 **Mental health management performance indicator framework**



12.3 Key performance indicator results

Different delivery contexts, locations and types of clients can affect the equity, effectiveness and efficiency of mental health management services.

Data Quality Information (DQI) is included where available for performance indicators in this Report. The purpose of DQI is to provide structured and consistent information about quality aspects of data used to report on performance indicators, in addition to material in the chapter or sector overview and attachment tables. All DQI for the 2016 Report can be found at www.pc.gov.au/rogs/2016.

Outputs

Outputs are the services delivered (while outcomes are the impact of these services on the status of an individual or group) (see chapter 1, section 1.5). Output information is also critical for equitable, efficient and effective management of government services.

Equity

Access — new client index

‘New client index’ is an indicator of governments’ objective to provide mental health services in an equitable manner (box 12.2). Where population treatment rates are relatively low it may be difficult for a new client to access services if already used by existing clients.

Box 12.2 New client index

‘New client index’ is defined by two measures, the proportions of total clients under the care of:

- State and Territory governments’ specialised public mental health services, who were new — clients include all people who received one or more community-based ambulatory contact or had one or more day of admitted patient or community-based residential care.
- MBS subsidised mental health services provided by private psychiatrists, GPs and allied health providers, who were new.

A new client is a consumer who has not received a mental health service in the five years preceding the initial contact with a service in the relevant reference period.

(continued next page)

Box 12.2 (continued)

A high or increasing proportion of total clients who are new might be desirable, as it can suggest better access for new clients. However, results are difficult to interpret as the appropriate balance between providing ongoing care to existing clients who have continuing needs and meeting the needs of new clients is unknown. In addition, a significant increase in the proportion of new clients accessing services might be the result of an increase in the prevalence of mental illness.

This indicator does not provide information on whether the services are appropriate or adequate for the needs of the people receiving them (new or existing clients), or correctly targeted to those clients who are most in need.

Data reported for the proportions of total clients under the care of State and Territory governments' specialised public mental health services who were new are:

- comparable (subject to caveats) within some jurisdictions over time but are not comparable across jurisdictions or over time for other jurisdictions (see caveats in attachment tables for specific jurisdictions)
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions.

Data reported for the proportions of total clients under the care of MBS subsidised mental health services who were new are:


- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required 2014-15 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally, the proportion of State and Territory governments' specialised mental health services clients who are new remained relatively stable over time (figure 12.10). These proportions tend to be higher than for clients of MBS subsidised mental health services nationally and for all jurisdictions, except the NT (figure 12.11).

For Aboriginal and Torres Strait Islander Australians, the proportion of clients who are new tend to be lower than for non-Indigenous Australians for State and Territory governments' specialised mental health services. By contrast, for MBS subsidised mental health services, the proportion of Aboriginal and Torres Strait Islander Australian clients who are new tends to be slightly higher than for non-Indigenous Australians (tables 12A.34 and 12A.36). Data are also reported by age, gender and remoteness in tables 12A.34 and 12A.36 and Socio-Economic Indexes for Areas (SEIFA) quintiles in table 12A.34.

Figure 12.10 Proportion of State and Territory governments' specialised mental health service clients who are new^{a, b}



^a Victorian 2011-12 and 2012-13 data are not available. ^b See box 12.2 and table 12A.33 for detailed definitions, footnotes and caveats.

Source: AIHW (unpublished) derived from State and Territory governments' data; table 12A.33.

Figure 12.11 Proportion of MBS subsidised mental health service clients who are new^a



^a See box 12.2 and table 12A.35 for detailed definitions, footnotes and caveats.

Source: Australian Government Department of Health (unpublished); table 12A.35.

Access — mental health service use by selected community groups

‘Mental health service use by selected community groups’ is an indicator of governments’ objective to provide mental health services in an equitable manner, including access to services by selected community groups such as Aboriginal and Torres Strait Islander Australians (box 12.3).

Box 12.3 Mental health service use by selected community groups

‘Mental health service use by selected community groups’ is defined by two measures:

- the proportion of the population in a selected community group using the service, compared to the proportion of the population outside the selected community group, for each of:
 - State and Territory governments’ specialised public mental health services
 - MBS subsidised mental health services.

The selected community groups reported are Aboriginal and Torres Strait Islander Australians, people from outer regional, remote and very remote locations and people residing in low socioeconomic areas.

This indicator is difficult to interpret. Variations in use could be due to variations in access, but could also be a result of differences in the prevalence of mental illness. It also does not provide information on whether the services are appropriate for the needs of the people receiving them, or correctly targeted to those most in need.

Data reported for the State and Territory governments’ specialised public mental health services measure:

- may not be comparable (subject to caveats) within jurisdictions over time and may not be comparable across jurisdictions (see caveats in DQI and attachment tables for details)
- are complete (subject to caveats) for the current reporting period (subject to caveats). All required 2013-14 data are available.

Data reported for the MBS subsidised mental health services measure are:

- comparable (subject to caveats) across jurisdictions but a break in series means that data from 2011-12 onwards by geographic location and SEIFA are not comparable to data for previous years’ (see caveats in DQI and attachment tables for details)
- complete (subject to caveats) for the current reporting period (subject to caveats). All required 2013-14 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

While a higher proportion of the population access MBS subsidised mental health services than State and Territory governments’ specialised mental health services, the pattern of service use across the selected community groups differs.


For State and Territory governments specialised public mental health services, across all the selected community groups, higher proportions of people within these groups (Aboriginal and Torres Strait Islander Australians, people from outer regional, remote and very remote areas and people residing in low socioeconomic areas) access these services

than those outside these groups (figures 12.12-13 and table 12A.39).

For MBS subsidised mental health services the results are mixed. Nationally, a similar proportion of Aboriginal and Torres Strait Islander Australians accessed these services to non-Indigenous Australians (figure 12.12), likewise for people across different socioeconomic areas (table 12A.39). Results varied across states and territories. However, for people in outer regional, remote and very remote areas, the proportions accessing MBS subsidised services were lower than for people in inner regional and major cities both nationally and across all states (figure 12.13).

Additional data on the use of State and Territory governments' specialised and MBS subsidised mental health services by community groups are in tables 12A.40–41. Data on the use of private hospital mental health services are also contained in tables 12A.37–39 and 12A.41–42.


Figure 12.12 Population using mental health services, by Indigenous status and service type, 2013-14^a



^a See box 12.3 and table 12A.37 for detailed definitions, footnotes and caveats.

Source: AIHW (unpublished), derived from data provided by State and Territory governments and the Australian Government Department of Health and Department of Veterans' Affairs (DVA); ABS (unpublished) Estimated Residential Population, 30 June 2013; table 12A.37.

Figure 12.13 Population using mental health services, by geographic location and service type, 2013-14^{a, b, c}



^a See box 12.3 and table 12A.38 for detailed definitions, footnotes and caveats. ^b Victoria does not have very remote areas. Tasmania does not have major cities. The ACT does not have outer regional, remote or very remote locations. The NT does not have major cities or inner regional locations. ^c ACT data are not published for inner regional areas.

Source: AIHW (unpublished), derived from data provided by State and Territory governments and the Australian Government Department of Health and DVA; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period); table 12A.38.

Effectiveness

Access — mental health service use by total population

‘Mental health service use by total population’ is an indicator of governments’ objective to provide equitable access to mental health services for all people who need them (box 12.4).

Box 12.4 Mental health service use by total population

'Mental health service use by total population' is defined as the proportion of the population using a State and Territory specialised public mental health service or a MBS subsidised mental health service. Data are reported separately for State and Territory specialised public mental health services and MBS subsidised mental health services.

This indicator is difficult to interpret. As a robust estimate of the population who need mental health services is not available, the indicator is reported as a proportion of the total population. Variations in use could be due to variations in access, but could also be a result of differences in the prevalence of mental illness.

This indicator does not provide information on whether the services are appropriate for the needs of the people receiving them, or correctly targeted to those most in need. People with a mental illness can have low rates of service use due to them choosing not to access services, appropriate services are unavailable, lack of awareness that services are available and negative experiences associated with the previous use of services (AHMC 2008). In addition, it might not be appropriate for all people with a mental illness to use a service, for example, some can seek and receive assistance from outside the health system (AHMC 2008).

Data reported for the State and Territory governments' specialised public mental health services measure are:

- comparable (subject to caveats) within most jurisdictions over time but are not comparable across jurisdictions or over time for Tasmania (see caveats in DQI and attachment tables for specific jurisdictions)
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available.


Data reported for the MBS subsidised mental health services measure are:

- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

In 2013-14, 1.8 per cent and 8.4 per cent of the total population received State and Territory governments' specialised mental health services and MBS subsidised services, respectively (figure 12.14). While the proportion of the population using State and Territory governments' specialised mental health services has remained relatively constant (figure 12.5), the proportion using MBS subsidised services has increased steadily over time from 4.9 per cent in 2007-08 to 8.4 per cent in 2013-14 (table 12A.42). Much of this growth has come from greater utilisation of GP mental health specific services (from 3.5 to 6.7 per cent) and other allied health services (1.3 per cent to 2.5 per cent) over that period (table 12A.42). Data from the 2007 SMHWB on the proportion of people with an mental illness who did/did not use services for their mental health are reported in table 12A.43.

Figure 12.14 **Population receiving mental health services, by service type, 2013-14^a**



^a See box 12.4 and table 12A.42 for detailed definitions, footnotes and caveats.

Source: AIHW (unpublished) derived from data provided by State and Territory governments and Australian Government, Department of Health and DVA; ABS (unpublished) *Estimated Residential Population*, 30 June (prior to relevant period); table 12A.42.

Data from the 2007 SMHWB on the proportion of people who had a lifetime mental disorder with symptoms in the 12 months before the survey who used any service for mental health are also reported in tables 12A.42–43.

Access — primary mental health care for children and young people

‘Primary mental health care for children and young people’ is an indicator of governments’ objective to prevent, where possible, the development of mental health problems and mental illness and undertake early intervention for mental health problems and mental illness (box 12.5).

Box 12.5 **Primary mental health care for children and young people**

‘Primary mental health care for children and young people’ is defined as the proportion of young people aged under 25 years who received a mental health care service subsidised through the MBS from a GP, psychologist or an allied health professional.

High or increasing proportions of young people who had contact with MBS subsidised primary mental health care services is desirable.

(continued next page)

Box 12.5 (continued)

This indicator does not provide information on whether the services are appropriate for the needs of the young people receiving them, or correctly targeted to those young people most in need. Variations in use could be due to variations in access, but could also be a result of differences in the prevalence of mental illness.

Results for this indicator should be interpreted with caution as some primary mental health services for children and young people are excluded; for example, community health centres, school and university counsellors and health nurses and some mental health care provided by State and Territory governments' specialised mental health services (NMHPSC 2011a).


Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required 2014-15 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

The proportion of all children and young people who receive MBS subsidised primary mental health care services has increased gradually over time (table 12A.45). The proportion increases as age increases, with the highest proportion for young people aged 18–24 years (10.2 per cent of this population receiving these primary mental health care services nationally in 2014-15) (figure 12.15).

Figure 12.15 Children and young people who received MBS subsidised primary mental health care, by age group, 2014-15^a



^a See table 12A.45 for detailed definitions, footnotes and caveats.

Source: Australian Government Department of Health (unpublished); table 12A.45.

Data on the proportion of young people who had received MBS subsidised primary mental health care services by Indigenous status, SEIFA, gender and service type are also available in tables 12A.46–47, with the pattern of results reflecting those for the total population (all ages).

Appropriateness — services reviewed against the National Standards

‘Services reviewed against the National Standards’ is an indicator of governments’ objective to provide mental health services that are appropriate (box 12.6).

Box 12.6 Services reviewed against the National Standards

‘Services reviewed against the National Standards’ is defined as the proportion of expenditure on State and Territory governments’ specialised public mental health services that had completed a review by an external accreditation agency against the National Standards for Mental Health Services (NSMHS) and were assessed at level 1. The assessment levels are defined as:

- *Services at level 1* — services reviewed by an external accreditation agency and judged to have met all National Standards.
- *Services at level 2* — services reviewed by an external accreditation agency and judged to have met some but not all National Standards.
- *Services at level 3* — services (i) in the process of being reviewed by an external accreditation agency but the outcomes are not known, or (ii) booked for review by an external accreditation agency.
- *Services at level 4* — services that do not meet criteria detailed under levels 1 to 3.

A high or increasing proportion of expenditure on specialised mental health services that had completed a review by an external accreditation agency and had been assessed against the NSMHS as level 1 is desirable.

This is a process indicator of appropriateness, reflecting progress made in meeting the NSMHS. It does not provide information on whether the standards or assessment process are appropriate. In addition, services that had not been assessed do not necessarily deliver services of lower quality. Some services that had not completed an external review included those that were undergoing a review and those that had booked for review and were engaged in self-assessment preparation.

Data reported for this indicator are:

- comparable (subject to caveats) within jurisdictions over time but are not comparable across jurisdictions
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

The first NSMHS were developed under the *First National Mental Health Plan 1993–1998* and revised NSMHS were released in September 2010 (box 12.7).

Box 12.7 The 2010 NSMHS


The 2010 NSMHS comprise 10 overarching standards:

- | | |
|-------------------------------------|--|
| 1. Rights and responsibilities | 6. Consumers |
| 2. Safety | 7. Carers |
| 3. Consumer and carer participation | 8. Governance, leadership and management |
| 4. Diversity responsiveness | 9. Integration |
| 5. Promotion and prevention | 10. Delivery of care |

Source: Australian Health Ministers' Conference (2010) *National Standards for Mental Health Services 2010*, Canberra.

Figure 12.16 shows the proportion of expenditure on specialised public mental health services that had completed an external review against the NSMHS and met 'all standards' (level 1).

Figure 12.16 **Share of expenditure on specialised public mental health services reviewed against the NSMHS, by assessment level, 30 June 2014^a**



^a See box 12.6 and table 12A.48 for detailed definitions, footnotes and caveats.

Source: AIHW (unpublished) MHE NMDS; table 12A.48.

Appropriateness — services provided in the appropriate setting

'Services provided in the appropriate setting' is an indicator of governments' objective to provide mental health services in community-based settings wherever possible (box 12.8).

Box 12.8 Services provided in the appropriate setting

'Services provided in the appropriate setting' is defined as the proportion of State and Territory governments' recurrent expenditure on specialised public mental health services that was on community-based services. Community-based services expenditure comprises that on ambulatory care, adult residential services, and NGOs. Older people's residential expenditure is excluded to improve comparability.

A high or increasing proportion of recurrent expenditure spent on community-based services is desirable, reflecting a greater reliance on services that are based in community settings.


Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

State and Territory governments' expenditure on community-based services as a proportion of total expenditure on specialised mental health services has remained stable over time (figure 12.17).

Figure 12.17 State and Territory governments' expenditure on community-based services as a proportion of total expenditure on specialised mental health services^a



^a See box 12.8 and table 12A.49 for detailed definitions, footnotes and caveats.

Source: AIHW (unpublished) MHE NMDS; table 12A.49.

Appropriateness — collection of information on consumers' outcomes

'Collection of information on consumers' outcomes' is an indicator of governments' objective that consumer outcomes be monitored (box 12.9).

Box 12.9 **Collection of information on consumers' outcomes**

'Collection of information on consumers' outcomes' is defined as the proportion of State and Territory governments' specialised public mental health service episodes (by client type) with completed clinical mental health outcome measures.

High or increasing proportions of episodes for which information on consumers' mental health outcomes is collected is desirable.

This is a process indicator and monitors the uptake of the routine National Outcomes Casemix Collection. It does not provide information on whether consumers had appropriate outcomes.


Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required data for 2013-14 are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

The estimated proportions of specialised mental health service episodes for which 'complete' consumers' mental health outcomes are collected are shown in figure 12.18.

Figure 12.18 **Estimated proportion of episodes for which ‘complete’ outcome measures were collected, by client type 2013-14^{a, b}**



^a See box 12.9 and table 12A.50 for detailed definitions, footnotes and caveats. ^b ACT data for consumers discharged from ambulatory care are not published.

Source: AIHW (unpublished) from data provided by the Australian Mental Health Outcomes and Classification Network; table 12A.50.

Quality — safety — rate of seclusion — acute inpatient units

‘Rate of seclusion — acute inpatient units’ is an indicator of governments’ objective that services are safe and of a high quality (box 12.10). The reduction, and where possible elimination of, seclusion in specialised mental health services is a national safety priority.

Box 12.10 **Rate of seclusion — acute inpatient units**

‘Rate of seclusion — acute inpatient units’ is defined as the number of seclusion events per 1000 bed days in State and Territory governments’ specialised mental health acute inpatient units.

Seclusion involves a patient being confined at any time of the day or night alone in a room or area from which it is not within their control to leave (see section 12.5 for further details on seclusion and ‘seclusion events’). Legislation or mandatory policy governs the use of seclusion in each State and Territory and may result in exceptions to the definition of a seclusion event and variations in the data collected across jurisdictions (NMHPSC 2011b).

(continued next page)

Box 12.10 (continued)

A low (or nil) or decreasing number of seclusion events per 1000 bed days in specialised public mental health inpatient units is desirable.

Data reported for this indicator are:


- comparable (subject to caveats) within jurisdictions over time but are not comparable across jurisdictions
- complete (subject to caveats) for the current reporting period. All required data for 2014-15 are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Supporting data on the duration of seclusion events are provided in table 12A.51. These data when considered with the rate of seclusion provide information on the use and management of seclusion within each jurisdiction. A low rate of seclusion events combined with shorter average durations is desirable.

Nationally, the number of seclusion events per 1000 bed days has decreased steadily from 11.8 in 2010-11 to 7.8 in 2014-15 (figure 12.19). This downward trend was reflected across most jurisdictions (figure 12.19) and target population groups (table 12A.52). The lowest seclusion rates were in older people's units and the highest were in children and adolescent units (table 12A.52).

Figure 12.19 Rate of seclusion^a



^a See box 12.10 and table 12A.51 for detailed definitions, footnotes and caveats.

Source: AIHW (2015) *Mental Health Services in Australia Online*, mhsa.aihw.gov.au/home/ (accessed 16 December 2015); table 12A.51.

Quality — responsiveness — consumer and carer experiences of services

‘Consumer and carer experiences of services’ is an indicator of governments’ objective that services are of a high quality and responsive to the needs of consumers and their carers (box 12.11).

Box 12.11 Consumer and carer experiences of services

‘Consumer and carer experiences of services’ is yet to be defined.

Data for this indicator were not available for the 2016 Report.

Quality — responsiveness — consumer and carer involvement in decision making

‘Consumer and carer involvement in decision making’ is an indicator of governments’ objective that consumers and carers are involved at the service delivery level, where they have the opportunity to influence the services they receive (box 12.12).

Box 12.12 Consumer and carer involvement in decision making

‘Consumer and carer involvement in decision making’ is defined by two measures, the number of paid FTE:

- consumer staff per 1000 FTE direct care staff
- carer staff per 1000 FTE direct care staff.

High or increasing proportions of paid FTE direct care staff who are consumers or carers implies better opportunities for consumers and carers to be involved at the service delivery level, where they can influence the services received.


Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions but a break in series means that data from 2010-11 are not comparable to data for previous years
- complete (subject to caveats) for the current reporting period. All required data for 2013-14 are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

The number of paid FTE consumer and carer staff per 1000 paid FTE direct care staff are reported in figures 12.20 and 12.21 respectively.


Figure 12.20 **Paid FTE consumer staff per 1000 paid FTE direct care staff**
a, b, c, d



a See box 12.12 and table 12A.53 for detailed definitions, footnotes and caveats. **b** Tasmania did not employ consumer staff in 2012-13. **c** The ACT do not employ consumer staff. **d** The NT did not employ consumer staff in 2010-11 and 2011-12.

Source: AIHW (unpublished) MHE NMDS; table 12A.53.

Figure 12.21 **Paid FTE carer staff per 1000 paid FTE direct care staff**^{a, b}



a See box 12.12 and table 12A.53 for detailed definitions, footnotes and caveats. **b** WA did not employ carer staff in 2013-14 and the ACT and the NT do not employ carer staff.

Source: AIHW (unpublished) MHE NMDS; table 12A.53.

Quality — continuity — specialised public mental health service consumers with nominated GP

‘Specialised public mental health service consumers with nominated GP’ is an indicator of governments’ objective to provide continuity of care in the delivery of mental health services. GPs can be an important point of contact for those with a mental illness (box 12.13).

Box 12.13 Specialised public mental health service consumers with nominated GP

‘Proportion of specialised public mental health service consumers with nominated GP’ is yet to be defined.

Data for this indicator were not available for the 2016 Report.

Quality — continuity — community follow-up after psychiatric admission/hospitalisation

‘Community follow-up after psychiatric admission/hospitalisation’ is an indicator of governments’ objective to provide continuity of care in the delivery of mental health services (box 12.14).

Box 12.14 Community follow-up after psychiatric admission/hospitalisation

‘Community follow-up after psychiatric admission/hospitalisation’ is defined as the proportion of State and Territory governments’ specialised public admitted patient overnight acute separations from psychiatric units for which a community-based ambulatory contact was recorded in the seven days following separation.

A high or increasing rate of community follow-up within the first seven days of discharge from hospital is desirable.

This indicator does not measure the frequency of contacts recorded in the seven days following separation. It also does not distinguish qualitative differences between the mode of contact. Only follow-up contacts made by State and Territory governments’ specialised public mental health services are included.


Data reported for this indicator are:

- comparable (subject to caveats) within some jurisdictions over time but are not comparable across jurisdictions or over time for other jurisdictions (see caveats in DQI and attachment tables for specific jurisdictions)
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

For people who are discharged from hospital after an acute psychiatric episode, it is essential to have prompt community follow-up in the vulnerable period following discharge in order to maintain clinical and functional stability and minimise the need for hospital readmission (NMHPSC 2011a). Nationally, the rate of community follow-up for people within the first seven days of discharge from an acute inpatient psychiatric unit has increased from 49.6 per cent in 2009-10 to 66.4 in 2013-14 (figure 12.22). Community follow-up rates data by Indigenous status, remoteness areas, SEIFA, age groups and gender are in tables 12A.55–56.

Figure 12.22 Community follow-up for people within the first seven days of discharge from acute inpatient psychiatric units^{a, b}



^a See box 12.14 and table 12A.54 for detailed definitions, footnotes and caveats. ^b Victorian data are not available for 2011-12 and 2012-13.

Source: AIHW (unpublished), from data provided by State and Territory governments; table 12A.54.

Quality — continuity — readmissions to hospital within 28 days of discharge

‘Readmissions to hospital within 28 days of discharge’ is an indicator of governments’ objective to provide effective and continuous care in mental health services (box 12.15).

Box 12.15 Readmissions to hospital within 28 days of discharge

‘Readmissions to hospital within 28 days of discharge’ is defined as the proportion of State and Territory governments’ admitted patient overnight separations from psychiatric acute inpatient units that were followed by readmission to a State and Territory governments’ psychiatric acute inpatient unit within 28 days of discharge.

(continued next page)

Box 12.15 (continued)

A low or decreasing rate of readmissions to hospital within 28 days of discharge is desirable. Readmissions following a recent discharge can indicate that inpatient treatment was either incomplete or ineffective, or that follow-up community care was inadequate to maintain people out of hospital (NMHPSC 2011a).

Readmission rates can be affected by factors other than deficiencies in specialised public mental health services, such as the cyclic and episodic nature of some illnesses (National Mental Health Working Group Information Strategy Committee Performance Indicator Drafting Group 2005).


Data reported for this indicator are:

- comparable (subject to caveats) within some jurisdictions over time but are not comparable across jurisdictions or over time for other jurisdictions (see caveats in DQI or attachment tables for specific jurisdictions)
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally, the rates of readmission to hospital acute psychiatric units within 28 days of discharge have remained relatively stable since 2009-10 (figure 12.23). Rates of readmission to hospital within 28 days of discharge by Indigenous status, remoteness areas, SEIFA, age group and sex are in table 12A.58.

Figure 12.23 Readmissions to hospital acute psychiatric units within 28 days^a



^a See box 12.15 and table 12A.57 for detailed definitions, footnotes and caveats.

Source: AIHW (unpublished), from data provided by State and Territory governments; table 12A.57.

Efficiency

The efficiency indicators reported are for State and territory governments' specialised mental health services. Mainstreaming has occurred at different rates across states and territories, with some treating a greater proportion of clients with severe mental illnesses in community-based services than other jurisdictions. This can create differences across states and territories in the mix of clients, and therefore the costs, within service types.

Efficiency — Sustainability

The Steering Committee has identified sustainability as an area for reporting but no indicators have yet been identified.

Efficiency — cost of inpatient care

'Cost of inpatient care' is an indicator of governments' objective that mental health services are delivered in an efficient manner (box 12.16).

Box 12.16 Cost of inpatient care

'Cost of inpatient care' has two measures:

- 'Cost per inpatient bed day' is defined as expenditure on inpatient services divided by the number of inpatient bed days — data are disaggregated by hospital type (psychiatric and general hospitals) and care type (acute and non-acute units) and by inpatient target population (acute units only).
- 'Average length of stay' is defined as the number of inpatient patient days divided by the number of separations in the reference period — data are disaggregated by inpatient target population (acute units only). Patient days for clients who separated in the reference period (2013-14) that were during the previous period (2012-13) are excluded. Patient days for clients who remain in hospital (that is, are not included in the separations data) are included.

These measures are considered together for the inpatient acute units by target population to provide a 'proxy' measure to improve understanding of service efficiency. Average inpatient bed day costs can be reduced with longer lengths of stay because the costs of admission, discharge and more intensive treatment early in a stay are spread over more days of care.

A low or decreasing cost per inpatient bed day combined with similar or shorter average lengths of stay can indicate more efficient service delivery, although efficiency data need to be interpreted with care as they do not provide any information on the quality of service provided.

This indicator does not account for differences in the client mix. The client mix in inpatient settings can differ — for example, some jurisdictions treat a higher proportion of less complex patients in inpatient settings as distinct from treating them in the community. Measures that adjust to take into account the type and complexity of cases would be more appropriate but the data needed are not yet available.

(continued next page)

Box 12.16 (continued)


Data reported for the two measures for this indicator are:

- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions providing the services.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally in 2013-14, the average cost per inpatient bed day was higher in acute than non-acute units and slightly higher in psychiatric hospitals than in general hospitals for both acute and non-acute units (figure 12.24).

Figure 12.24 Average recurrent cost per inpatient bed day, by public hospital and care type, 2013-14^{a, b, c, d}




^a See box 12.16 and table 12A.62 for detailed definitions, footnotes and caveats. ^b Queensland does not provide acute services in psychiatric hospitals. ^c Tasmania, the ACT and the NT do not have psychiatric hospitals. ^d SA, the ACT and the NT do not have non-acute units in general hospitals.

Source: AIHW (unpublished) MHE NMDS; table 12A.62.

Older people's units have lower costs per inpatient day, but have considerably longer lengths of stay than general adult or child and adolescent units (figure 12.25). Data for forensic services are included for costs per inpatient bed day only, as the length of stay is dependent on factors outside the control of these services. Data on the average cost per inpatient bed day by target population for all care types are reported in table 12A.59.

Figure 12.25 **Inpatient care in acute units of public hospitals, by target population, 2013-14^{a, b, c, d, e}**



^a See box 12.16 and tables 12A.60–61 for detailed definitions, footnotes and caveats. ^b Queensland does not report any acute forensic services. ^c Tasmania does not provide, or cannot separately identify, child and adolescent mental health services or older people’s mental health services. ^d The ACT does not have separate forensic or child and adolescent mental health inpatient services. ^e The NT has general mental health services only.

Source: AIHW (unpublished) MHE NMDS; tables 12A.60–61.

Efficiency — cost of community-based residential care

‘Cost of community-based residential care’ is an indicator of governments’ objective that specialised mental health services be delivered in an efficient manner (box 12.17).

Box 12.17 Cost of community-based residential care

'Cost of community-based residential care' is defined as the average cost per patient day. Data are reported for both the care of adults and older people.

A low or decreasing average cost per patient day can indicate efficiency, although efficiency data need to be interpreted with care as they do not provide any information on the quality of service provided.

The indicator does not account for differences in the client mix. The client mix in community-based services can differ across jurisdictions and can reflect differences in the rate of institutional change (that is, the mainstreaming of mental health services).

Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions and over time
- complete (subject to caveats) for the current reporting period. All required 2013-14 data are available for all jurisdictions providing the services.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally in 2013-14, the average cost for 24 hour staffed residential care is higher for general adult units (\$535.58 per patient day) compared to older people's care units (\$397.86 per patient day), those this varied across states and territories (table 12A.63). Nationally and for all relevant jurisdictions, the costs for general adults units were higher for those staffed 24 hours compared to those that were non-24 hours staffed (table 12A.63).

Efficiency — cost of ambulatory care

'Cost of ambulatory care' is an indicator of governments' objective that specialised mental health services be delivered in an efficient manner (box 12.18).

Box 12.18 Cost of ambulatory care

'Cost of ambulatory care' is defined by two measures:

- average cost per treatment day
- average number of treatment days per episode — this measure is provided, along with average costs, as frequency of servicing is the main driver of variation in care costs.

An episode of ambulatory care is a three-month period of ambulatory care for an individual registered consumer where the consumer was under 'active care' (one or more treatment days in the period). Community-based periods relate to the following four fixed three-monthly periods: January to March, April to June, July to September, and October to December. Treatment day refers to any day on which one or more community contacts (direct or indirect) are recorded for a registered client during an ambulatory care episode.

(continued next page)

Box 12.18 (continued)

Low or decreasing average cost and/or fewer treatment days can indicate greater efficiency. Efficiency data need to be interpreted with care as they do not provide information on the quality of services provided.

The measures do not account for differences in the consumer mix. The consumer mix in community-based services can differ across jurisdictions and can reflect differences in the rate of institutional change (that is, the mainstreaming of mental health services) — for example, some State and Territory governments treat a higher proportion of consumers with more complex conditions in ambulatory care.


Data reported for the two measures are:

- comparable (subject to caveats) within some jurisdictions over time but are not comparable across jurisdictions or over time for other jurisdictions (see caveats in attachment tables for specific jurisdictions)
- complete (subject to caveats) for the current reporting period. All required data for 2013-14 are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally, the average recurrent cost per treatment day of ambulatory care declined slightly over the period 2009-10 to 2013-14 (figure 12.26), whereas the average treatment days per episode of ambulatory care increased slightly (figure 12.27).


Figure 12.26 Average recurrent cost per treatment day of ambulatory care (2013-14 dollars)^{a, b}



^a See box 12.18 and table 12A.64 for detailed definitions, footnotes and caveats. ^b Victorian 2011-12 and 2012-13 data are not available.

Source: AIHW (unpublished) Community Mental Health Care (CMHC) NMDS; AIHW (unpublished) MHE NMDS; table 12A.64.

Figure 12.27 **Average treatment days per episode of ambulatory care^{a, b}**



^a See box 12.18 and table 12A.64 for detailed caveats. ^b Victorian 2011-12 and 2012-13 data are not available.

Source: AIHW (unpublished) CMHC NMDS; AIHW (unpublished) MHE NMDS; table 12A.64.

Outcomes

Outcomes are the impact of services on the status of an individual or group (see chapter 1, section 1.5). The outcome indicators identified and/or reported here reflect the performance of governments (including the mental health sector) against the broad objectives of the NMHS. The whole-of-government approach within the *Fourth National Mental Health Plan 2009–2014* acknowledges that many of the determinants of good mental health, and of mental illness, are influenced by factors beyond the health system.

Rates of licit and illicit drug use

‘Rates of licit and illicit drug use’ is an indicator of governments’ objective to prevent the development of mental health problems and mental illness where possible, by reducing the prevalence of risk factors that contribute to the onset of mental illness and prevent longer term recovery (box 12.19).

Box 12.19 Rates of licit and illicit drug use

'Rates of licit and illicit drug use' is defined as the proportion of people aged 14 years or over who used alcohol at 'risky' levels and illicit drugs in the preceding 12 months. 'Risky' alcohol use is defined as more than two standard drinks per day on average. The specific illicit drugs include: cannabis, ecstasy, cocaine, meth/amphetamine, hallucinogens, Gamma hydroxybutyrate (GHB), inhalants, and heroin.

A low or decreasing proportion of people using alcohol at risky levels or using illicit drugs is desirable.

Many of the risk and protective factors that affect a person's propensity to consume these drugs lie outside the control of the mental health system. These include environmental, sociocultural and economic factors.


Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions but a break in series means that data for 2013 and 2010 are not comparable to data for earlier years
- complete (subject to caveats) for the current reporting period. All required 2013 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

In 2013, 18.2 per cent of people aged 14 years or over drank alcohol at levels considered 'risky' for developing long-term health problems (figure 12.28).

Figure 12.28 Use of alcohol in last 12 months by people aged 14 years or over, lifetime risk status, 2013^a



^a See box 12.19 and table 12A.65 for detailed definitions, footnotes and caveats.


Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra; table 12A.65.

The proportion of people who drank at these ‘risky’ levels slightly decreased between 2007 and 2013, a trend that was consistent across both males and females and most age groups (table 12A.71). Further data on alcohol use, and lifetime and single occasion risk status for 2013, 2010 and 2007, and by Indigenous status and remoteness areas are in tables 12A.66 and 12A.69–72.

Cannabis, ecstasy, cocaine and meth/amphetamines are the most widely used illicit drugs in Australia (table 12A.67). Data across the 2007, 2010 and 2013 surveys show that people using illicit drugs had higher levels of psychological distress and a higher proportion of people with a mental illness used illicit drugs than those without a mental illness (table 12A.74). Data by Indigenous status and remoteness areas are in table 12A.75.

Younger people’s use of cannabis and meth/amphetamines is of particular concern. Cannabis can precipitate schizophrenia in people who have a family history, increase the risk of psychosis symptoms and also exacerbate the schizophrenia symptoms (AHMC 2012). Psychosis symptoms are also associated with meth/amphetamine use and dependent meth/amphetamine users can also suffer from a range of co-morbid mental health problems (AHMC 2012). Since 2001, the proportions of younger people aged 14–19 years and 20–29 years who used these drugs decreased (figure 12.29). Cannabis use in 2013 by state and territory and age group are reported in table 12A.68.

Figure 12.29 Young people’s use of cannabis and meth/amphetamines, by age group^a



^a See box 12.19 and table 12A.73 for detailed definitions, footnotes and caveats.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra; table 12A.73.

Prevalence of mental illness

‘Prevalence of mental illness’ is an indicator of governments’ objective to prevent the development of mental health problems and mental illness where possible (box 12.20).

Box 12.20 Prevalence of mental illness

‘Prevalence of mental illness’ is defined as the proportion of the total population who have a mental illness. Proportions are reported for all people by sex, age group and disorder type.

A low or decreasing prevalence of mental illness can indicate that measures to prevent mental illness have been effective.

A reduction in the prevalence of mental illness can be brought about by preventative activities to stop an illness occurring, or by increasing access to effective treatments for those who have an illness (AHMC 2012). Many of the risk and protective factors that can affect the development of mental health problems and mental illness are outside the scope of the mental health system, in sectors that affect the daily lives of individuals and communities. These include environmental, sociocultural and economic factors, of which some can increase the risk of mental illness whilst others can support good mental health.

Not all mental illnesses are preventable and a reduction in the effect of symptoms and an improved quality of life will be a positive outcome for many people with a mental illness.

Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions (no time series data are reported)
- complete (subject to caveats) for the current reporting period. All required 2007 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally in 2007, the prevalence of a mental illness (with symptoms in the previous 12 months) was 20.0 per cent for people aged 16–85 years, with a further 25.5 per cent reported as having a mental illness at some point in their life (no symptoms in the last 12 months) (table 12A.76). National data on the prevalence of mental illness by disorder, age, sex and social characteristics are reported in tables 12A.76–78. These prevalence estimates are for the mental disorders that are considered to have the highest incidence rates in the population, but not some severe mental disorders, such as schizophrenia and bipolar disorder. The *National Survey of Psychotic Illness 2010* provides information on the one month treated prevalence of psychotic illnesses. In 2010, there were an estimated 3.1 cases of psychotic illness per 1000 adult population (aged 18–64 years) (Morgan et al. 2011).

The prevalence of mental illness among children and young people aged 4–17 years was an estimated 13.9 per cent in 2013–14 (Lawrence et al. 2015). Attention deficit/hyperactivity disorder (ADHD) was the most common mental illness overall for this age group, with 7.4 per cent assessed as having ADHD in the previous 12 months, followed by anxiety disorders (6.9 per cent) (Lawrence et al. 2015).

Mortality due to suicide

‘Mortality due to suicide’ is an indicator of governments’ objective to prevent mental health problems, mental illness and suicide, and identify and intervene early with people at risk (box 12.21).

Box 12.21 Mortality due to suicide

‘Mortality due to suicide’ is defined as the suicide rate per 100 000 people. The suicide rate is reported for all people by sex, age group, Indigenous status and significant urban areas. Deaths from suicide are defined as causes of death with the International Classification of Diseases (ICD)-10 codes X60–X84 and Y87.0.

A low or decreasing suicide rate per 100 000 people is desirable.

While mental health services contribute to reducing suicides, other government services also have a significant role. Public mental health programs are primarily concerned with providing treatment and support services for individual clients affected by severe mental illness, some of whom have either attempted, or indicated an intention, to commit suicide. Suicide prevention targeted at the wider population is also addressed through the initiatives of other government agencies, NGOs and other special interest groups.

Many factors outside the control of mental health services can influence a person’s decision to commit suicide. These include environmental, sociocultural and economic risk factors. Often a combination of these factors can increase the risk of suicidal behaviour.

Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions and over time for some years and disaggregations but not comparable for other years and disaggregations (see the DQI and attachment tables 12A.82–84 for details)
- complete (subject to caveats) for the current reporting period. All required 2013 or 2009–2013 data are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

People with a mental illness are at a higher risk of suicide than are the general population. For the period 2009–2013, 12 306 deaths by suicide were recorded in Australia (table 12A.81) — equivalent to 10.9 deaths per 100 000 people (figure 12.30). Nationally, the suicide rate increased as remoteness increased (ranging from 9.6 suicides per 100 000 people in capital cities to 14.8 suicides per 100 000 people in rural areas), with a similar pattern in most states and territories (table 12A.84).

Suicide data disaggregated by Indigenous status are available for NSW, Queensland, WA, SA and the NT only (figure 12.31). For the period 2009–2013, after adjusting for differences in population age structures, the rate of deaths for Aboriginal and Torres Strait Islander Australians due to suicide was almost twice the rate for non-Indigenous Australians (figure 12.31).


Figure 12.30 Suicide rates, 5 year average, 2009–2013^a



^a See box 12.21 and table 12A.82 for detailed definitions, footnotes and caveats.

Source: ABS (unpublished) *Causes of Deaths, Australia*, Cat. no. 3303.0; table 12A.82.

Figure 12.31 Suicide rates, by Indigenous status, 2009–2013^a



^a See box 12.21 and table 12A.85 for detailed definitions, footnotes and caveats.

Source: ABS (unpublished) *Causes of Deaths, Australia*, Cat. no. 3303.0; table 12A.85.

National data available by age and sex show that the rate for males (16.8 per 100 000 males) was over three times that for females (5.3 per 100 000 females) — a ratio that was

relatively constant over all age groups, except for those aged 85 years or over where the male suicide rate was around six times the female rate (table 12A.81).

Historical data are available by sex in table 12A.80 and by state and territory in tables 12A.82–84.

Physical health outcomes for people with a mental illness

‘Physical health outcomes for people with a mental illness’ is an indicator of governments’ objective to promote the recovery of people with a mental illness and to provide high quality co-ordinated services that are appropriate to the conditions and circumstances of people with a mental illness (box 12.22).

Box 12.22 Physical health outcomes for people with a mental illness

‘Physical health outcomes for people with a mental illness’ is defined by two measures, the proportion of adults with a mental illness (compared with the proportion of adults without a mental illness):

- who are exposed to particular health risk factors: obese/overweight, daily smokers and at risk of long term harm from alcohol.
- who experienced a long-term physical health condition: cancer, diabetes, arthritis, cardiovascular disease and asthma.

Low or decreasing proportions of people with a mental illness who are subject to particular health risk factors and who experience a long-term physical health condition are desirable.

People with a mental illness have worse physical health outcomes than people without mental illness (Coghlan et al. 2001; Happell et al. 2015; Joukamaa et al. 2001; Sartorius 2007; Lawrence, Hancock and Kisely 2013). However, the relationship between a physical and mental health is complex. Poor physical health can exacerbate mental health problems and poor mental health can lead to poor physical health. In addition, some psychiatric medications that are prescribed to treat mental health conditions are known to lead to worse physical health outcomes.

Data for these measures include 95 per cent confidence intervals (in the form of error bars in figures).

Data reported for this indicator are:


- comparable (subject to caveats) across jurisdictions (no time series data are reported)
- complete (subject to caveats) for the current reporting period. All required 2011-12 data are available for all jurisdictions.

The total and non-Indigenous components of the *2011-13 AHS* does not include people living in discrete Aboriginal and Torres Strait Islander communities and very remote areas, which affects the comparability of the NT results.

Data quality information for this indicator is under development.

Greater exposure to particular health risk factors can contribute to poorer physical health. In 2011-12, people with a mental illness had significantly higher daily smoking rates (26.1 per cent) compared to people without a mental illness (14.7 per cent) (figure 12.32). The proportions of people who are obese/overweight or at risk of long term harm from alcohol are similar for those with and without a mental illness (table 12A.86).

Figure 12.32 Adults who are daily smokers, by mental illness status, 2011-12^a




^a See box 12.22 and table 12A.86 for detailed definitions, footnotes and caveats.

Source: ABS (unpublished) *AHS 2011–13 (2011-12 NHS component)*, Cat. no. 4364.0; table 12A.86.

A higher proportion of people with a mental illness had long-term health conditions compared to people without mental illness. Nationally in 2011-12, the proportions of people with a mental illness who had arthritis (26.9 per cent) and cardiovascular disease (9.5 per cent) were higher than those without mental illness (16.7 per cent and 5.2 per cent respectively) (figure 12.33). Table 12A.87 provides data for cancer, asthma and diabetes.

Figure 12.33 **Adults with long-term health conditions, by mental illness status, 2011-12^a**



^a See box 12.22 and table 12A.87 for detailed definitions, footnotes and caveats.

Source: ABS (unpublished) *AHS 2011–13 (2011-12 NHS component)*, Cat. no. 4364.0; table 12A.87.

Social and economic inclusion of people with a mental illness

‘Social and economic inclusion of people with a mental illness’ is an indicator of governments’ objective to improve mental health and facilitate recovery from illness through encouraging meaningful participation in recreational, social, employment and other activities in the community (box 12.23).

Box 12.23 **Social and economic inclusion of people with a mental illness**

'Social and economic inclusion of people with a mental illness' is defined by three measures, the proportion of people:

- aged 16–64 years with a mental illness who are employed, compared with the proportion for people without a mental illness
- aged 16–30 years with a mental illness who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (studying full or part time), compared with the proportion for people without a mental illness
- aged 15 years or over with a mental illness who had face-to-face contact with family or friends living outside the household in the last week, compared with the proportion for people without a mental illness.

High or increasing proportions for people with a mental illness, and proportions that are similar to those without a mental illness, of people who are employed, who had face-to-face contact with family or friends and who are employed and/or are enrolled for study are desirable.

This indicator measures employment participation relative to the total population, as distinct from the labour force. Some people can choose not to participate in the labour force (that is, they are not working or actively looking for work). It also does not provide information on whether the employment, education or social activities participated in were appropriate or meaningful.

Data for these measures include 95 per cent confidence intervals (in the form of error bars in figures).

Data reported for this indicator are:

- comparable (subject to caveats) across jurisdictions for all surveys and over time for the 2011-12 and 2007-08 NHS data
- complete (subject to caveats) for the current reporting period. All required 2011-12 data and 2014 data are available for all jurisdictions.

The total and non-Indigenous components of the *2011-13 AHS* and the *2014 General Social Survey* does not include people living in discrete Aboriginal and Torres Strait Islander communities and very remote areas, which affects the comparability of the NT results.


Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

While employment is recognised as important in supporting good mental health, having a mental illness can act as a barrier to gaining and maintaining employment (AHMC 2012). Nationally in 2011-12, the proportion of people employed was lower for those with a mental illness (61.7 per cent) compared to those without a mental illness (80.3 per cent) (figure 12.34). The significantly higher proportion of people with a mental illness who do not participate in the labour force, compared to those without a mental illness, is a major contributing factor (32.0 per cent compared to 16.7 per cent). Historical data are available in tables 12A.90, 12A.92 and 12A.94.

Mental illness in early adult years can lead to disrupted and premature exit from education or disrupt the transition from school to work. These disruptions can have long term effects

on the person's ability to participate in a range of vocational activities (AHMC 2012). Nationally in 2011-12, the proportion of people aged 16–30 years with a mental illness who were employed and/or are enrolled for study in a formal secondary or tertiary qualification was 79.2 per cent, compared to 90.2 per cent for those without a mental illness (table 12A.89). Historical data are available in tables 12A.91 and 12A.93–94.

Figure 12.34 People aged 16–64 years who are employed, by mental illness status, 2011-12^a




^a See box 12.23 and table 12A.88 for detailed definitions, footnotes and caveats.

Source: ABS (unpublished) *AHS 2011–13 (2011-12 NHS component)*, Cat. no. 4364.0; table 12A.88.

Contact with immediate household, extended family, friends, and neighbours is related to lower levels of psychological distress and therefore can protect against the onset and adverse course of mental illnesses (Berry et al. 2007). Nationally in 2014, the proportion of people aged 15 years or over with a mental illness who had face-to-face contact with family or friends living outside the household in the last week (76.5 per cent) was similar to the proportion for people without a mental illness (77.1 per cent) (figure 12.35).

Figure 12.35 People who had face-to-face contact with family or friends living outside the household in the last week, by mental illness status, 2014^a



^a See box 12.23 and table 12A.95 for detailed definitions, footnotes and caveats.

Source: ABS (unpublished) *General Social Survey 2014*, Cat. no. 4159.0 (derived using Table Builder product); table 12A.95.

Mental health outcomes of consumers of specialised public mental health services

‘Mental health outcomes of consumers of specialised public mental health services’ is an indicator of governments’ objective to improve the effectiveness and quality of service delivery and outcomes and promote recovery from mental health problems and mental illness (box 12.24).

Box 12.24 Mental health outcomes of consumers of specialised public mental health services

‘Mental health outcomes of consumers of specialised public mental health services’ is defined as the proportion of people receiving care who had a significant improvement in their clinical mental health outcomes. See section 12.5 for information on how the consumer outcomes average score is derived. Data are also reported on the proportion who experienced no significant change or a significant deterioration in their mental health outcomes. Data are reported by three consumer types: people in ongoing community-based ambulatory care, people discharged from community-based ambulatory care and people discharged from a hospital psychiatric unit.

(continued next page)

Box 12.24 (continued)

A high or increasing proportion of people receiving care in State and Territory governments' specialised public mental health services who had a significant improvement in their clinical mental health outcomes is desirable.

This indicator has a number of technical and conceptual issues. The outcome measurement tool is imprecise. A single 'average score' does not reflect the complex service system in which services are delivered across multiple settings (inpatient, ambulatory and residential) and provided as both discrete, short term episodes of care and prolonged care over indefinite periods (AHMC 2012). The approach separates a consumer's care into segments (hospital versus the community) rather than tracking the person's overall outcomes across treatment settings. In addition, consumers' outcomes are measured from the clinician's perspective and not as the 'lived experience' from the consumer's viewpoint (AHMC 2012).


Data reported for this indicator:

- may not be comparable (subject to caveats) within jurisdictions over time and may not be comparable across jurisdictions due to differences in the quality of the data and the proportion of episodes for which completed outcomes data are available
- are complete (subject to caveats) for the current reporting period. All required data for 2013-14 are available for all jurisdictions.

Data quality information for this indicator is at www.pc.gov.au/rogs/2016.

Nationally in 2013-14, 26.6 per cent of people in ongoing ambulatory care, 48.4 per cent of people discharged from ambulatory care and 72.4 per cent of people discharged from a hospital psychiatric inpatient unit showed a significant improvement in their mental health clinical outcomes (figure 12.36). Across age groups and over the reporting years from 2007-08, for those discharged from hospital and community care, a larger proportion of people aged 18–64 years old showed a significant improvement compared to those in other age groups. Whereas, for those in ongoing community care, younger people aged 0–17 years had the highest proportion who showed a significant improvement compared to other age groups (table 12A.97).

Figure 12.36 **Mental health outcomes of consumers of State and Territory governments' specialised mental health services, 2013-14^a**



^a See box 12.24 and table 12A.96 for detailed definitions, footnotes and caveats. ^b The ACT and NT data are not published due to insufficient observations.

Source: AIHW (unpublished) from data provided by the Australian Mental Health Outcomes and Classification Network; table 12A.96.

12.4 Future directions in performance reporting

Priorities for future reporting on mental health management include the following:

- developing an estimate of the number of people who need mental health services so that access to services can be measured in terms of need
- improving reporting on government funded non-government entities to include information on their activity and the outcomes of the consumers of these services
- identifying indicators that relate to the performance framework dimension of sustainability
- further developing the measurement and reporting on the clinical mental health outcomes of consumers of specialised public mental health services.

12.5 Definitions of key terms

General terms

General practice

The organisational structure in which one or more GPs provide and supervise health care for a 'population' of patients. This definition includes medical practitioners who work solely with one specific population, such as women's health or Aboriginal and Torres Strait Islander health.

Health management

The ongoing process beginning with initial client contact and including all actions relating to the client. Includes assessment/evaluation, education of the person, family or carer(s), and diagnosis and treatment. Involves problems with adherence to treatment and liaison with, or referral to, other agencies.

Separation

An episode of care for an admitted patient, which can be a total hospital stay, or a portion of a hospital stay beginning or ending in a change of type of care (for example, from acute to rehabilitation). Separation also means the process by which an admitted patient completes an episode of care.

Mental health

Acute services

Services that primarily provide specialised psychiatric care for people with acute episodes of mental illness. These episodes are characterised by recent onset of severe clinical symptoms of mental illness that have potential for prolonged dysfunction or risk to self and/or others. The key characteristic of acute services is that the treatment effort focuses on symptom reduction with a reasonable expectation of substantial improvement. In general, acute psychiatric services provide relatively short term treatment. Acute services can:

- focus on assisting people who have had no prior contact or previous psychiatric history, or individuals with a continuing psychiatric illness for whom there has been an acute exacerbation of symptoms
- target the general population or be specialised in nature, targeting specific clinical populations. The latter group include psychogeriatric, child and adolescent, youth and forensic mental health services.

Accrued mental health patient days

Mental health care days are days of admitted patient care provided to admitted patients in psychiatric hospitals, designated psychiatric units and days of residential care provided to residents in residential mental health services. Accrued mental health care days can also be referred to as occupied bed days in specialised mental health services. The days to be counted are only those days occurring within the reference period, that is from 1 July to the

	<p>following 30 June for the relevant period, even if the patient/resident was admitted prior to the reference period or discharged after the reference period. The key basic rules to calculate the number of accrued mental health care days are as follows:</p> <ul style="list-style-type: none"> • For a patient admitted and discharged on different days, all days are counted as mental health care days except the day of discharge and any leave days. • Admission and discharge on the same day are equal to one patient day. • Leave days involving an overnight absence are not counted. • A patient day is recorded on the day of return from leave.
Affective disorders	A mood disturbance, including mania, hypomania, bipolar affective disorder, depression and dysthymia.
Ambulatory care services	Mental health services dedicated to the assessment, treatment, rehabilitation or care of non-admitted inpatients, including but not confined to crisis assessment and treatment services, mobile assessment and treatment services, outpatient clinic services (whether provided from a hospital or community mental health centre), child and adolescent outpatient treatment teams, social and living skills programs (including day programs, day hospitals and living skills centres), and psychogeriatric assessment teams and day programs.
Anxiety disorders	Feelings of tension, distress or nervousness. Includes agoraphobia, social phobia, panic disorder, generalised anxiety disorder, obsessive–compulsive disorder and post-traumatic stress disorder.
Average available beds	The number of beds available to provide overnight accommodation for patients admitted to hospital (other than neonatal cots [non-special-care] and beds occupied by hospital-in-the-home patients) or to specialised residential mental health care, averaged over the counting period. Beds are available only if they are suitably located and equipped to provide care and the necessary financial and human resources can be provided.
Child and adolescent mental health services	Services principally targeted at children and young people up to the age of 18 years. Classification of services in this category requires recognition by the regional or central funding authority of the special focus of the inpatient service on children or adolescents. These services can include a forensic component.
Community-based residential services	Staffed residential units established in community settings that provide specialised treatment, rehabilitation or care for people affected by a mental illness or psychiatric disability. To be defined as community-based residences, the services must: provide residential care to people with mental illnesses or psychiatric disability; be located in a community setting external to the campus of a general hospital or psychiatric institution; employ onsite staff for at least some part of the day; and be government funded.
Co-morbidity	The simultaneous occurrence of two or more illnesses such as depressive illness with anxiety disorder, or depressive disorder with anorexia.
Comparability	Data are considered comparable if (subject to caveats) they can be used to inform an assessment of comparative performance. Typically, data are considered comparable when they are collected in the same way and in accordance with the same definitions. For comparable indicators or measures, significant differences in reported results allow an assessment of differences in performance, rather than being the result of anomalies in the data.
Completeness	Data are considered complete if all required data are available for all jurisdictions that provide the service.
Cost per inpatient bed day	The average patient day cost according to the inpatient type.
Depression	A state of gloom, despondency or sadness lasting at least two weeks. The person usually suffers from low mood, loss of interest and enjoyment, and reduced energy. Sleep, appetite and concentration can be affected.
Forensic mental health services	Services principally providing assessment, treatment and care of mentally ill individuals whose behaviour has led them to commit criminal offences or makes it likely that they will offend in the future if not adequately treated and contained. This includes prison-based services, but excludes services that are primarily for children and adolescents and for older people even where they

	include a forensic component.
General mental health services	<p>Services that principally target the general adult population (18–65 years old) but that can provide services to children, adolescents or older people. Includes, therefore, those services that cannot be described as specialised child and adolescent, youth, older people's or forensic services.</p> <p>General mental health services include hospital units whose principal function is to provide some form of specialised service to the general adult population (for example, inpatient psychotherapy) or to focus on specific clinical disorders within the adult population (for example, postnatal depression, anxiety disorders).</p>
Mental illness	A diagnosable illness that significantly interferes with an individual's cognitive, emotional and/or social abilities.
Mental health	The capacity of individuals within groups and the environment to interact with one another in ways that promote subjective wellbeing, the optimal development and use of mental abilities (cognitive, affective and relational) and the achievement of individual and collective goals consistent with justice.
Mental health problems	Diminished cognitive, emotional or social abilities, but not to the extent of meeting the criteria for a mental illness.
Mental health promotion	Actions taken to maximise mental health and wellbeing among populations and individuals. It is aimed at changing environments (social, physical, economic, educational, cultural) and enhancing the 'coping' capacity of communities, families and individuals by giving power, knowledge, skills and necessary resources.
Mental illness prevention	Interventions that occur before the initial onset of an illness to prevent its development. The goal of prevention interventions is to reduce the incidence and prevalence of mental health problems and mental illnesses.
Non-acute services	<p>Non-acute services are defined by two categories:</p> <ul style="list-style-type: none"> • Rehabilitation services that have a primary focus on intervention to reduce functional impairments that limit the independence of patients. Rehabilitation services are focused on disability and the promotion of personal recovery. They are characterised by an expectation of substantial improvement over the short to mid-term. Patients treated by rehabilitation services usually have a relatively stable pattern of clinical symptoms. • Extended care services that primarily provide care over an indefinite period for patients who have a stable but severe level of functional impairment and an inability to function independently, thus requiring extensive care and support. Patients of extended care services present a stable pattern of clinical symptoms, which can include high levels of severe unremitting symptoms of mental illness. Treatment is focused on preventing deterioration and reducing impairment; improvement is expected to occur slowly.
Non-government organisations (NGOs)	Private not-for-profit community managed organisations that receive State and Territory government funding specifically for the purpose of providing community support services for people affected by a mental illness or psychiatric disability. Programs provided by the NGOs sector can include supported accommodation services (including community-based crisis and respite beds), vocational rehabilitation programs, advocacy programs (including system advocacy), consumer self-help services, and support services for families and primary carers.
Older people's mental health services	Services principally targeting people in the age group 65 years or over. Classification of services in this category requires recognition by the regional or central funding authority of the special focus of the inpatient service on aged people. These services can include a forensic component. Excludes general mental health services that may treat older people as part of a more general service.
Outpatient services — community-based	Services primarily provided to non-admitted patients on an appointment basis and delivered from health centres located in community settings, physically separated within hospital sites. They can include outreach or domiciliary care as an adjunct to services provided from the centre base.
Outpatient services — hospital-based	Services primarily provided to non-admitted patients on an appointment basis and delivered from clinics located within hospitals. They can include outreach or domiciliary care as an adjunct to services provided from the clinic base.

Outcomes measurement – calculating the consumers ‘score’.	The assessment of a consumer’s clinical mental health outcomes is based on the changes reported in a consumer’s ‘score’ on a rating scale known as the Health of the Nation Outcomes Scale (HoNOS), or for children and adolescents, the Health of the Nation Outcome Scales for Children and Adolescents (HoNOSCA). Outcome scores are classified based on effect size — a statistic used to assess the magnitude of a treatment effect (AHMC 2012). The effect size is based on the ratio of the difference between the pre- and post-scores to the standard deviation of the pre-score. Individual episodes are classified as ‘significant improvement’ if the effect size index is greater than or equal to positive 0.5; ‘no change’ if the index is between 0.5 and -0.5; and ‘significant deterioration’ if the effect size index is less than or equal to -0.5 (AHMC 2012)
Prevalence	The number of cases of a disease present in a population at a given time (point prevalence) or during a given period (period prevalence).
Preventive interventions	Programs designed to decrease the incidence, prevalence and negative outcomes of illnesses.
Psychiatrist	A medical practitioner with specialist training in psychiatry.
Public health	The organised, social response to protect and promote health, and to prevent illness, injury and disability. The starting point for identifying public health issues, problems and priorities, and for designing and implementing interventions, is the population as a whole or population subgroups. Public health is characterised by a focus on the health of the population (and particular at-risk groups) and complements clinical provision of health care services.
Public (non-psychiatric) hospital	A hospital that provides at least minimum medical, surgical or obstetric services for inpatient treatment and/or care, and around-the-clock, comprehensive, qualified nursing services, as well as other necessary professional services.
Schizophrenia	A combination of signs and symptoms that can include delusions, hallucinations, disorganised speech or behaviour, a flattening in emotions, and restrictions in thought, speech and goal directed behaviour.
Seclusion	<p>Seclusion is the confinement of the consumer at any time of the day or night alone in a room or area from which free exit is prevented. The intended purpose of the confinement is not relevant in determining what is or is not seclusion. Seclusion applies even if the consumer agrees or requests the confinement (NMHPSC 2011b).</p> <p>The awareness of the consumer that they are confined alone and denied exit is not relevant in determining what is or is not seclusion. The structure and dimensions of the area to which the consumer is confined is not relevant in determining what is or is not seclusion. The area may be an open area, for example, a courtyard. Seclusion does not include confinement of consumers to High Dependency sections of gazetted mental health units, unless it meets the definition (AIHW 2015).</p>
Seclusion event	An event is when a consumer enters seclusion and when there is a clinical decision to cease seclusion. Following the clinical decision to cease seclusion, if a consumer re-enters seclusion within a short period of time this would be considered a new seclusion event. The term ‘seclusion event’ is utilised to differentiate it from the different definitions of ‘seclusion episode’ used across jurisdictions (NMHPSC 2011b).
Specialised mental health inpatient services	Services provided to admitted patients in stand-alone psychiatric hospitals or specialised psychiatric units located within general hospitals.
Specialised mental health services	Services whose primary function is specifically to provide treatment, rehabilitation or community support targeted towards people affected by a mental illness or psychiatric disability. Further, such activities are delivered from a service or facility that is readily identifiable as both specialised and serving a mental health function. This criterion applies regardless of the source of funds.
Specialised residential services	Services provided in the community that are staffed by mental health professionals on a non-24 or 24-hour basis.
Staffing categories (mental health)	Medical officers: all medical officers employed or engaged by the organisation on a full time or part time basis. Includes visiting medical officers who are engaged on an hourly, sessional or fee-for-service basis.

Psychiatric hospitals

Psychiatrists and consultant psychiatrists: medical officers who are registered to practice psychiatry under the relevant State or Territory medical registration board; or who are fellows of the Royal Australian and New Zealand College of Psychiatrists or registered with Health Insurance Commission as a specialist in Psychiatry.

Psychiatry registrars and trainees: medical officers who are formal trainees within the Royal Australian and New Zealand College of Psychiatrists' Postgraduate Training Program.

Other medical officers: medical officers employed or engaged by the organisation who are not registered as psychiatrists within the State or Territory, or as formal trainees within the Royal Australian and New Zealand College of Psychiatrists' Postgraduate Training Program.

Nursing staff: all categories of registered nurses and enrolled nurses, employed or engaged by the organisation.

Registered nurses: people with at least a three year training certificate or tertiary qualification who are certified as being a registered nurse with the State or Territory registration board. This is a comprehensive category and includes general and specialised categories of registered nurses.

Enrolled nurses: refers to people who are second level nurses who are enrolled in all states except Victoria where they are registered by the state registration board to practise in this capacity. Includes general enrolled nurse and specialist enrolled nurse (e.g. mothercraft nurses in some states).

Diagnostic and health professionals (allied health professionals): qualified staff (other than qualified medical or nursing staff) who are engaged in duties of a diagnostic, professional or technical nature. This category covers all allied health professionals, such as social workers, psychologists, occupational therapists, physiotherapists, and other diagnostic and health professionals.

Social workers: people who have completed a course of recognised training and are eligible for membership of the Australian Association of Social Workers.

Psychologists: people who are registered as psychologists with the relevant State or Territory registration board.

Occupational therapists: people who have completed a course of recognised training and who are eligible for membership of the Australian Association of Occupational Therapists.

Other personal care staff: attendants, assistants, home companions, family aides, ward helpers, warders, orderlies, ward assistants and nursing assistants who are engaged primarily in the provision of personal care to patients or residents, and who are not formally qualified or who are undergoing training in nursing or allied health professions.

Administrative and clerical staff: staff engaged in administrative and clerical duties. Excludes medical, nursing, diagnostic and health professional and domestic staff wholly or partly involved in administrative and clerical duties, who should be counted under their appropriate occupational categories. Civil engineers and computing staff are included in this category.

Domestic and other staff: staff involved in the provision of food and cleaning services including domestic staff primarily engaged in administrative duties such as food services manager. Dieticians are excluded.

Health establishments that are primarily devoted to the treatment and care of inpatients with psychiatric, mental or behavioural disorders, and that are situated at physically separate locations from a general hospital. Stand-alone hospitals may or may not be managed by the mainstream health system. Psychiatric hospitals situated at physically separate locations from a general hospital are included within the 'stand-alone' category regardless of whether they are under the management control of a general hospital. A health establishment that operates in a separate building but is located on, or immediately adjoining, the acute care hospital campus can also be a stand-alone hospital if the following criteria are not met:

- a single organisational or management structure covers the acute care hospital and the psychiatric hospital
- a single employer covers the staff of the acute care hospital and the psychiatric hospital

	<ul style="list-style-type: none"> • the location of the acute care hospital and psychiatric hospital can be regarded as part of a single overall hospital campus • the patients of the psychiatric hospital are regarded as patients of the single integrated health service.
Substance use disorders	Disorders in which drugs or alcohol are used to such an extent that behaviour becomes maladaptive, social and occupational functioning is impaired, and control or abstinence becomes impossible. Reliance on the drug can be psychological (as in substance misuse) or physiological (as in substance dependence).
Youth mental health services	Services principally targeting children and young people generally aged 16-25 years. The classification of a service into this category requires recognition by the regional or central funding authority of the special focus of the service. These services may include a forensic component.

12.6 List of attachment tables

Attachment tables are identified in references throughout this chapter by a ‘12A’ prefix (for example, table 12A.1). Attachment tables are available on the website (www.pc.gov.au/rogs/2016).

Table 12A.1	Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million)
Table 12A.2	Real estimated recurrent expenditure on State and Territory governments specialised mental health services (2013-14 dollars)
Table 12A.3	Real estimated expenditure on State and Territory governments’ specialised mental health services, by funding source (2013-14 dollars) (\$million)
Table 12A.4	Real Australian, State and Territory governments expenditure on mental health services (2013-14 dollars) (\$million),
Table 12A.5	Depreciation expenditure on State and Territory governments’ specialised mental health services (current prices) (\$million)
Table 12A.6	Total state and territory recurrent expenditure on specialised mental health services (current prices)
Table 12A.7	Functioning and quality of life measures, by 12-month mental disorder status, 2007 (per cent)
Table 12A.8	Age-standardised rate of adults with very high levels of psychological distress, by State and Territory, 2011-12
Table 12A.9	Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2011-12
Table 12A.10	Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2011-12
Table 12A.11	Age-standardised rate of adults with very high levels of psychological distress, by State and Territory, 2007-08
Table 12A.12	Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2007-08

Table 12A.13	Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2007-08
Table 12A.14	Level of psychological distress K10, 2007-08 (per cent)
Table 12A.15	Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, by Indigenous status, 2011–13
Table 12A.16	Level of psychological distress K10, 2004-05 (per cent)
Table 12A.17	Mental health care specific MBS items processed
Table 12A.18	GP mental health-related encounters (general and mental health specific)
Table 12A.19	GP mental health-related encounters (general and mental health specific), by patient demographics, 2013-14
Table 12A.20	The 10 most frequent GP managed mental health-related problems, by gender, 2013-14
Table 12A.21	Mental health patient days
Table 12A.22	Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type
Table 12A.23	Ambulatory-equivalent public mental health-related separations with specialised psychiatric care, by principal diagnosis, 2012-13
Table 12A.24	Community mental health service contacts, by sex and age group
Table 12A.25	Specialised mental health care reported, by Indigenous status
Table 12A.26	Available beds in State and Territory governments' specialised mental health services
Table 12A.27	Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people)
Table 12A.28	FTE direct care staff employed in specialised mental health services, by service setting (per 100 000 people)
Table 12A.29	Targeted Community Care (Mental Health) program participants (number)
Table 12A.30	Admitted patient mental health-related separations without specialised psychiatric care, by principal diagnosis in ICD-10-AM groupings, 2012-13
Table 12A.31	Ambulatory-equivalent mental health-related separations without specialised psychiatric care, by principal diagnosis, 2012-13
Table 12A.32	Mental health-related emergency department occasions of service in public hospitals, by episode end status, 2012-13
Table 12A.33	New clients as a proportion of total clients under the care of State or Territory specialised public mental health services
Table 12A.34	New clients as a proportion of total clients under the care of State or Territory specialised public mental health services, by selected characteristics, 2013-14
Table 12A.35	New clients as a proportion of total clients of MBS subsidised mental health services
Table 12A.36	New clients as a proportion of total clients under the care of MBS subsidised mental health services, by selected characteristics, 2014-15
Table 12A.37	Proportion of people receiving clinical mental health services by service type and Indigenous status

Table 12A.38	Proportion of people receiving clinical mental health services by service type and remoteness area
Table 12A.39	Proportion of people receiving clinical mental health services by service type and SEIFA
Table 12A.40	People receiving clinical public mental health services, by age group and gender, 2013-14
Table 12A.41	Proportion of people receiving clinical mental health services, by service type and SEIFA IRSD deciles (age-standardised rate)
Table 12A.42	Proportion of people receiving clinical mental health services by service type
Table 12A.43	Services used for mental health problems, Australia, 2007 (per cent)
Table 12A.44	Services used for mental health, by mental disorder status, 2007 (per cent)
Table 12A.45	Young people who had contact with MBS subsidised primary mental health care services, by age group
Table 12A.46	Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by selected characteristics (per cent)
Table 12A.47	Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by service type (per cent)
Table 12A.48	Specialised public mental health services reviewed against National Standards for Mental Health Services, 30 June
Table 12A.49	Recurrent expenditure on community-based services as a proportion of total spending on mental health services (per cent)
Table 12A.50	Specialised public mental health services episodes with completed consumer outcomes measures collected
Table 12A.51	Rate and duration of seclusion events in public specialised mental health acute inpatient units
Table 12A.52	Rate and duration of seclusion events in public specialised mental health acute inpatient units (per 1000 patient days), by target population
Table 12A.53	Consumer and carer participation
Table 12A.54	Rates of community follow-up for people within the first seven days of discharge from hospital
Table 12A.55	Rates of community follow-up within first seven days of discharge from a psychiatric admission, by State and Territory, by Indigenous status and remoteness
Table 12A.56	Rates of community follow-up within first seven days of discharge from a psychiatric admission, by age group, gender and SEIFA quintiles, 2013-14
Table 12A.57	Readmissions to hospital within 28 days of discharge
Table 12A.58	Readmissions to hospital within 28 days of discharge, by selected characteristics, 2013-14
Table 12A.59	Average recurrent real costs per inpatient bed day, public hospitals, by target population (2013-14 dollars)
Table 12A.60	Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars)

Table 12A.61	Average length of stay, public hospitals acute units, by target population (no. of days)
Table 12A.62	Average recurrent cost per inpatient bed day, by public hospital type (2013-14 dollars)
Table 12A.63	Average recurrent cost per patient day for community residential services (2013-14 dollars)
Table 12A.64	Average cost, and treatment days per episode, of ambulatory care
Table 12A.65	Risk status recent drinkers (in last 12 months) aged 14 years or over, 2013 (per cent)
Table 12A.66	Recent drinkers lifetime and single occasion risk, people aged 14 years or older, by social characteristics, 2013 (per cent)
Table 12A.67	Recent alcohol and illicit drug use, people aged 14 years or over, by substance, 2013 (per cent)
Table 12A.68	Use of cannabis, by age group, 2013 (per cent)
Table 12A.69	Risk status recent drinkers (in last 12 months) aged 14 years or over, 2010 (per cent)
Table 12A.70	Recent alcohol and illicit drug use, people aged 14 years or over, by substance, 2010 (per cent)
Table 12A.71	Lifetime risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent)
Table 12A.72	Single occasion risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent)
Table 12A.73	Selected illicit drug use, by substance and age group (per cent)
Table 12A.74	Selected illicit drug use by people aged 18 years or over, by level of psychological distress and self-reported health conditions (per cent)
Table 12A.75	Illicit drug use, people aged 14 years or older, by social characteristics (per cent)
Table 12A.76	Prevalence of lifetime mental disorders among adults aged 16–85 years, 2007 (per cent)
Table 12A.77	Prevalence of lifetime mental disorders among adults aged 16–85 years, by sex, 2007 (per cent)
Table 12A.78	Prevalence of lifetime mental disorders among adults, by age, 2007 (per cent)
Table 12A.79	Prevalence of lifetime mental disorders among adults, by disadvantage and section of state, 2007 (per cent)
Table 12A.80	Suicides and mortality rate, by sex, Australia
Table 12A.81	Suicides and mortality rate, by age and sex, Australia
Table 12A.82	Suicide deaths and death rate
Table 12A.83	Suicide deaths and death rate of people aged 15–24 years
Table 12A.84	Suicide deaths and suicide death rate, by area
Table 12A.85	Suicide deaths, by Indigenous status, 2009–2013
Table 12A.86	Age-standardised proportions of adults by health risk factors and mental illness status, 2011-12

-
- Table 12A.87** Age-standardised proportions of adults by long-term health conditions and mental illness status, 2011-12
- Table 12A.88** Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2011-12 (per cent)
- Table 12A.89** Age-standardised proportion of the population aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status, 2011-12 (per cent)
- Table 12A.90** Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2007-08 (per cent)
- Table 12A.91** Population aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status, 2007-08 (per cent)
- Table 12A.92** Labour force and employment participation among adults aged 16–64 years, by mental disorder status, 2007 (per cent)
- Table 12A.93** Education, training and employment participation among adults aged 16–30 years, by mental disorder status, 2007 (per cent)
- Table 12A.94** Labour force and employment participation among adults aged 16–30 years, by mental disorder status, 2007 (per cent)
- Table 12A.95** Proportion of people who had face-to-face contact with family or friends living outside the household in the last week, by mental illness status, 2014 (per cent)
- Table 12A.96** Clinical outcomes of people receiving various types of mental health care provided by State and Territory public mental health services (per cent)
- Table 12A.97** People who received mental health care provided by State and Territory public mental health services and who significantly improved, by service type and age group (per cent)
- Table 12A.98** Deflators used to calculate real State and Territory mental health expenditure
- Table 12A.99** Estimated resident populations used in mental health per head calculations

12.7 References

ABS (Australian Bureau of Statistics) 2012, *Information Paper: Use of the Kessler Psychological Distress Scale in ABS Health Surveys*, Australia, 2007-08, Cat. no. 4817.0.55.001, Canberra.

Australian Health Ministers 1991, *Mental Health Statement of Rights and Responsibilities*, Australian Government Publishing Service, Canberra.

AHMC (Australian Health Ministers' Conference) 2008, *Council of Australian Governments National Action Plan for Mental Health 2006–2011: Progress Report 2006-07*, Report prepared under the auspice of the Mental Health Standing Committee of the Australian Health Ministers' Advisory Council, www.coag.gov.au/node/195 (accessed 16 November 2012).

— 2010, *National Standards for Mental Health Services 2010*, Canberra.

-
- 2012, *Council of Australian Governments National Action Plan for Mental Health 2006–2011: Progress Report 2009–10*, Report prepared under the auspice of the Mental Health Standing Committee of the Australian Health Ministers' Advisory Council, www.coag.gov.au/node/441 (accessed 16 November 2012).
- AIHW (Australian Institute of Health and Welfare) 2014, *Mental Health Services in Australia Online*, mhsa.aihw.gov.au/home/ (accessed 1 October 2014 and 17 December 2014).
- 2015, *Mental Health Services in Australia Online*, mhsa.aihw.gov.au/home/ (accessed 28 September 2015).
- Begg, S., Vos, T., Barker, B., Stevenson, C., Stanley, L. and Lopez, A. 2007, *The burden of disease and injury in Australia, 2003*, Cat. no. PHE 82, AIHW, Canberra.
- Berry, H., Rodgers, B. and Dear, K. 2007, 'Preliminary development and validation of an Australian community participation questionnaire: Types of participation and associations with distress in a coastal community', *Social Science & Medicine*, vol. 64, pp. 1719–1737.
- Coghlan, R., Lawrence D., Holman D. and Jablensky A. 2001, *Duty to Care: Physical Illness in People with Mental Illness*, University of Western Australia, Perth.
- Department of Health 2014, *National mental health strategy* (available at www.health.gov.au/internet/main/publishing.nsf/content/mental-strat).
- DoHA (Department of Health and Ageing) 2010, *National Mental Health Report 2010: Summary of 15 years of reform in Australia's Mental Health Services under the National Mental Health Strategy 1993–2008*, Australian Government, Canberra.
- DHAC (Australian Government Department of Health and Community Services) and AIHW 1999, *National Health Priority Areas Report: Mental Health 1998*, AIHW Cat. no. PHE 13, Canberra.
- Happell, B., Platania-Phung, C., Webster, S., McKenna, B., Millar, F., Stanton, R., Galletly, C., Castle, D., Furness, T., Liu, D. and Scott, D 2015, 'Applying the World Health Organization Mental Health Action Plan to evaluate policy on addressing co-occurrence of physical and mental illnesses in Australia', *Australian Health Review*, Vol. 39, pp. 370–378, CSRIO Publishing (available at <http://dx.doi.org/10.1071/AH14098>).
- Joukamaa, M., Heliövaara, M., Knekt, P., Aromaa, A., Raitasalo, R. and Lehtinen, V. 2001, 'Mental disorders and cause specific mortality', *The British Journal of Psychiatry*, vol. 179, no. 6, pp. 498–502.
- Lawrence, D., Hancock, K. and Kisely S. 2013, 'The gap in life expectancy from preventable physical illness in psychiatric patients in Western Australia: retrospective analysis of population based registers', *British Medical Journal*, Vol. 346.
- Lawrence, D., Johnson, S., Hafekost, J., Boterhoven, K., Sawyer, M., Ainley, J., Zubrick, S. 2015, *The Mental Health of Children and Adolescents. Report on the*

second Australian Child and Adolescent Survey of Mental Health and Wellbeing, Department of Health, Canberra.

Morgan, V., Waterreus, A., Jablensky, A., Mackinnon, A., McGrath, J., Carr, V., Bush, R., Castle, D., Cohen, M., Harvey, C., Galletly, C., Stain, H., Neil, A. McGorry, P., Hocking, B., Shah, S., Saw, S. 2011, *People Living With Psychotic Illness 2010, Report on the Second Australian National Survey*, Australian Government, Canberra.

NMHPSC (National Mental Health Performance Subcommittee) 2011a, *The Fourth National Mental Health Plan Measurement Strategy 2011*, Report produced for the Australian Health Ministers Advisory Council Mental Health Standing Committee, www.health.gov.au/internet/mhsc/publishing.nsf/Content/publications_1 (accessed 30 September 2011).

— 2011b, *Key performance indicators for Australian public mental health services: second edition 2011*, Report produced for the Australian Health Ministers Advisory Council Mental Health Standing Committee, [www.health.gov.au/internet/mhsc/Australian Health Ministers Advisory Council Mental Health Standing Committee publishing.nsf/Content/publications_1](http://www.health.gov.au/internet/mhsc/Australian%20Health%20Ministers%20Advisory%20Council%20Mental%20Health%20Standing%20Committee%20publishing.nsf/Content/publications_1) (accessed 6 October 2013).

National Mental Health Working Group Information Strategy Committee Performance Indicator Drafting Group 2005, *Key Performance Indicators for Australian Public Mental Health Services*, ISC Discussion Paper no. 6, Australian Government Department of Health and Ageing, Canberra.

Sartorius, N. 2007, 'Physical illness in people with mental disorders', *World Psychiatry*, vol. 6, no. 1, pp. 3–4.

WHO (World Health Organization) 2001, *Strengthening mental health promotion*, Fact sheet no. 220, Geneva.

12A Mental health management — attachment

Definitions for the indicators and descriptors in this attachment are in section 12.5 of the chapter. Unsourced information was obtained from the Australian, State and Territory governments.

Data in this Report are examined by the Health Working Group, but have not been formally audited by the Secretariat.

Data reported in the attachment tables are the most accurate available at the time of data collection. Historical data may have been updated since the last edition of RoGS.

This file is available in Adobe PDF format on the Review web page (www.pc.gov.au/rogs/2016).

Attachment contents

Table 12A.1	Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million)
Table 12A.2	Real estimated recurrent expenditure on State and Territory governments specialised mental health services (2013-14 dollars)
Table 12A.3	Real estimated expenditure on State and Territory governments' specialised mental health services, by funding source (2013-14 dollars) (\$million)
Table 12A.4	Real Australian, State and Territory governments expenditure on mental health services (2013-14 dollars) (\$million),
Table 12A.5	Depreciation expenditure on State and Territory governments' specialised mental health services (current prices) (\$million)
Table 12A.6	Total state and territory recurrent expenditure on specialised mental health services (current prices)
Table 12A.7	Functioning and quality of life measures, by 12-month mental disorder status, 2007 (per cent)
Table 12A.8	Age-standardised rate of adults with very high levels of psychological distress, by State and Territory, 2011-12
Table 12A.9	Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2011-12
Table 12A.10	Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2011-12
Table 12A.11	Age-standardised rate of adults with very high levels of psychological distress, by State and Territory, 2007-08
Table 12A.12	Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2007-08
Table 12A.13	Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2007-08
Table 12A.14	Level of psychological distress K10, 2007-08 (per cent)
Table 12A.15	Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, by Indigenous status, 2011-13
Table 12A.16	Level of psychological distress K10, 2004-05 (per cent)
Table 12A.17	Mental health care specific MBS items processed
Table 12A.18	GP mental health-related encounters (general and mental health specific)
Table 12A.19	GP mental health-related encounters (general and mental health specific), by patient demographics, 2013-14
Table 12A.20	The 10 most frequent GP managed mental health-related problems, by gender, 2013-14
Table 12A.21	Mental health patient days
Table 12A.22	Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type
Table 12A.23	Ambulatory-equivalent public mental health-related separations with specialised psychiatric care, by principal diagnosis, 2012-13
Table 12A.24	Community mental health service contacts, by sex and age group
Table 12A.25	Specialised mental health care reported, by Indigenous status
Table 12A.26	Available beds in State and Territory governments' specialised mental health services
Table 12A.27	Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people)

Attachment contents

Table 12A.28	Full time equivalent (FTE) direct care staff employed in specialised mental health services, by service setting (per 100 000 people)
Table 12A.29	Targeted Community Care (Mental Health) program participants (number)
Table 12A.30	Admitted patient mental health-related separations without specialised psychiatric care, by principal diagnosis in ICD-10-AM groupings, 2012-13
Table 12A.31	Ambulatory-equivalent mental health-related separations without specialised psychiatric care, by principal diagnosis, 2012-13
Table 12A.32	Mental health-related emergency department occasions of service in public hospitals, by episode end status, 2012-13
Table 12A.33	New clients as a proportion of total clients under the care of State or Territory specialised public mental health services
Table 12A.34	New clients as a proportion of total clients under the care of State or Territory specialised public mental health services, by selected characteristics, 2013-14
Table 12A.35	New clients as a proportion of total clients of MBS subsidised mental health services
Table 12A.36	New clients as a proportion of total clients under the care of MBS subsidised mental health services, by selected characteristics, 2013-14
Table 12A.37	Proportion of people receiving clinical mental health services by service type and Indigenous status
Table 12A.38	Proportion of people receiving clinical mental health services by service type and remoteness area
Table 12A.39	Proportion of people receiving clinical mental health services by service type and SEIFA
Table 12A.40	People receiving clinical public mental health services, by age group and gender, 2013-14
Table 12A.41	Proportion of people receiving clinical mental health services, by service type and SEIFA IRSD deciles (age-standardised rate)
Table 12A.42	Proportion of people receiving clinical mental health services by service type
Table 12A.43	Services used for mental health problems, Australia, 2007 (per cent)
Table 12A.44	Services used for mental health, by mental disorder status, 2007 (per cent)
Table 12A.45	Young people who had contact with MBS subsidised primary mental health care services, by age group
Table 12A.46	Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by selected characteristics (per cent)
Table 12A.47	Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by service type (per cent)
Table 12A.48	Specialised public mental health services reviewed against National Standards for Mental Health Services, 30 June
Table 12A.49	Recurrent expenditure on community-based services as a proportion of total spending on mental health services (per cent)
Table 12A.50	Specialised public mental health services episodes with completed consumer outcomes measures collected
Table 12A.51	Rate and duration of seclusion events in public specialised mental health acute inpatient units
Table 12A.52	Rate and duration of seclusion events in public specialised mental health acute inpatient units (per 1000 patient days), by target population
Table 12A.53	Consumer and carer participation

Attachment contents

Table 12A.54	Rates of community follow-up for people within the first seven days of discharge from hospital
Table 12A.55	Rate of community follow up within first seven days of discharge from a psychiatric admission, by State and Territory, by Indigenous status and remoteness
Table 12A.56	Rate of community follow up within first seven days of discharge from a psychiatric admission, by age group, gender and SEIFA quintiles, 2013-14
Table 12A.57	Readmissions to hospital within 28 days of discharge
Table 12A.58	Readmissions to hospital within 28 days of discharge, by selected characteristics, 2013-14
Table 12A.59	Average recurrent real costs per inpatient bed day, public hospitals, by target population (2013-14 dollars)
Table 12A.60	Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars)
Table 12A.61	Average length of stay, public hospitals acute units, by target population (no. of days)
Table 12A.62	Average recurrent cost per inpatient bed day, by public hospital type (2013-14 dollars)
Table 12A.63	Average recurrent cost per patient day for community residential services (2013-14 dollars)
Table 12A.64	Average cost, and treatment days per episode, of ambulatory care
Table 12A.65	Risk status recent drinkers (in last 12 months) aged 14 years or over, 2013 (per cent)
Table 12A.66	Recent drinkers lifetime and single occasion risk, people aged 14 years or older, by social characteristics, 2013 (per cent)
Table 12A.67	Recent alcohol and illicit drug use, people aged 14 years or over, by substance, 2013 (per cent)
Table 12A.68	Use of cannabis, by age group, 2013 (per cent)
Table 12A.69	Risk status recent drinkers (in last 12 months) aged 14 years or over, 2010 (per cent)
Table 12A.70	Recent alcohol and illicit drug use, people aged 14 years or over, by substance, 2010 (per cent)
Table 12A.71	Lifetime risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent)
Table 12A.72	Single occasion risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent)
Table 12A.73	Selected illicit drug use, by substance and age group (per cent)
Table 12A.74	Selected illicit drug use by people aged 18 years or over, by level of psychological distress and self-reported health conditions (per cent)
Table 12A.75	Illicit drug use, people aged 14 years or older, by social characteristics (per cent)
Table 12A.76	Prevalence of lifetime mental disorders among adults aged 16–85 years, 2007 (per cent)
Table 12A.77	Prevalence of lifetime mental disorders among adults aged 16–85 years, by sex, 2007 (per cent)
Table 12A.78	Prevalence of lifetime mental disorders among adults, by age, 2007 (per cent)
Table 12A.79	Prevalence of lifetime mental disorders among adults, by disadvantage and section of state, 2007 (per cent)
Table 12A.80	Suicides and mortality rate, by sex, Australia
Table 12A.81	Suicides and mortality rate, by age and sex, Australia
Table 12A.82	Suicide deaths and death rate

Attachment contents

Table 12A.83	Suicide deaths and death rate of people aged 15–24 years
Table 12A.84	Suicide deaths and suicide death rate, by area
Table 12A.85	Suicide deaths, by Indigenous status, 2009–2013
Table 12A.86	Age-standardised proportions of adults by health risk factors and mental illness status, 2011-12
Table 12A.87	Age-standardised proportions of adults by long-term health conditions and mental illness status, 2011-12
Table 12A.88	Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2011-12 (per cent)
Table 12A.89	Age-standardised proportion of the population aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status, 2011-12 (per cent)
Table 12A.90	Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2007-08 (per cent)
Table 12A.91	Population aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status, 2007-08 (per cent)
Table 12A.92	Labour force and employment participation among adults aged 16–64 years, by mental disorder status, 2007 (per cent)
Table 12A.93	Education, training and employment participation among adults aged 16–30 years, by mental disorder status, 2007 (per cent)
Table 12A.94	Labour force and employment participation among adults aged 16–30 years, by mental disorder status, 2007 (per cent)
Table 12A.95	Proportion of people who had face-to-face contact with family or friends living outside the household in the last week, by mental illness status, 2014 (per cent)
Table 12A.96	Clinical outcomes of people receiving various types of mental health care provided by State and Territory public mental health services (per cent)
Table 12A.97	People who received mental health care provided by State and Territory public mental health services and who significantly improved, by service type and age group (per cent)
Table 12A.98	Deflators used to calculate real State and Territory mental health expenditure
Table 12A.99	Estimated resident populations used in mental health per head calculations

TABLE 12A.1

Table 12A.1 **Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million) (a), (b), (c)**

	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14
Mental health specific payments to states and territories (d)	89.9	88.8	94.4	92.4	4.0	6.9	19.6	51.8	62.5
National programs and initiative (Department of Health managed) (e)	92.4	105.5	230.2	208.3	203.7	248.0	295.0	410.5	450.8
National programs and initiative (Department of Social Services [DSS] or Families, Housing, Community Services and Indigenous Affairs [FaHCSIA] managed) (f)	..	10.5	97.9	163.2	153.8	153.2	160.0	185.6	210.9
National programs and initiative (DVA managed) (g)	158.2	170.5	170.0	179.7	172.8	170.1	166.6	173.6	174.5
Department of Defence-funded programs (h)	na	na	na	na	16.4	23.0	22.6	25.0	na
National Suicide Prevention Program (i)	10.9	21.0	22.0	23.4	23.8	26.2	54.6	57.0	56.5
Indigenous social and emotional wellbeing programmes (j)	27.8	29.2	31.6	37.5	38.7	43.8	42.7	44.5	44.5
MBS — Psychiatrists (k)	278.0	279.5	282.5	283.8	284.4	292.6	299.4	309.1	320.4
MBS — General practitioners (l)	293.3	182.5	158.9	198.3	215.4	248.2	209.6	199.8	216.4
MBS — Psychologists/Allied Health (m)	3.0	67.6	208.1	279.0	331.4	382.2	391.0	422.6	434.3
Pharmaceutical Benefits Schedule (n)	800.9	804.1	825.1	845.7	847.2	876.3	878.6	789.3	735.0
Private Health Insurance Premium Rebates (o)	73.1	78.1	87.4	83.6	102.3	101.6	118.0	109.3	108.6
Research (p)	34.5	37.0	43.3	51.3	56.8	63.0	65.2	69.3	75.3
National Mental Health Commission (q)	2.8	7.0	6.0
TOTAL	1 861.9	1 874.4	2 251.6	2 446.2	2 450.8	2 635.1	2 725.7	2 854.5	2 895.6
<i>Per cent</i>									
Mental health specific payments to states and territories (d)	4.8	4.7	4.2	3.8	0.2	0.3	0.7	1.8	2.2
National programs and initiative (DoHA managed) (e)	5.0	5.6	10.2	8.5	8.3	9.4	10.8	14.4	15.6
National programs and initiative (FaHCSIA managed) (f)	..	0.6	4.3	6.7	6.3	5.8	5.9	6.5	7.3
National programs and initiative (DVA managed) (g)	8.5	9.1	7.6	7.3	7.1	6.5	6.1	6.1	6.0
Department of Defence-funded programs (h)	na	na	na	na	0.7	0.9	0.8	0.9	na

TABLE 12A.1

Table 12A.1 **Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million) (a), (b), (c)**

	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14
National Suicide Prevention Program (i)	0.6	1.1	1.0	1.0	1.0	1.0	2.0	2.0	2.0
Indigenous social and emotional wellbeing programmes (j)	1.5	1.6	1.4	1.5	1.6	1.7	1.6	1.6	1.5
MBS — Psychiatrists (k)	14.9	14.9	12.5	11.6	11.6	11.1	11.0	10.8	11.1
MBS — General practitioners (l)	15.8	9.7	7.1	8.1	8.8	9.4	7.7	7.0	7.5
MBS — Psychologists/Allied Health (m)	0.2	3.6	9.2	11.4	13.5	14.5	14.3	14.8	15.0
Pharmaceutical Benefits Schedule (n)	43.0	42.9	36.6	34.6	34.6	33.3	32.2	27.7	25.4
Private Health Insurance Premium Rebates (o)	3.9	4.2	3.9	3.4	4.2	3.9	4.3	3.8	3.8
Research (p)	1.9	2.0	1.9	2.1	2.3	2.4	2.4	2.4	2.6
National Mental Health Commission (q)	0.1	0.2	0.2
TOTAL	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

(a) Detailed notes on how estimates specific to Commonwealth mental health specific expenditure are derived are provided in the AIHW Mental Health Services in Australia on-line publication. See <http://mhsa.aihw.gov.au/resources/expenditure/data-source/>.

(b) Estimated Australian Government expenditure shown in the table covers only those areas of expenditure that have a clear and identifiable mental health purpose. A range of other expenditure, both directly and indirectly related to provision of support for people affected by mental illness, is not covered in the table.

(c) Time series financial data are adjusted to 2013-14 dollars using the implicit price deflator for general government final consumption expenditure on hospital and nursing home services (table 12A.98).

(d) *Mental health specific payments to states and territories:* For years up to 2008-09, this category covers specific payments made to states and territories by the Australian Government for mental health reform under the Medicare Agreements 1993–1998, and Australian Health Care Agreements 1998–2003 and 2008-09. From July 2009 the Australian Government provided special purpose payments (SPP) to State and Territory governments under the National Healthcare Agreement (NHA) that do not specify the amount to be spent on mental health or any other health area. As a consequence, specific mental health funding cannot be identified under the NHA. From 2008-09 onwards, the amounts include: National Perinatal Depression Plan — Payments to States, National Partnership — Supporting Mental Health Reform and specific payments to Tasmania under the Tasmanian Health Assistance Package. Note that the expenditure reported here excludes payments to states and territories for the development of subacute mental health beds made under Schedule E of the National Partnership Agreement - Improving Public Hospital Services, which will total \$175 million over the period 2010-11 to 2013-14. Mental-health specific payments cannot be separately identified from payments for other categories of subacute beds made to states and territories.

Table 12A.1 **Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million) (a), (b), (c)**

	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14
(e) <i>National programs and initiatives (Department of Health managed)</i> : This category of expenditure includes the expenditure groups described in the AIHW Mental Health Services in Australia on-line publication. See http://mhsa.aihw.gov.au/resources/expenditure/data-source/ . Expenditure reported under the item 'Indigenous social and emotional wellbeing programmes' has previously been reported under 'National programs and initiatives (Department of Health managed)'. This expenditure is now separately reported following the transfer of the former Office for Aboriginal and Torres Strait Islander Health (OATSIH) Social and Emotional Wellbeing program to the Department of Prime Minister and Cabinet. Adjustments have been made to all years.									
(f) <i>National programs and initiatives (Department of Social Services [DSS] managed, before September 2013 it was Families, Housing, Community Services and Indigenous Affairs [FaHCSIA])</i> : Expenditure on DSS/FaHCSIA-managed Council of Australian Governments (COAG) Action Plan programs refers to funding outlays on three initiatives funded by the Australian Government under the COAG Action Plan on Mental Health (Personal Helpers and Mentors, More Respite Care Places to Help Families and Carers, Family Mental Health Support Services (previously referred to as Community based programmes to help families coping with mental illness). 2013-14 expenditure also includes expenditure on new Community Mental Health – drought assistance measure.									
(g) <i>National programs and initiatives (Department of Veterans' Affairs [DVA] managed)</i> : This category of expenditure includes the groups described in the AIHW Mental Health Services in Australia on-line publication. See http://mhsa.aihw.gov.au/resources/expenditure/data-source/ .									
(h) <i>Department of Defence-funded programs</i> : Expenditure reporting commences at 2009-10, and covers a range of mental health programs and services delivered to ADF personnel. Increased expenditure over the period reflects, in part, increased accuracy of data capture. Details of the ADF Mental Health Strategy are available at http://www.defence.gov.au/health/dmh/i-dmh.htm . Data are not available for 2013-14.									
(i) <i>National Suicide Prevention Program</i> : Expenditure reported includes all Australian Government allocations made under the national program, including additional funding made available under the COAG Action Plan and the 2010-11 and 2011-12 Federal Budgets.									
(j) <i>Indigenous social and emotional wellbeing programmes</i> : Previously reported under 'National programmes and initiatives (Health managed)' up to 2012-13 but has been separately identified following the transfer of the former OATSIH Social and Emotional Wellbeing program to the Department of Prime Minister and Cabinet.									
(k) <i>Medicare Benefits Schedule – Psychiatrists</i> : Expenditure reported refers to benefits paid for services by consultant psychiatrists processed in each of the index years. The amounts reported exclude payments made by the Department of Veterans' Affairs under the Repatriation Medical Benefits Schedule. These are included under the Department of Veterans' Affairs expenditure.									

Table 12A.1 **Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million) (a), (b), (c)**

	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14
(l) <i>Medicare Benefits Schedule – General Practitioner</i> : Prior to 2006-07, General Practitioner mental health-related expenditure was based on a crude estimate of 6.1 per cent of total MBS benefits paid for GP attendances, and derived from data and assumptions as detailed in the National Mental Health Report 2007. This estimate was historical and aimed to recognise that, although few mental health specific items were available in the MBS to accurately monitor GP mental health service provision, GPs are a significant provider of services to people with mental illness. Commencing November 2006, new mental health specific GP items were introduced under the Better Access to Mental Health Care initiative. To incorporate these changes, GP expenditure reported for 2006-07 is based on total MBS benefits paid against these new mental health specific items, plus an additional 6.1 per cent of total GP Benefits paid in the period preceding the introduction of the new items (July and November 2006). From 2007-08 onwards, expenditure on GP mental health care is based solely on benefits paid against MBS mental health specific GP items, which are predominantly the Better Access GP mental health items plus a small number of other items that were created in the years preceding the introduction of the Better Access initiative. This method provides a significantly lower expenditure figure than obtained using the 6.1 per cent estimate of previous years because it is conservative and does not attempt to assign a cost to the range of GP mental health work that is not billed as a specific mental health item. Comparisons of GP mental health related expenditure reported pre- and post-2006-07 are therefore not valid as the apparent decrease reflects the different approach to counting GP mental health services.									
(m) <i>Medicare Benefits Schedule – Psychologists/Allied Health</i> : Expenditure refers to MBS benefits paid for Clinical Psychologists, Psychologists, Social Workers and Occupational Therapists under the new items introduced through the Better Access to Mental Health Care initiative on 1 November 2006, plus a small number of Psychologist/Allied health items that were created under the Enhanced Primary Care program in the years preceding the introduction of the Better Access initiative.									
(n) <i>Pharmaceutical Benefits Scheme</i> : Expenditure under the Pharmaceutical Benefits Scheme refers to all Australian Government benefits for psychiatric medication in each of the index years, defined as drugs included in the following classes of the Anatomical Therapeutic Chemical Drug Classification System: antipsychotics (except prochlorperazine); anxiolytics; hypnotics and sedatives; psychostimulants; and antidepressants. Expenditure on Clozapine, funded under the Highly Specialised Drugs Program, has been included for all years, including Clozapine dispensed through public hospitals. The amounts reported exclude payments made by the Department of Veterans' Affairs under the Repatriation Pharmaceutical Benefits Schedule. These are included under the Department of Veterans' Affairs expenditure.									
(o) <i>Private Health Insurance Premium Rebates</i> : Estimates of the 'mental health share' of Australian Government Private Health Insurance Rebates are derived from a combination of sources and based on the assumption that a proportion of Australian Government outlays designed to increase public take up of private health insurance have subsidised private psychiatric care in hospitals. The methodology underpinning these estimates is described in the AIHW Mental Health Services in Australia on-line publication. See http://mhsa.aihw.gov.au/resources/expenditure/data-source/									
(p) <i>Research</i> : Research funding represents the value of mental health related grants administered by the National Health and Medical Research Council (NHMRC) during the relevant year. Data were sourced from the NHMRC website: http://www.nhmrc.gov.au/grants-funding/research-funding-statistics-and-data/mental-health-nhpa , accessed 28 September 2015. Historical adjustments were made to most years preceding 2013-14.									

Table 12A.1 **Real estimated Australian Government expenditure on mental health services (2013-14 dollars) (\$million) (a), (b), (c)**

	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14
--	---------	---------	---------	---------	---------	---------	---------	---------	---------

(q) *National Mental Health Commission*: The Commission commenced operation in January 2012.

na Not available. .. Not applicable.

Source: Department of Health (Australian Government) (unpublished).

TABLE 12A.2

Table 12A.2 **Real estimated recurrent expenditure on State and Territory governments specialised mental health services (2013-14 dollars) (a), (b), (c), (d)**

	<i>NSW (e)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Real recurrent expenditure (\$ million)</i>									
2005-06	1 096.9	867.6	596.2	416.0	287.9	87.7	57.5	35.6	3 445.3
2006-07	1 143.1	895.8	665.4	436.5	319.3	99.8	66.7	39.0	3 665.7
2007-08	1 196.0	929.2	752.0	473.4	337.2	107.0	70.4	40.9	3 906.1
2008-09	1 255.5	963.1	794.4	505.7	352.4	109.4	75.3	43.0	4 098.8
2009-10	1 327.0	1 003.3	853.7	514.6	357.4	119.0	75.3	43.9	4 294.1
2010-11	1 419.2	1 053.9	909.6	572.2	370.4	125.6	78.8	47.0	4 576.6
2011-12	1 475.9	1 065.6	946.0	616.9	368.6	113.2	83.9	51.4	4 721.5
2012-13	1 506.4	1 074.1	901.0	643.3	362.0	114.1	88.4	51.2	4 740.3
2013-14	1 567.1	1 124.9	892.0	666.5	398.6	111.7	90.7	54.2	4 905.7
<i>Real expenditure per person (\$)</i>									
2005-06	163.27	172.72	150.39	204.92	186.37	179.66	172.50	171.49	169.62
2006-07	168.45	175.50	164.07	210.17	204.53	203.07	197.25	184.91	177.71
2007-08	173.73	178.71	180.77	221.74	213.61	215.76	204.68	188.74	185.86
2008-09	179.32	181.26	185.80	228.95	220.53	217.94	214.59	193.17	190.86
2009-10	186.86	185.13	195.47	227.32	220.80	234.97	210.38	192.79	196.39
2010-11	197.67	191.77	205.00	246.74	226.87	246.09	216.08	203.96	206.41
2011-12	203.64	191.15	209.62	258.43	224.07	221.23	226.20	221.19	209.98
2012-13	204.98	189.11	195.41	260.15	217.76	222.57	232.80	216.18	206.94
2013-14	209.91	194.25	190.16	261.28	237.65	217.27	236.17	223.51	210.37

(a) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).

Table 12A.2 Real estimated recurrent expenditure on State and Territory governments specialised mental health services (2013-14 dollars) (a), (b), (c), (d)

	<i>NSW (e)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
--	----------------	------------	------------	-----------	-----------	------------	------------	-----------	-------------

(b) Estimates of expenditure on State and Territory governments' specialised mental health services include revenue from other sources (including patient fees and reimbursement by third party compensation insurers), Australian government funding provided under the Australian Health Care Agreement base grants/NHA SPP, 'other Australian Government funds', Australian Government mental health specific payments to states and territories and funding provided through the Department of Veterans' Affairs.

(c) Depreciation is excluded for all years.

(d) Due to the ongoing validation of National Minimum Data Set (NMDS), data could differ from previous reports.

(e) The quality of the NSW 2010-11 Mental Health Establishments (MHE) NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.

Source: Australian Institute of Health and Welfare (AIHW) (unpublished) Mental Health Establishments National Minimum Data Set (MHE NMDS); Australian Government (unpublished); ABS (various issues), Australian Demographic Statistics, December (various years), Cat. no. 3101.0.

TABLE 12A.3

Table 12A.3 **Real estimated expenditure on State and Territory governments' specialised mental health services, by funding source (2013-14 dollars) (\$million) (a), (b), (c), (d)**

	<i>NSW (e)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (f)</i>
<i>2005-06</i>									
State/Territory funds	1 028.2	815.2	564.1	400.6	271.4	81.8	54.6	33.6	3 249.5
Australian Government funds									
Mental health specific payments to states and territories (g)	27.4	21.6	18.0	9.2	7.2	2.7	1.9	1.8	89.8
Department of Veterans' Affairs (h)	10.9	10.4	1.2	2.1	4.1	0.2	0.2	–	29.0
Total Australian Government funds	38.2	32.0	19.1	11.4	11.2	2.9	2.1	1.8	118.8
Other revenue	30.5	20.5	13.0	4.0	5.2	3.0	0.8	0.1	77.0
Total funds	1 096.9	867.6	596.2	416.0	287.9	87.7	57.5	35.6	3 445.3
<i>2006-07</i>									
State/Territory funds	1 083.1	830.1	632.2	420.2	304.1	93.9	63.9	37.1	3 464.6
Australian Government funds									
Mental health specific payments to states and territories (g)	28.7	20.5	17.6	8.9	7.3	2.1	1.8	1.9	88.8
Department of Veterans' Affairs (h)	8.9	9.4	3.5	3.3	3.8	0.5	0.2	0.0	29.6
Total Australian Government funds	37.6	29.8	21.0	12.2	11.1	2.6	2.0	1.9	118.3
Other revenue	22.4	35.8	12.2	4.1	4.1	3.3	0.8	0.0	82.8
Total funds	1 143.1	895.8	665.4	436.5	319.3	99.8	66.7	39.0	3 665.7
<i>2007-08</i>									
State/Territory funds	1 136.8	872.3	719.7	457.6	320.3	100.3	67.1	38.2	3 712.2
Australian Government funds									
Mental health specific payments to states and territories (g)	29.5	22.2	18.7	9.4	7.3	2.6	2.6	2.2	94.3
Department of Veterans' Affairs (h)	8.8	7.2	2.7	2.8	4.4	0.4	0.3	–	26.6
Total Australian Government funds	38.3	29.4	21.4	12.2	11.6	2.9	2.8	2.2	120.9
Other revenue	20.9	27.4	10.9	3.6	5.3	3.8	0.6	0.4	73.0

TABLE 12A.3

Table 12A.3 **Real estimated expenditure on State and Territory governments' specialised mental health services, by funding source (2013-14 dollars) (\$million) (a), (b), (c), (d)**

	<i>NSW (e)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (f)</i>
Total funds	1 196.0	929.2	752.0	473.4	337.2	107.0	70.4	40.9	3 906.1
<i>2008-09</i>									
State/Territory funds	1 201.1	899.4	759.8	489.6	331.7	100.8	72.2	41.1	3 895.7
Australian Government funds									
Mental health specific payments to states and territories (g)	29.1	21.8	18.7	9.4	7.0	2.2	2.2	1.9	92.3
Department of Veterans' Affairs (h)	8.9	11.2	4.1	4.1	5.3	0.6	0.1	–	34.3
Total Australian Government funds	38.1	32.9	22.8	13.5	12.3	2.9	2.2	1.9	126.6
Other revenue	16.4	30.8	11.7	2.6	8.3	5.7	0.9	0.0	76.5
Total funds	1 255.5	963.1	794.4	505.7	352.4	109.4	75.3	43.0	4 098.8
<i>2009-10</i>									
State/Territory funds	1 304.0	957.5	837.3	508.6	346.9	113.9	73.9	43.7	4 185.8
Australian Government funds									
Mental health specific payments to states and territories (g)	1.1	0.9	0.7	0.5	0.3	0.2	0.2	0.2	3.9
Department of Veterans' Affairs (h)	9.8	9.9	4.3	2.6	4.2	0.6	0.4	–	31.8
Total Australian Government funds	10.9	10.8	5.0	3.1	4.5	0.8	0.5	0.2	35.7
Other revenue	12.1	35.0	11.4	2.9	6.0	4.3	0.8	–	72.6
Total funds	1 327.0	1 003.3	853.7	514.6	357.4	119.0	75.3	43.9	4 294.1
<i>2010-11</i>									
State/Territory funds	1 379.0	1 001.3	891.3	566.7	361.7	122.4	77.1	46.7	4 446.2
Australian Government funds									
Mental health specific payments to states and territories (g)	1.9	1.6	1.4	0.9	0.5	0.2	0.2	0.2	6.9
Department of Veterans' Affairs (h)	10.3	10.7	3.8	2.2	4.7	0.4	0.3	–	32.4

TABLE 12A.3

Table 12A.3 **Real estimated expenditure on State and Territory governments' specialised mental health services, by funding source (2013-14 dollars) (\$million) (a), (b), (c), (d)**

	<i>NSW (e)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (f)</i>
Total Australian Government funds	12.3	12.3	5.1	3.1	5.2	0.6	0.5	0.2	39.4
Other revenue	28.0	40.3	13.1	2.5	3.5	2.5	1.2	0.1	91.1
Total funds	1 419.2	1 053.9	909.6	572.2	370.4	125.6	78.8	47.0	4 576.6
<i>2011-12</i>									
State/Territory funds	1 442.4	1 005.5	922.3	605.5	357.8	109.6	82.1	50.8	4 576.0
Australian Government funds									
Mental health specific payments to states and territories (g)	6.4	4.2	3.5	2.8	1.4	0.4	0.6	0.4	19.6
Department of Veterans' Affairs (h)	10.0	9.5	3.0	1.8	4.1	0.5	0.3	0.0	29.3
Total Australian Government funds	16.4	13.7	6.5	4.6	5.6	0.8	0.9	0.5	48.9
Other revenue	17.1	46.4	17.3	6.8	5.2	2.7	0.9	0.1	96.5
Total funds	1 475.9	1 065.6	946.0	616.9	368.6	113.2	83.9	51.4	4 721.5
<i>2012-13</i>									
State/Territory funds	1 456.8	997.5	867.7	628.4	350.8	107.6	86.2	50.0	4 544.9
Australian Government funds									
Mental health specific payments to states and territories (g)	15.7	10.9	10.4	6.2	3.8	3.0	0.8	1.1	51.8
Department of Veterans' Affairs (h)	11.5	11.9	4.0	4.1	4.3	0.5	0.5	–	36.9
Total Australian Government funds	27.2	22.7	14.4	10.3	8.2	3.4	1.3	1.1	88.7
Other revenue	22.4	53.8	18.9	4.6	3.0	3.0	0.9	0.1	106.6
Total funds	1 506.4	1 074.1	901.0	643.3	362.0	114.1	88.4	51.2	4 740.3
<i>2013-14</i>									
State/Territory funds	1 520.0	1 050.9	852.0	647.5	387.4	98.7	88.7	52.9	4 698.1

TABLE 12A.3

Table 12A.3 **Real estimated expenditure on State and Territory governments' specialised mental health services, by funding source (2013-14 dollars) (\$million) (a), (b), (c), (d)**

	NSW (e)	Vic	Qld	WA	SA	Tas	ACT	NT	Aust (f)
Australian Government funds									
Mental health specific payments to states and territories (g)	16.6	11.9	17.4	7.3	4.1	3.0	0.9	1.2	62.5
Department of Veterans' Affairs (h)	13.1	10.4	3.9	3.8	4.8	0.6	0.3	–	36.9
Total Australian Government funds	29.8	22.3	21.3	11.1	8.9	3.6	1.2	1.2	99.4
Other revenue	17.3	51.8	18.8	7.9	2.3	9.3	0.8	0.1	108.2
Total funds	1 567.1	1 124.9	892.0	666.5	398.6	111.7	90.7	54.2	4 905.7

- (a) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).
- (b) Estimates of State and Territory government funds include Australian government funding provided under the Australian Health Care Agreement base grants/NHA SPP.
- (c) Depreciation is excluded for all years.
- (d) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (e) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.
- (f) The Australian total for mental health specific payments to states and territories can differ slightly to those in table 12A.1 as in that table the deflator for Australia is used, whereas in this table State or Territory specific deflators are used and the Australian total is the sum of states and territories.
- (g) *Mental health specific payments to states and territories*: For years up to 2008-09, this category covers specific payments made to states and territories by the Australian Government for mental health reform under the Medicare Agreements 1993–1998, and Australian Health Care Agreements 1998–2003 and 2008-09. From July 2009, the Australian Government provided SPP to State and Territory governments under the NHA that do not specify the amount to be spent on mental health or any other health area. As a consequence, specific mental health funding cannot be identified under the NHA. From 2008-09 onwards, the amounts include: National Perinatal Depression Plan – Payments to States, National Partnership – Supporting Mental Health Reform and specific payments to Tasmania under the Tasmanian Health Assistance Package. Note that the expenditure reported here excludes payments to states and territories for the development of subacute mental health beds made under Schedule E of the National Partnership Agreement – Improving Public Hospital Services, which will total \$175 million over the period 2010-11 to 2013-14. Mental-health specific payments cannot be separately identified from payments for other categories of subacute beds made to states and territories.
- (h) *Department of Veterans' Affairs*: This category of expenditure includes the groups described in the AIHW Mental Health Services in Australia on-line publication. See <http://mhsa.aihw.gov.au/resources/expenditure/data-source/>.

Table 12A.3 **Real estimated expenditure on State and Territory governments' specialised mental health services, by funding source (2013-14 dollars) (\$million) (a), (b), (c), (d)**

	<i>NSW (e)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (f)</i>
--	----------------	------------	------------	-----------	-----------	------------	------------	-----------	-----------------

– Nil or rounded to zero.

Source: AIHW (unpublished) MHE NMDS; Department of Health (Australian Government) (unpublished).

Table 12A.4 Real Australian, State and Territory governments expenditure on mental health services (2013-14 dollars) (\$million), (a), (b), (c), (d)

	<i>State and Territory governments</i>	<i>Australian Government</i>	<i>Total</i>
<i>Real expenditure (\$million)</i>			
2005-06	3 326.6	1 861.9	5 188.4
2006-07	3 547.3	1 874.4	5 421.8
2007-08	3 785.2	2 251.6	6 036.7
2008-09	3 972.2	2 446.2	6 418.4
2009-10	4 258.4	2 450.8	6 709.1
2010-11	4 537.3	2 635.1	7 172.3
2011-12	4 672.5	2 725.7	7 398.3
2012-13	4 651.6	2 854.5	7 506.0
2013-14	4 806.3	2 895.6	7 701.9
<i>Expenditure per person</i>			
2005-06	163.78	91.67	255.44
2006-07	171.97	90.87	262.84
2007-08	180.11	107.14	287.24
2008-09	184.96	113.91	298.87
2009-10	194.75	112.08	306.83
2010-11	204.64	118.84	323.48
2011-12	207.80	121.22	329.03
2012-13	203.07	124.61	327.68
2013-14	206.11	124.17	330.28
<i>Proportion of expenditure</i>			
2005-06	64.1	35.9	100.0
2006-07	65.4	34.6	100.0
2007-08	62.7	37.3	100.0
2008-09	61.9	38.1	100.0
2009-10	63.5	36.5	100.0
2010-11	63.3	36.7	100.0
2011-12	63.2	36.8	100.0
2012-13	62.0	38.0	100.0
2013-14	62.4	37.6	100.0

(a) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).

(b) The estimate of State and Territory governments' expenditure relates to expenditure on specialised mental health services (tables 12A.2 and 12A.3) less Australian Governments expenditure on 'Mental health specific payments to states and territories' and the Department of Veterans' Affairs as reported in table 12A.3. It includes expenditure sourced from other revenue (as reported in table 12A.3) and Australian Government funding provided under the Australian Health Care Agreement base grants/NHA SPP.

(c) Depreciation is excluded for all years.

(d) Due to the ongoing validation of NMDS, data could differ from previous reports.

Table 12A.4 Real Australian, State and Territory governments expenditure on mental health services (2013-14 dollars) (\$million), (a), (b), (c), (d)

	<i>State and Territory governments</i>	<i>Australian Government</i>	<i>Total</i>
<i>Source:</i>	AIHW (unpublished) MHE NMDS; Department of Health (Australian Government), unpublished; table 12A.98.		

Table 12A.5 Depreciation expenditure on State and Territory governments' specialised mental health services (current prices) (\$million) (a), (b)

	<i>NSW (c)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
2005-06	15.3	7.4	8.5	4.3	0.1	–	0.3	–	35.7
2006-07	12.4	7.2	9.7	4.1	–	–	–	–	33.4
2007-08	13.8	11.3	9.1	3.5	0.4	–	–	0.5	38.8
2008-09	9.0	12.9	8.2	4.1	3.2	–	–	–	37.5
2009-10	14.4	19.7	7.7	4.3	2.5	–	–	–	48.5
2010-11	13.2	29.6	9.2	4.3	1.5	–	–	–	57.9
2011-12	13.6	26.3	9.1	4.8	1.1	–	–	–	54.8
2012-13	16.3	20.3	7.1	4.0	–	–	–	–	47.8
2013-14	17.0	18.5	8.3	8.4	5.9	–	–	–	58.0

(a) See the *AIHW Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure estimates.

(b) Due to the ongoing validation of NMDS, data could differ from previous reports.

(c) The quality of the NSW 2010-11 MHE NMDS data has been affected by the reconfiguration of the service system during the year.

– Nil or rounded to zero.

Source: AIHW (unpublished) MHE NMDS.

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
<i>2005-06</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	191.2	30.2	65.7	63.1	81.3	431.4
Public acute hospital	266.1	193.6	177.1	92.1	36.6	22.5	9.0	10.4	807.6
<i>Total admitted patient (i)</i>	457.3	223.8	242.7	155.2	117.9	22.5	9.0	10.4	1 239.0
Community residential	24.4	121.9	..	4.9	2.8	16.0	5.9	0.3	176.3
Ambulatory	307.7	269.6	159.8	142.2	79.3	25.2	23.9	12.4	1 020.2
Non-government organisations	31.7	61.1	25.3	16.5	14.7	1.7	5.1	3.1	159.3
Indirect	64.2	32.2	26.8	7.2	5.1	5.0	2.2	2.0	144.7
Total expenditure	885.4	708.6	454.6	326.1	219.9	70.4	46.2	28.2	2 739.4
<i>Per cent</i>									
Public psychiatric hospital	21.6	4.3	14.4	19.3	37.0	15.7
Public acute hospital	30.1	27.3	39.0	28.3	16.7	32.0	19.5	36.8	29.5
<i>Total admitted patient (i)</i>	51.7	31.6	53.4	47.6	53.6	32.0	19.5	36.8	45.2
Community residential	2.8	17.2	..	1.5	1.3	22.8	12.9	0.9	6.4
Ambulatory	34.8	38.0	35.1	43.6	36.1	35.7	51.7	44.1	37.2
Non-government organisations	3.6	8.6	5.6	5.1	6.7	2.4	11.1	10.9	5.8
Indirect	7.2	4.5	5.9	2.2	2.3	7.1	4.8	7.3	5.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2006-07</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	189.2	32.9	70.4	66.9	80.0	439.4
Public acute hospital	310.2	206.2	190.0	98.6	55.0	29.8	14.2	10.3	914.3
<i>Total admitted patient (i)</i>	499.4	239.1	260.4	165.5	135.0	29.8	14.2	10.3	1 353.8
Community residential	27.8	124.7	..	6.5	3.0	18.5	6.9	0.3	187.7

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
Ambulatory	332.9	283.9	208.9	154.5	88.5	27.7	27.4	15.2	1 139.0
Non-government organisations	40.5	64.3	32.5	18.0	21.8	3.3	5.3	4.1	189.8
Indirect	63.2	43.0	29.3	10.7	4.8	4.3	1.9	2.1	159.1
Total expenditure	963.8	754.9	531.1	355.2	253.1	83.5	55.7	32.1	3 029.3
<i>Per cent</i>									
Public psychiatric hospital	19.6	4.4	13.3	18.8	31.6	14.5
Public acute hospital	32.2	27.3	35.8	27.8	21.7	35.7	25.5	32.1	30.2
<i>Total admitted patient (i)</i>	51.8	31.7	49.0	46.6	53.4	35.7	25.5	32.1	44.7
Community residential	2.9	16.5	..	1.8	1.2	22.1	12.4	1.1	6.2
Ambulatory	34.5	37.6	39.3	43.5	35.0	33.2	49.3	47.5	37.6
Non-government organisations	4.2	8.5	6.1	5.1	8.6	3.9	9.5	12.8	6.3
Indirect	6.6	5.7	5.5	3.0	1.9	5.1	3.4	6.6	5.3
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2007-08</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	181.5	39.7	77.8	67.5	80.6	447.1
Public acute hospital	340.7	221.1	221.5	113.0	60.2	34.2	16.0	11.7	1 018.5
<i>Total admitted patient (i)</i>	522.2	260.8	299.3	180.6	140.8	34.2	16.0	11.7	1 465.5
Community residential	15.1	131.3	..	9.1	6.3	19.3	7.4	0.5	189.1
Ambulatory	372.7	303.4	249.2	174.6	98.7	29.2	27.1	16.4	1 271.3
Non-government organisations	60.4	65.6	39.4	21.1	24.5	4.7	6.1	3.8	225.6
Indirect	66.8	42.0	33.2	13.7	5.7	4.7	3.3	2.4	171.9
Total expenditure	1 037.1	803.2	621.1	399.1	276.0	92.1	59.9	34.8	3 323.4
<i>Per cent</i>									
Public psychiatric hospital	17.5	4.9	12.5	16.9	29.2	13.5

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
Public acute hospital	32.9	27.5	35.7	28.3	21.8	37.1	26.7	33.5	30.6
<i>Total admitted patient (i)</i>	50.3	32.5	48.2	45.2	51.0	37.1	26.7	33.5	44.1
Community residential	1.5	16.3	..	2.3	2.3	21.0	12.4	1.3	5.7
Ambulatory	35.9	37.8	40.1	43.7	35.8	31.7	45.2	47.1	38.3
Non-government organisations	5.8	8.2	6.3	5.3	8.9	5.1	10.2	11.0	6.8
Indirect	6.4	5.2	5.3	3.4	2.1	5.2	5.5	7.0	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2008-09</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	181.5	37.2	80.6	74.3	80.7	454.4
Public acute hospital	405.6	241.0	227.8	124.4	66.0	35.9	16.4	13.0	1 130.1
<i>Total admitted patient (i)</i>	587.2	278.2	308.4	198.7	146.8	35.9	16.4	13.0	1 584.5
Community residential	13.9	142.2	..	12.9	9.1	19.1	9.9	0.9	208.0
Ambulatory	401.9	323.5	285.2	193.4	113.3	32.1	31.4	17.2	1 397.8
Non-government organisations	57.7	70.0	46.1	23.7	24.0	4.7	6.2	3.6	236.0
Indirect	54.0	45.3	41.8	14.4	6.7	5.3	2.7	3.2	173.3
Total expenditure	1 114.6	859.2	681.5	443.0	299.9	97.0	66.5	37.9	3 599.6
<i>Per cent</i>									
Public psychiatric hospital	16.3	4.3	11.8	16.8	26.9	12.6
Public acute hospital	36.4	28.0	33.4	28.1	22.0	37.0	24.7	34.2	31.4
<i>Total admitted patient (i)</i>	52.7	32.4	45.3	44.9	48.9	37.0	24.7	34.2	44.0
Community residential	1.2	16.6	..	2.9	3.0	19.7	14.8	2.3	5.8
Ambulatory	36.1	37.7	41.9	43.6	37.8	33.0	47.2	45.4	38.8
Non-government organisations	5.2	8.1	6.8	5.3	8.0	4.8	9.3	9.6	6.6
Indirect	4.8	5.3	6.1	3.2	2.2	5.5	4.0	8.4	4.8

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2009-10</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	221.2	43.8	86.9	75.4	72.8	500.2
Public acute hospital	416.8	252.1	244.1	131.5	74.8	43.1	16.6	12.9	1 192.0
<i>Total admitted patient (i)</i>	638.0	296.0	331.0	206.8	147.7	43.1	16.6	12.9	1 692.2
Community residential	11.9	152.3	..	14.9	9.0	20.2	10.6	1.3	220.3
Ambulatory	434.3	344.6	338.4	206.6	123.4	34.0	30.9	19.4	1 531.6
Non-government organisations	68.3	74.7	50.3	25.8	30.2	5.5	7.9	3.7	266.3
Indirect	65.5	56.6	46.9	14.1	6.9	6.3	2.5	2.8	201.7
Total expenditure	1 218.1	924.2	766.6	468.2	317.2	109.2	68.6	40.1	3 912.1
<i>Per cent</i>									
Public psychiatric hospital	18.2	4.7	11.3	16.1	23.0	12.8
Public acute hospital	34.2	27.3	31.8	28.1	23.6	39.5	24.3	32.2	30.5
<i>Total admitted patient (i)</i>	52.4	32.0	43.2	44.2	46.6	39.5	24.3	32.2	43.3
Community residential	1.0	16.5	..	3.2	2.9	18.6	15.5	3.2	5.6
Ambulatory	35.7	37.3	44.1	44.1	38.9	31.2	45.1	48.3	39.1
Non-government organisations	5.6	8.1	6.6	5.5	9.5	5.0	11.5	9.3	6.8
Indirect	5.4	6.1	6.1	3.0	2.2	5.8	3.6	7.0	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2010-11</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	244.0	42.2	90.4	82.1	66.8	525.5
Public acute hospital	449.7	271.3	254.0	151.1	75.0	44.5	18.6	14.3	1 278.5
<i>Total admitted patient (i)</i>	693.7	313.5	344.4	233.2	141.8	44.5	18.6	14.3	1 804.0

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
Community residential	12.3	164.4	..	17.7	11.8	21.0	10.0	1.5	238.7
Ambulatory	467.4	368.8	364.4	221.4	135.7	36.2	32.3	20.9	1 647.2
Non-government organisations	72.6	80.4	65.6	28.5	36.5	7.7	8.6	3.4	303.2
Indirect	69.2	58.3	60.7	24.6	6.3	7.0	3.0	3.3	232.2
Total expenditure	1 315.3	985.4	835.0	525.5	331.9	116.4	72.5	43.3	4 225.4
<i>Per cent</i>									
Public psychiatric hospital	18.6	4.3	10.8	15.6	20.1	12.4
Public acute hospital	34.2	27.5	30.4	28.8	22.6	38.2	25.6	33.0	30.3
<i>Total admitted patient (i)</i>	52.7	31.8	41.2	44.4	42.7	38.2	25.6	33.0	42.7
Community residential	0.9	16.7	..	3.4	3.5	18.1	13.8	3.4	5.6
Ambulatory	35.5	37.4	43.6	42.1	40.9	31.1	44.6	48.3	39.0
Non-government organisations	5.5	8.2	7.9	5.4	11.0	6.6	11.9	7.8	7.2
Indirect	5.3	5.9	7.3	4.7	1.9	6.0	4.1	7.5	5.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2011-12</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	238.2	40.8	97.4	88.8	63.8	529.0
Public acute hospital	510.0	274.6	267.5	167.2	73.6	40.6	19.4	16.0	1 368.9
<i>Total admitted patient (i)</i>	748.2	315.4	364.9	256.0	137.5	40.6	19.4	16.0	1 897.9
Community residential	12.7	164.1	..	21.6	18.4	19.8	11.0	1.5	249.2
Ambulatory	499.5	394.4	401.5	240.3	144.5	34.3	35.4	23.3	1 773.1
Non-government organisations	70.3	83.6	69.4	31.8	33.5	6.5	10.5	3.6	309.3
Indirect	68.4	56.1	55.5	31.9	8.6	6.3	2.8	4.3	233.9
Total expenditure	1 399.2	1 013.6	891.3	581.5	342.5	107.5	79.2	48.6	4 463.4
<i>Per cent</i>									

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
Public psychiatric hospital	17.0	4.0	10.9	15.3	18.6	11.9
Public acute hospital	36.5	27.1	30.0	28.8	21.5	37.7	24.5	32.9	30.7
<i>Total admitted patient (i)</i>	53.5	31.1	40.9	44.0	40.1	37.7	24.5	32.9	42.5
Community residential	0.9	16.2	..	3.7	5.4	18.5	13.9	3.1	5.6
Ambulatory	35.7	38.9	45.0	41.3	42.2	31.9	44.7	47.9	39.7
Non-government organisations	5.0	8.3	7.8	5.5	9.8	6.1	13.3	7.3	6.9
Indirect	4.9	5.5	6.2	5.5	2.5	5.9	3.5	8.8	5.2
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>2012-13</i>									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	251.0	40.5	94.4	91.4	57.9	535.2
Public acute hospital	561.7	286.3	277.7	186.7	72.3	39.7	22.0	17.8	1 464.3
<i>Total admitted patient (i)</i>	812.7	326.8	372.2	278.1	130.1	39.7	22.0	17.8	1 999.4
Community residential	10.9	172.1	..	23.2	19.2	22.2	11.7	2.1	261.4
Ambulatory	498.8	401.6	402.0	248.0	157.0	36.1	35.7	23.5	1 802.6
Non-government organisations	76.1	89.4	58.9	41.7	31.7	6.5	13.3	3.6	321.2
Indirect	72.3	56.4	41.9	32.7	11.2	6.8	3.0	3.3	227.5
Total expenditure	1 470.8	1 046.3	875.0	623.6	349.2	111.2	85.7	50.3	4 612.2
<i>Per cent</i>									
Public psychiatric hospital	17.1	3.9	10.8	14.7	16.6	11.6
Public acute hospital	38.2	27.4	31.7	29.9	20.7	35.7	25.7	35.4	31.7
<i>Total admitted patient (i)</i>	55.3	31.2	42.5	44.6	37.3	35.7	25.7	35.4	43.4
Community residential	0.7	16.4	..	3.7	5.5	20.0	13.7	4.2	5.7
Ambulatory	33.9	38.4	45.9	39.8	45.0	32.4	41.6	46.7	39.1
Non-government organisations	5.2	8.5	6.7	6.7	9.1	5.8	15.5	7.1	7.0

TABLE 12A.6

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g)	WA	SA	Tas (h)	ACT	NT	Aust
Indirect	4.9	5.4	4.8	5.2	3.2	6.1	3.5	6.5	4.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2013-14 (j)									
<i>Recurrent expenditure (\$million)</i>									
Public psychiatric hospital	255.0	44.4	87.2	86.3	58.7	531.8
Public acute hospital	615.7	303.4	294.4	205.7	98.2	39.6	23.7	18.6	1 599.2
<i>Total admitted patient (i)</i>	870.7	347.9	381.6	292.0	157.0	39.6	23.7	18.6	2 131.0
Community residential	10.3	190.8	..	26.7	23.0	21.7	12.3	3.1	288.0
Ambulatory	518.9	426.1	402.9	259.8	171.6	37.4	36.2	25.4	1 878.4
Non-government organisations	84.3	101.3	66.5	44.4	36.7	8.3	15.6	4.2	361.4
Indirect	82.9	58.8	41.0	43.5	10.3	4.6	2.9	2.9	247.0
Total expenditure	1 567.1	1 124.9	892.0	666.5	398.6	111.7	90.7	54.2	4 905.7
<i>Per cent</i>									
Public psychiatric hospital	16.3	4.0	9.8	13.0	14.7	10.8
Public acute hospital	39.3	27.0	33.0	30.9	24.6	35.5	26.1	34.3	32.6
<i>Total admitted patient (i)</i>	55.6	30.9	42.8	43.8	39.4	35.5	26.1	34.3	43.4
Community residential	0.7	17.0	..	4.0	5.8	19.5	13.5	5.8	5.9
Ambulatory	33.1	37.9	45.2	39.0	43.1	33.5	39.9	46.8	38.3
Non-government organisations	5.4	9.0	7.5	6.7	9.2	7.4	17.2	7.8	7.4
Indirect	5.3	5.2	4.6	6.5	2.6	4.1	3.2	5.4	5.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

(a) Expenditure is current prices for all years and includes all spending, regardless of source of funds.

(b) Depreciation is excluded for all years.

(c) See the AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure estimates.

Table 12A.6 **Total state and territory recurrent expenditure on specialised mental health services (current prices) (a), (b), (c), (d), (e)**

	<i>NSW (f)</i>	<i>Vic</i>	<i>Qld (g)</i>	<i>WA</i>	<i>SA</i>	<i>Tas (h)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
(d)	Due to the ongoing validation of NMDS, data could differ from previous reports.								
(e)	Totals may not add due to rounding.								
(f)	The quality of the NSW 2010-11 <i>MHE NMDS</i> data has been affected by the reconfiguration of the service system during the year.								
(g)	Queensland does not fund community residential services, however, it funds a number of extended treatment services, both campus and non-campus based, which provide longer term inpatient treatment and rehabilitation services with a full clinical staffing 24 hours a day seven days a week. In addition, Queensland have advised that funding to non-government services for psychiatric disability support services is administered either by Queensland Health or Disability Services Queensland (DSQ).								
(h)	For Tasmania, in 2005-06, non-government organisations (NGOs) providing residential services were included for the first time in the community residential category. As these NGOs are now categorised as residential services, NGO funding decreased from previous years. Indirect/residual expenditure represents State indirect/residual expenditure. If organisational indirect expenditure were included this expenditure would have been \$10 719 100.								
(i)	Includes expenditure on public hospital services managed and operated by private and non-government entities.								
(j)	For SA, the increases in admitted patient and ambulatory care expenditure in 2013-14 partly relate to genuine increases in mental health services. However, a significant proportion of the increases relate to improved identification and allocation of direct care and general overhead expenditure to mental health services.								

.. Not applicable.

Source: AIHW (unpublished) MHE NMDS.

Table 12A.7 Functioning and quality of life measures, by 12-month mental disorder status, 2007 (per cent) (a)

	<i>Any 12-month mental disorder (b)</i>	<i>No 12-month mental disorder</i>	<i>Total</i>
<i>Level of psychological distress (c)</i>			
Low	10.9 ± 1.1	89.1 ± 1.0	100.0
Moderate	32.0 ± 2.6	68.0 ± 2.5	100.0
High	57.1 ± 5.1	42.9 ± 5.1	100.0
Very high	79.6 ± 7.2	20.4 ± 7.1	100.0
<i>Disability status (d)</i>			
Profound/severe	42.9 ± 8.2	57.1 ± 8.2	100.0
Moderate/mild	32.1 ± 5.5	67.9 ± 5.6	100.0
Schooling/employment restriction only	43.4 ± 7.1	56.6 ± 7.1	100.0
No disability/no specific limitations or restrictions	16.6 ± 1.1	83.4 ± 1.1	100.0
<i>Days out of role (e)</i>			
0 days	14.7 ± 1.3	85.3 ± 1.3	100.0
1 to 7 days	28.5 ± 2.5	71.5 ± 2.5	100.0
More than 7 days	42.0 ± 5.2	58.0 ± 5.2	100.0
<i>Suicidal behaviour</i>			
Ideation (f)	71.7 ± 8.7	28.3 ± 8.7	100.0
Plans	77.5 ± 12.6	22.5* ± 12.6	100.0
Attempts	94.2 ± 9.0	5.8** ± 8.9	100.0
No suicidal behaviours	18.7 ± 1.1	81.3 ± 1.1	100.0
Total people aged 16–85 years	20.0 ± 1.1	80.0 ± 1.1	100.0

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent). A '*' indicates a relative standard error (RSE) of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution. A '**' indicates a RSE of greater than 50 per cent. Estimates with RSEs greater than 50 per cent are considered too unreliable for general use.

(b) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(c) Level of psychological distress is measured by the Kessler Psychological Distress Scale (K10), from which a score of 10 to 50 is produced. Higher scores indicate a higher level of distress; low scores indicate a low level of distress. Scores are grouped as follows: Low 10–15, Moderate 16–21, High 22–29, and Very high 30–50.

(d) Disability status relates to whether a person has disability, a core-activity limitation (mild, moderate, severe or profound), or a schooling or employment restriction.

(e) People who were unable to carry out or had to cut down on their usual activities in the 30 days prior to interview. Total includes 'not stated'.

(f) Suicidal ideation refers to the presence of serious thoughts about committing suicide.

Source: ABS (2008) *National Survey of Mental Health and Wellbeing: Summary of Results, 2007*, Cat. no. 4326.0.

TABLE 12A.8

Table 12A.8 **Age-standardised rate of adults with very high levels of psychological distress, by State and Territory, 2011-12 (a), (b), (c), (d)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (e)</i>	<i>Aust</i>
<i>Age standardised proportion</i>										
Males	%	2.5	3.3	2.9	2.1	2.8	2.3*	2.5	2.4*	2.7
Females	%	3.8	4.0	4.8	3.8	3.7	4.0	3.1	4.0	4.1
Total	%	3.2	3.7	3.9	3.0	3.3	3.2	2.8	3.2	3.4
<i>Relative standard errors</i>										
Males	%	20.2	15.5	18.6	23.9	20.3	32.2	22.1	34.5	9.5
Females	%	13.4	15.7	13.2	14.1	16.8	20.7	20.7	23.0	7.1
Total	%	12.0	11.7	12.1	13.0	12.7	17.7	15.4	20.1	5.9
<i>95 per cent confidence intervals</i>										
Males	±	1.0	1.0	1.1	1.0	1.1	1.5	1.1	1.6	0.5
Females	±	1.0	1.2	1.2	1.0	1.2	1.6	1.3	1.8	0.6
Total	±	0.7	0.8	0.9	0.8	0.8	1.1	0.9	1.3	0.4

(a) Denominator includes a small number of persons for whom levels of psychological distress were unable to be determined.

(b) Adults are defined as persons aged 18 years and over.

(c) Rates are age standardised by State and Territory, to the June 2001 Estimated Resident Population (5 year ranges from 18 for adults).

(d) Estimates with a "*" have a relative standard error between 25 per cent and 50 per cent and should be used with caution.

(e) Data for NT should be interpreted with caution as the *Australian Health Survey (AHS) (2011-12 National Health Survey [NHS] component)* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

Source: ABS (unpublished) *AHS 2011-13* (2011-12 NHS component), Cat. no. 4364.0.

Table 12A.9 Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2011-12 (a), (b), (c), (d)

	<i>Age standardised proportion (%)</i>	<i>Relative standard error (%)</i>	<i>95 % confidence interval (±)</i>
<i>Remoteness of residence</i>			
Major cities	3.3	8.0	0.5
Inner regional	3.8	12.8	0.9
Outer regional	3.5	19.2	1.3
Remote	2.9*	42.1	2.4
Very remote (e)
<i>SEIFA of residence (quintiles) (f)</i>			
Quintile 1	5.4	12.6	1.3
Quintile 2	4.1	8.8	0.7
Quintile 3	3.5	12.7	0.9
Quintile 4	2.8	13.3	0.7
Quintile 5	1.9	17.2	0.6
<i>SEIFA of residence (deciles) (f)</i>			
Decile 1	5.7	15.9	1.8
Decile 2	5.2	17.4	1.8
Decile 3	3.9	14.8	1.1
Decile 4	4.2	14.5	1.2
Decile 5	4.1	17.5	1.4
Decile 6	2.9	15.8	0.9
Decile 7	3.0	18.8	1.1
Decile 8	2.7	21.5	1.1
Decile 9	2.0	23.8	1.0
Decile 10	1.7*	25.3	0.9
<i>Disability status</i>			
With disability or restrictive long-term health condition	8.2	6.7	1.1
No disability or restrictive long-term health condition	1.1	9.4	0.2

SEIFA = Socio-Economic Indexes for Areas

- (a) Denominator includes a small number of persons for whom levels of psychological distress were unable to be determined.
- (b) Adults are defined as persons aged 18 years and over.
- (c) Rates are age standardised by State and Territory to the June 2001 Estimated Resident Population (5 year ranges from 18).
- (d) Estimates with a "*" have a relative standard error between 25 per cent and 50 per cent and should be used with caution.
- (e) Very remote data was not collected in the 2011-12 NHS component of the 2011-13 AHS.
- (f) For the Socioeconomic Index for Areas, Index of relative disadvantage data, quintile/decile 1 contains areas of most disadvantage.

Table 12A.9 **Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2011-12 (a), (b), (c), (d)**

	<i>Age standardised proportion (%)</i>	<i>Relative standard error (%)</i>	<i>95 % confidence interval (\pm)</i>
--	--	--	--

.. Not applicable.

Source: ABS (unpublished) *AHS 2011–13 (2011-12 NHS component)*, Cat. no. 4364.0.

TABLE 12A.10

Table 12A.10 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2011-12 (a), (b), (c), (d)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (e)</i>	<i>Aust</i>
<i>Age standardised proportion</i>										
<i>Remoteness of residence</i>										
Major cities	%	10.6	10.7	10.6	10.3	10.5	..	9.1	..	10.6
Inner regional	%	9.9	13.1	11.9	13.3	11.0*	8.8	–	..	11.4
Outer regional/remote	%	8.3*	13.2*	9.9	9.8	16.8	10.4	..	9.0	10.8
Very remote (e)	%
<i>SEIFA of residence (quintiles) (f)</i>										
Quintile 1	%	15.9	16.4	19.6	16.5	17.6	11.2	np	11.1	16.7
Quintile 2	%	14.0	13.0	11.9	13.4	12.5	9.3	11.4*	6.8*	12.9
Quintile 3	%	11.0	11.6	11.3	10.3	8.2	10.2	11.0*	10.0*	10.9
Quintile 4	%	8.3	9.6	7.7	6.7	5.9*	6.7*	10.6	9.1*	8.1
Quintile 5	%	5.7	7.8	8.1	8.3	10.1	5.9*	7.3	6.8*	7.4
<i>Disability status</i>										
With disability or restrictive long-term health condition	%	21.2	26.6	21.4	22.1	24.3	17.4	17.5	20.4	22.7
No disability or restrictive long-term health condition	%	5.2	4.8	5.1	4.7	5.1	3.8	4.4	3.8	5.0
Total	%	10.4	11.4	10.8	10.6	11.4	9.1	9.1	9.0	10.8
<i>Relative standard errors</i>										
<i>Remoteness of residence</i>										
Major cities	%	6.5	6.9	8.4	8.0	8.5	–	9.7	..	3.4
Inner regional	%	16.9	13.4	13.7	22.4	29.3	10.5	–	..	7.3
Outer regional/remote	%	44.2	31.7	22.0	19.2	19.2	16.9	..	15.0	11.5
Very remote (e)	%

TABLE 12A.10

Table 12A.10 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2011-12 (a), (b), (c), (d)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (e)</i>	<i>Aust</i>
<i>SEIFA of residence (quintiles) (f)</i>										
Quintile 1	%	12.7	11.9	17.3	13.4	13.1	14.2	np	23.5	7.8
Quintile 2	%	12.2	12.6	11.9	15.2	11.9	17.4	34.7	36.8	5.2
Quintile 3	%	17.6	12.2	10.6	17.4	21.2	16.5	26.4	30.6	6.1
Quintile 4	%	17.1	15.9	16.1	16.5	29.7	28.7	15.9	25.6	9.6
Quintile 5	%	19.8	20.9	16.4	19.2	24.7	47.7	16.9	28.4	9.0
<i>Disability status</i>										
With disability or restrictive long-term health condition	%	9.5	7.1	7.7	8.1	8.5	11.5	13.1	15.3	3.7
No disability or restrictive long-term health condition	%	11.0	10.6	11.7	14.2	12.6	18.7	16.1	21.8	4.8
Total	%	6.7	6.2	6.3	7.2	7.3	8.8	9.7	15.0	3.2
<i>95 per cent confidence intervals</i>										
<i>Remoteness of residence</i>										
Major cities	±	1.4	1.4	1.7	1.6	1.8	..	1.7	..	0.7
Inner regional	±	3.3	3.4	3.2	5.8	6.3	1.8	–	..	1.6
Outer regional/remote	±	7.2	8.2	4.3	3.7	6.3	3.5	..	2.7	2.4
Very remote (e)	±
<i>SEIFA of residence (quintiles) (f)</i>										
Quintile 1	±	4.0	3.8	6.6	4.3	4.5	3.1	np	5.1	2.5
Quintile 2	±	3.3	3.2	2.8	4.0	2.9	3.2	7.7	4.9	1.3
Quintile 3	±	3.8	2.8	2.4	3.5	3.4	3.3	5.7	6.0	1.3
Quintile 4	±	2.8	3.0	2.4	2.1	3.4	3.8	3.3	4.5	1.5
Quintile 5	±	2.2	3.2	2.6	3.1	4.9	5.5	2.4	3.8	1.3

TABLE 12A.10

Table 12A.10 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2011-12 (a), (b), (c), (d)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (e)</i>	<i>Aust</i>
<i>Disability status</i>										
With disability or restrictive long-term health condition	±	3.9	3.7	3.2	3.5	4.1	3.9	4.5	6.1	1.7
No disability or restrictive long-term health condition	±	1.1	1.0	1.2	1.3	1.3	1.4	1.4	1.6	0.5
Total	±	1.4	1.4	1.3	1.5	1.6	1.6	1.7	2.7	0.7

SEIFA = Socio-Economic Indexes for Areas

(a) Total includes a small number of persons for whom levels of psychological distress were unable to be determined.

(b) Adults are defined as persons aged 18 years and over.

(c) Rates are age standardised by State and Territory, to the June 2001 Estimated Resident Population (5 year ranges from 18).

(d) Estimates with a "*" have a relative standard error between 25 per cent and 50 per cent and should be used with caution.

(e) Data for NT should be interpreted with caution as the *AHS (2011-12 NHS component)* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

(f) For the Socioeconomic Index for Areas, Index of relative disadvantage data, quintile/decile 1 contains areas of most disadvantage.

.. Not applicable. – Nil or rounded to zero. **np** Not published.

Source: ABS (unpublished) *AHS 2011-13 (2011-12 NHS component)*, Cat. no. 4364.

TABLE 12A.11

Table 12A.11 **Age-standardised rate of adults with very high levels of psychological distress, by State and Territory, 2007-08 (a), (b)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Age standardised proportion (c), (d)</i>										
Males	%	3.2	3.0	2.0	2.3	3.5	*2.5	np	np	2.8
Females	%	4.8	4.0	4.1	3.3	3.5	*4.0	np	np	4.1
Total	%	4.0	3.5	3.1	2.8	3.5	3.3	3.4	np	3.5
<i>Relative standard errors (d)</i>										
Males	%	18.0	23.0	20.3	22.1	19.8	31.4	np	np	9.2
Females	%	16.1	16.0	15.5	17.8	18.6	26.0	np	np	9.3
Total	%	11.9	13.3	13.5	13.6	13.8	20.0	17.6	np	6.7
<i>95 per cent confidence intervals</i>										
Males	±	1.1	1.3	0.8	1.0	1.4	1.5	np	np	0.5
Females	±	1.5	1.2	1.2	1.2	1.3	2.0	np	np	0.8
Total	±	0.9	0.9	0.8	0.8	1.0	1.3	1.2	np	0.5

(a) Psychological distress levels derived from the K10. Denominator includes a small number of people for whom levels of psychological distress were unable to be determined.

(b) Adults are defined as people aged 18 years and over.

(c) Rates are age standardised by State and Territory, to the June 2001 Estimated Resident Population (5 year ranges from 18 for adults).

(d) Estimates with a "*" have a relative standard error between 25 per cent and 50 per cent and should be used with caution.

np Not published.

Source: ABS (unpublished) *NHS, 2007-08*, Cat. no. 4364.0.

Table 12A.12 Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2007-08 (a), (b)

	<i>Proportion (c)</i>	<i>Relative standard error (%) (d)</i>	<i>95 % confidence interval (±)</i>
<i>Remoteness of residence</i>			
Major cities	3.6	8.0	0.6
Inner regional	3.3	11.5	0.8
Outer regional	3.0	14.7	0.9
Remote	*3.2	32.5	2.0
Very remote (e)
<i>SEIFA of residence (quintiles)</i>			
Quintile 1	6.5	9.5	1.2
Quintile 2	3.7	12.7	0.9
Quintile 3	3.3	15.1	1.0
Quintile 4	2.1	16.1	0.7
Quintile 5	2.3	19.0	0.9
<i>SEIFA of residence (deciles)</i>			
Decile 1	8.1	12.2	1.9
Decile 2	5.1	12.3	1.2
Decile 3	4.1	16.1	1.3
Decile 4	3.2	19.3	1.2
Decile 5	3.7	23.7	1.7
Decile 6	2.7	17.0	0.9
Decile 7	2.1	22.6	0.9
Decile 8	2.2	22.1	1.0
Decile 9	*2.9	25.2	1.4
Decile 10	*1.5	27.0	0.8
<i>Disability status</i>			
With disability or restrictive long-term health condition	7.3	6.4	0.9
No disability or restrictive long-term health condition	1.0	16.4	0.3
Total	3.5	6.7	0.5

SEIFA = Socio-Economic Indexes for Areas

(a) Adults are defined as people aged 18 years and over.

(b) Psychological distress levels derived from the K10. Denominator includes a small number of people for whom levels of psychological distress were unable to be determined.

(c) Rates are age standardised by State and Territory, to the June 2001 Estimated Resident Population (5 year ranges from 18 for adults).

(d) Estimate with a "*" have a relative standard error between 25 per cent and 50 per cent and should be used with caution.

(e) Very remote data were not collected in the 2007-08 NHS.

.. Not applicable.

Table 12A.12 Age-standardised rate of adults with very high levels of psychological distress, by remoteness, SEIFA IRSD quintiles, SEIFA IRSD deciles, and disability status, 2007-08 (a), (b)

	<i>Proportion (c)</i>	<i>Relative standard error (%) (d)</i>	<i>95 % confidence interval (±)</i>
--	-----------------------	--	-------------------------------------

Source: ABS (unpublished) *NHS, 2007-08*, Cat. no. 4364.0.

TABLE 12A.13

Table 12A.13 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2007-08 (a), (b)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
<i>Age standardised proportion (d)</i>										
<i>Remoteness of residence</i>										
Major cities	%	13.4	11.9	11.2	9.7	12.3	..	10.9	..	12.1
Inner regional	%	12.1	11.7	11.9	10.9	*13.3	11.6	–	..	11.9
Outer regional/remote	%	*12.2	8.5	13.0	*9.6	14.2	9.9	..	*13.4	11.8
Very remote (c)	%
<i>SEIFA of residence (quintiles)</i>										
Quintile 1	%	20.1	18.6	15.8	19.3	20.4	15.9	np	np	18.6
Quintile 2	%	13.2	14.0	12.4	9.3	13.8	8.7	np	np	12.6
Quintile 3	%	11.4	11.5	11.4	14.3	13.1	9.0	*20.5	np	11.9
Quintile 4	%	9.8	8.5	*7.8	8.2	9.0	*6.7	12.4	np	8.9
Quintile 5	%	10.1	10.0	9.5	*3.9	9.9	*9.4	7.1	*23.4	9.2
<i>Disability status</i>										
With disability or restrictive long-term health condition	%	23.4	21.0	18.7	17.9	24.8	19.9	19.4	np	21.0
No disability or restrictive long-term health condition	%	6.3	5.3	6.8	5.1	5.2	4.8	4.6	np	5.9
<i>Gender</i>										
Males	%	10.2	8.5	9.0	8.6	12.2	9.0	9.8	np	9.6
Females	%	15.4	15.0	14.0	11.4	13.8	12.5	12.0	15.1	14.4
Total	%	12.8	11.8	11.5	10.0	13.0	10.8	10.9	*13.4	12.0
<i>Relative standard errors (e)</i>										
<i>Remoteness of residence</i>										
Major cities	%	6.6	7.9	10.1	8.7	8.3	..	9.3	..	3.6
Inner regional	%	14.9	15.8	14.1	22.3	26.3	12.6	–	..	7.0

TABLE 12A.13

Table 12A.13 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2007-08 (a), (b)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
Outer regional/remote	%	26.4	24.4	12.2	27.4	19.8	14.0	..	36.8	7.3
Very remote (c)	%
<i>SEIFA of residence (quintiles)</i>										
Quintile 1	%	8.2	12.6	11.3	13.7	12.9	12.6	np	np	5.1
Quintile 2	%	15.3	14.3	11.6	16.9	18.5	16.9	np	np	7.0
Quintile 3	%	15.5	13.7	12.0	16.3	17.0	24.2	29.9	np	6.9
Quintile 4	%	13.6	17.8	25.7	17.0	22.1	28.8	15.9	np	8.6
Quintile 5	%	15.2	17.6	21.5	29.8	16.6	32.4	16.1	44.5	7.8
<i>Disability status</i>										
With disability or restrictive long-term health condition	%	6.7	7.8	9.2	8.5	8.0	11.8	9.4	np	3.9
No disability or restrictive long-term health condition	%	9.4	12.5	14.5	14.0	15.5	19.6	17.4	np	5.5
<i>Gender</i>										
Males	%	9.6	11.3	12.9	10.8	12.1	14.3	14.3	np	4.5
Females	%	7.2	8.0	7.8	9.3	9.9	14.1	10.4	18.3	4.0
Total	%	5.7	6.6	7.7	7.2	8.3	9.3	9.4	36.8	3.1
<i>95 per cent confidence intervals</i>										
<i>Remoteness of residence</i>										
Major cities	±	1.7	1.8	2.2	1.6	2.0	..	2.0	..	0.9
Inner regional	±	3.5	3.6	3.3	4.7	6.9	2.8	–	..	1.6
Outer regional/remote	±	6.3	4.1	3.1	5.2	5.5	2.7	..	9.7	1.7
Very remote (c)	±
<i>SEIFA of residence (quintiles)</i>										
Quintile 1	±	3.2	4.6	3.5	5.2	5.2	3.9	np	np	1.8

TABLE 12A.13

Table 12A.13 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, 2007-08 (a), (b)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
Quintile 2	±	4.0	3.9	2.8	3.1	5.0	2.9	np	np	1.7
Quintile 3	±	3.5	3.1	2.7	4.6	4.4	4.3	12.0	np	1.6
Quintile 4	±	2.6	3.0	3.9	2.7	3.9	3.8	3.9	np	1.5
Quintile 5	±	3.0	3.5	4.0	2.3	3.2	5.9	2.2	20.4	1.4
<i>Disability status</i>										
With disability or restrictive long-term health condition	±	3.1	3.2	3.4	3.0	3.9	4.6	3.6	np	1.6
No disability or restrictive long-term health condition	±	1.2	1.3	1.9	1.4	1.6	1.8	1.6	np	0.6
<i>Gender</i>										
Males	±	1.9	1.9	2.3	1.8	2.9	2.5	2.7	np	0.8
Females	±	2.2	2.4	2.1	2.1	2.7	3.4	2.4	5.4	1.1
Total	±	1.4	1.5	1.7	1.4	2.1	2.0	2.0	9.7	0.7

SEIFA = Socio-Economic Indexes for Areas

(a) Adults are defined as people aged 18 years and over.

(b) Psychological distress levels derived from the K10. Denominator includes a small number of people for whom levels of psychological distress were unable to be determined.

(c) Data for NT should be interpreted with caution as the *NHS 2007-08* excluded very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

(d) Rates are age standardised by State and Territory, to the June 2001 Estimated Resident Population (5 year ranges from 18 for adults).

(e) Estimate with a "*" have a relative standard error between 25 per cent and 50 per cent and should be used with caution.

.. Not applicable. – Nil or rounded to zero. **np** Not published.

Source: ABS (unpublished) *NHS 2007-08*, Cat. no. 4364.0.

TABLE 12A.14

Table 12A.14 Level of psychological distress K10, 2007-08 (per cent) (a), (b)

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (c)	Aust
<i>Males</i>									
18–64 years									
Low (10–15)	70.6 ± 3.4	72.4 ± 3.4	70.2 ± 4.2	68.3 ± 3.8	67.9 ± 4.4	75.2 ± 5.3	69.3 ± 3.6	np	70.5 ± 1.7
Moderate (16–21)	19.1 ± 2.7	18.7 ± 3.4	19.9 ± 3.5	22.8 ± 4.0	19.2 ± 2.4	15.7 ± 4.5	20.6 ± 3.2	np	19.6 ± 1.6
High (22–29) & Very high (30–50)	10.3 ± 2.1	8.9 ± 2.3	9.9 ± 2.5	8.9 ± 2.1	12.9 ± 3.5	9.1 ± 2.7	10.1 ± 2.8	np	9.9 ± 1.0
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	np	100.0
65 years or over									
Low (10–15)	77.1 ± 5.2	83.7 ± 4.8	75.2 ± 6.7	82.8 ± 7.0	74.0 ± 6.4	73.1 ± 8.5	74.2 ± 10.8	np	78.7 ± 2.8
Moderate (16–21)	12.6 ± 4.2	9.5 ± 4.1	19.3 ± 6.0	10.7* ± 6.0	16.4 ± 5.9	15.6 ± 6.7	18.5* ± 12.3	np	13.3 ± 2.4
High (22–29) & Very high (30–50)	10.3 ± 4.1	6.8 ± 2.7	5.5* ± 3.1	6.5* ± 4.5	9.6* ± 4.7	11.3* ± 6.8	7.4* ± 6.3	np	8.0 ± 1.8
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	np	100.0
Total									
Low (10–15)	71.7 ± 3.2	74.1 ± 2.8	71.0 ± 3.9	70.4 ± 3.3	69.0 ± 3.5	74.8 ± 4.8	69.8 ± 3.7	63.9 ± 17.1	71.8 ± 1.6
Moderate (16–21)	18.1 ± 2.4	17.3 ± 2.8	19.8 ± 3.2	21.1 ± 3.5	18.7 ± 2.0	15.7 ± 4.1	20.3 ± 3.6	23.9* ± 14.1	18.6 ± 1.4
High (22–29) & Very high (30–50)	10.3 ± 1.9	8.6 ± 1.9	9.2 ± 2.3	8.6 ± 1.8	12.3 ± 3.1	9.5 ± 2.7	9.8 ± 2.7	12.3* ± 12.2	9.6 ± 0.9
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
<i>Females</i>									
18–64 years									
Low (10–15)	60.9 ± 3.0	61.4 ± 3.5	58.4 ± 3.9	68.1 ± 3.3	62.1 ± 4.0	65.0 ± 4.8	61.0 ± 3.7	np	61.4 ± 1.5
Moderate (16–21)	23.7 ± 2.9	22.7 ± 2.9	26.6 ± 3.8	19.7 ± 3.2	23.0 ± 3.1	21.3 ± 4.0	27.4 ± 3.5	np	23.6 ± 1.5
High (22–29) & Very high (30–50)	15.4 ± 2.4	15.8 ± 2.4	15.1 ± 2.5	12.2 ± 2.4	14.8 ± 3.1	13.7 ± 4.0	11.6 ± 2.5	np	15.0 ± 1.2
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	np	100.0
65 years and over									
Low (10–15)	65.1 ± 5.1	70.7 ± 7.2	75.1 ± 6.2	75.4 ± 5.6	76.5 ± 5.2	70.4 ± 7.2	67.7 ± 8.4	np	70.5 ± 3.0
Moderate (16–21)	19.5 ± 5.1	18.1 ± 5.6	16.6 ± 5.2	16.5 ± 5.2	15.6 ± 4.3	22.4 ± 6.5	18.3 ± 7.4	np	18.1 ± 2.6

TABLE 12A.14

Table 12A.14 Level of psychological distress K10, 2007-08 (per cent) (a), (b)

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (c)	Aust
High (22–29) & Very high (30–50)	15.4 ± 3.7	11.2* ± 5.5	8.3 ± 3.8	8.1 ± 3.5	7.8 ± 3.4	7.2* ± 3.9	14.0 ± 6.7	np	11.5 ± 2.0
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	np	100.0
Total									
Low (10–15)	61.7 ± 2.6	63.1 ± 3.3	61.0 ± 3.7	69.3 ± 2.8	65.0 ± 3.5	66.0 ± 3.9	61.9 ± 3.4	59.0 ± 19.6	63.0 ± 1.4
Moderate (16–21)	23.0 ± 2.6	21.9 ± 2.5	25.0 ± 3.4	19.2 ± 2.6	21.6 ± 2.8	21.5 ± 3.6	26.2 ± 3.2	26.6 ± 14.5	22.7 ± 1.4
High (22–29) & Very high (30–50)	15.4 ± 2.2	15.0 ± 2.4	14.0 ± 2.1	11.5 ± 2.2	13.4 ± 2.6	12.4 ± 3.2	11.9 ± 2.4	14.4* ± 9.7	14.4 ± 1.1
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
People									
18–64 years									
Low (10–15)	65.8 ± 2.2	66.9 ± 2.2	64.2 ± 2.9	68.2 ± 2.8	65.0 ± 3.2	70.0 ± 3.4	65.1 ± 2.6	np	66.0 ± 1.1
Moderate (16–21)	21.4 ± 1.9	20.7 ± 2.1	23.3 ± 2.6	21.3 ± 2.7	21.1 ± 1.9	18.5 ± 2.6	24.0 ± 2.4	np	21.6 ± 1.1
High (22–29) & Very high (30–50)	12.8 ± 1.5	12.4 ± 1.7	12.5 ± 1.9	10.5 ± 1.7	13.8 ± 2.5	11.5 ± 2.2	10.9 ± 2.1	np	12.4 ± 0.8
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	np	100.0
65 years and over									
Low (10–15)	70.7 ± 3.7	76.7 ± 4.6	75.2 ± 4.9	78.9 ± 4.5	75.4 ± 4.0	71.6 ± 5.5	70.7 ± 6.1	np	74.3 ± 2.3
Moderate (16–21)	16.3 ± 3.5	14.1 ± 3.3	17.9 ± 4.2	13.7 ± 4.1	16.0 ± 3.4	19.3 ± 4.4	18.4 ± 6.2	np	15.9 ± 1.9
High (22–29) & Very high (30–50)	13.0 ± 3.0	9.2 ± 3.4	6.9 ± 2.7	7.3 ± 2.7	8.6 ± 3.0	9.1 ± 3.7	10.9 ± 4.8	np	9.9 ± 1.4
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	np	100.0
Total									
Low (10–15)	66.6 ± 2.0	68.5 ± 2.0	65.9 ± 2.7	69.8 ± 2.4	67.0 ± 2.6	70.3 ± 3.0	65.8 ± 2.5	61.6 ± 15.4	67.3 ± 1.0
Moderate (16–21)	20.5 ± 1.8	19.6 ± 1.8	22.4 ± 2.3	20.1 ± 2.3	20.2 ± 1.6	18.7 ± 2.2	23.3 ± 2.4	25.1 ± 12.0	20.7 ± 1.0
High (22–29) & Very high (30–50)	12.9 ± 1.4	11.9 ± 1.5	11.6 ± 1.7	10.0 ± 1.4	12.9 ± 2.1	11.0 ± 1.9	10.9 ± 2.0	13.3* ± 8.1	12.0 ± 0.7
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table 12A.14 **Level of psychological distress K10, 2007-08 (per cent) (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
--	------------	------------	------------	-----------	-----------	------------	------------	---------------	-------------

- (a) Derived from the Kessler Psychological Distress Scale–10 items (K10). This is a scale of non-specific psychological distress based on 10 questions about negative emotional states in the 4 weeks prior to interview. The K10 is scored from 10 to 50, with higher scores indicating a higher level of distress; low scores indicate a low level of distress. Scores are grouped as follows: Low 10–15, Moderate 16–21, High 22–29, and Very high 30–50.
- (b) A '*' indicates a RSE of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution. Estimates with RSEs greater than 50 per cent are considered too unreliable for general use. These estimates are not published.
- (c) Separate estimates for the NT are not available for some estimates from this survey, but the NT contributes to national estimates. Data for NT should be interpreted with caution as the *NHS 2007-08* excluded very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.
- (d) Totals include not stated.

np Not published.

Source: ABS (unpublished) *NHS 2007-08*, Cat. no. 4364.0.

TABLE 12A.15

Table 12A.15 **Age-standardised rate of adults with high/ very high levels of psychological distress, by State and Territory, by Indigenous status, 2011–13 (a), (b), (c), (d)**

	<i>unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (e)</i>	<i>Aust</i>
<i>Age standardised proportion (%)</i>										
Aboriginal and Torres Strait Islander	rate	30.5	31.5	30.3	28.5	32.8	26.3	30.9	21.6	29.4
Non-Indigenous	rate	9.9	11.3	11.5	10.9	12.2	9.9	8.9	8.2	10.8
<i>Relative standard errors</i>										
Aboriginal and Torres Strait Islander	%	8.2	7.8	7.0	5.9	7.4	10.4	16.8	8.8	3.6
Non-Indigenous	%	6.8	6.1	6.5	7.8	7.4	9.0	9.1	13.2	3.0
<i>95 per cent confidence intervals</i>										
Aboriginal and Torres Strait Islander	±	4.9	4.8	4.1	3.3	4.8	5.4	10.2	3.7	2.1
Non-Indigenous	±	1.3	1.3	1.5	1.7	1.8	1.8	1.6	2.1	0.6

(a) Levels of psychological distress are derived from the Kessler Psychological Distress Scale (K5). Denominator includes a small number of persons for whom levels of psychological distress were unable to be determined.

(b) Rates are age standardised by State and Territory, to the June 2001 Estimated Resident Population (10 year ranges from 18 years).

(c) Adults are defined as persons aged 18 years and over.

(d) Totals for Aboriginal and Torres Strait Islander people exclude a small number of persons for whom responses were provided by proxy but who were not present at interview.

(e) Data for NT should be interpreted with caution as the *AHS 2011–13 (2011-12 NHS component)* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

Source: ABS (unpublished) *Australian Aboriginal and Torres Strait Islander Health Survey, 2012-13* (National Aboriginal and Torres Strait Islander Health Survey component); ABS (unpublished) *AHS 2011–13 (2011-12 NHS component)*.

TABLE 12A.16

Table 12A.16 Level of psychological distress K10, 2004-05 (per cent) (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
Males									
18–64 years									
Low (10–15)	65.9	64.8	64.5	68.1	64.7	68.9	65.8	na	65.6
Moderate (16–21)	23.3	23.8	23.4	22.1	24.5	19.7	24.8	na	23.3
High (22–29) & Very high (30–50)	10.7	11.1	11.9	9.8	10.5	11.1	9.4	na	11.0
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
65 years and over									
Low (10–15)	71.4	73.9	66.0	80.9	76.1	74.4	65.5	na	72.4
Moderate (16–21)	17.7	15.7	19.8	13.0	18.5	15.7	25.4	na	17.2
High (22–29) & Very high (30–50)	10.9*	9.8*	14.0*	6.0	5.3*	9.9*	9.1	na	10.2
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
Total									
Low (10–15)	66.8	66.2	64.7	69.9	66.7	69.8	65.8	na	66.6
Moderate (16–21)	22.4	22.6	22.9	20.9	23.5	19.0	24.8	na	22.4
High (22–29) & Very high (30–50)	10.8	10.9	12.2	9.2	9.6	10.9	9.4	na	10.8
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
Females									
18–64 years									
Low (10–15)	58.6	55.0	58.1	63.3	58.4	63.8	55.5	na	58.1
Moderate (16–21)	26.6	28.2	25.1	21.2	26.1	21.0	29.2	na	26.0
High (22–29) & Very high (30–50)	14.6	16.5	16.8	15.4	15.5	15.3	15.3	na	15.8
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
65 years and over									
Low (10–15)	65.0	63.8	61.9	75.1	69.5	68.3	60.9	na	65.4
Moderate (16–21)	21.8	26.4	23.8	16.7	19.3	21.5	29.0	na	22.8
High (22–29) & Very high (30–50)	13.1	9.3	14.1	8.1	11.2	10.2	10.1	na	11.6

TABLE 12A.16

Table 12A.16 Level of psychological distress K10, 2004-05 (per cent) (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
Total									
Low (10–15)	59.8	56.6	58.7	65.2	60.6	64.6	56.2	na	59.4
Moderate (16–21)	25.8	27.9	24.9	20.5	24.7	21.1	29.2	na	25.5
High (22–29) & Very high (30–50)	14.4	15.2	16.4	14.3	14.6	14.3	14.6	na	15.0
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
People									
18–64 years									
Low (10–15)	62.3	59.9	61.3	65.7	61.6	66.3	60.6	na	61.8
Moderate (16–21)	25.0	26.0	24.3	21.7	25.3	20.4	27.0	na	24.7
High (22–29) & Very high (30–50)	12.7	13.8	14.4	12.6	13.0	13.2	12.4	na	13.4
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
65 years and over									
Low (10–15)	67.9	68.4	63.8	77.8	72.5	71.1	63.0	na	68.6
Moderate (16–21)	20.0	21.6	21.9	15.0	18.9	18.9	27.3	na	20.2
High (22–29) & Very high (30–50)	12.1	9.5	14.1	7.2*	8.6	10.0*	9.7	na	11.0
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0
Total									
Low (10–15)	63.2	61.3	61.6	67.5	63.6	67.2	60.9	na	62.9
Moderate (16–21)	24.1	25.3	23.9	20.7	24.1	20.1	27.0	na	24.0
High (22–29) & Very high (30–50)	12.6	13.1	14.3	11.8	12.2	12.6	12.1	na	13.0
Total (d)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	na	100.0

(a) Psychological distress as measured by the Kessler 10 scale.

(b) A '*' indicates that an estimate has a RSE of between 25 per cent and 50 per cent and should be used with caution.

(c) Separate estimates for the NT are not available for this survey, but the NT contributes to national estimates.

(d) Totals include not stated.

Table 12A.16 **Level of psychological distress K10, 2004-05 (per cent) (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (c)</i>	<i>Aust</i>
--	------------	------------	------------	-----------	-----------	------------	------------	---------------	-------------

na Not available.

Source: ABS (2006) *National Health Survey 2004-05*, Cat. no. 4362.0, Canberra.

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aust
<i>2007-08</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	28 805	23 009	16 671	7 022	7 423	1 513	1 182	317	85 942
Patient attendances (d)	546 004	585 193	327 045	114 737	162 878	36 725	18 471	3 547	1 794 600
Group psychotherapy	15 850	18 137	2 898	870	567	2 877	146	15	41 360
Interview with non-patient	1 982	1 987	1 601	439	433	126	48	18	6 634
Telepsychiatry	643	92	334	15	9	2	11	19	1 125
Case conferencing	80	763	41	42	47	38	4	–	1 015
Electroconvulsive therapy (e)	5 280	5 327	4 886	1 480	1 216	790	45	2	19 026
Total psychiatrist services	598 644	634 508	353 476	124 605	172 573	42 071	19 907	3 918	1 949 702
GP mental health specific services									
GP mental health care plans	407 865	335 835	209 549	106 349	80 756	23 307	14 934	5 095	1 183 690
Focussed psychological strategies	13 254	10 350	7 051	1 474	4 168	414	376	46	37 133
Total GP mental health specific services	421 119	346 185	216 600	107 823	84 924	23 721	15 310	5 141	1 220 823
Psychologist services									
Psychological therapy — clinical psychologists	208 032	174 404	69 774	114 269	49 556	20 361	11 577	1 404	649 377
Focussed psychological strategies — psychologists	402 284	431 801	237 281	54 998	48 985	22 620	19 001	3 699	1 220 669
Enhanced primary care — psychologists	3 056	2 213	1 731	263	356	127	28	14	7 788
Total psychologist services	613 372	608 418	308 786	169 530	98 897	43 108	30 606	5 117	1 877 834
Other allied health services									

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Focussed psychological strategies — occupational therapist	5 830	3 826	1 999	1 563	1 642	494	3	82	15 439
Focussed psychological strategies — social worker	26 594	25 519	12 119	5 505	5 052	1 571	135	375	76 870
Enhanced Primary Care — mental health worker (f)	1 045	599	323	37	375	18	3	—	2 400
Total allied health services	33 469	29 944	14 441	7 105	7 069	2 083	141	457	94 709
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	86.4	120.9	83.6	58.5	108.4	84.9	58.4	18.0	92.1
GP mental health specific services	60.8	66.0	51.2	50.6	53.3	47.8	44.9	23.6	57.6
Psychologist services	88.5	116.0	73.0	79.6	62.1	87.0	89.8	23.5	88.7
Other allied health services	4.8	5.7	3.4	3.3	4.4	4.2	0.4	2.1	4.5
<i>2008-09</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	31 484	25 495	17 220	8 055	7 418	1 785	1 266	306	93 029
Patient attendances (d)	543 800	583 020	330 605	117 929	162 032	37 344	17 961	3 831	1 796 522
Group psychotherapy	20 082	17 924	2 479	678	574	3 106	201	30	45 074
Interview with non-patient	2 848	2 594	1 948	439	552	112	73	15	8 581
Telepsychiatry	752	78	447	26	8	1	15	29	1 356
Case conferencing	190	734	97	44	37	31	9	2	1 144
Electroconvulsive therapy (e)	5 425	6 326	5 462	1 852	1 628	589	103	6	21 391
Assessment and treatment of pervasive developmental disorder	32	65	22	5	—	—	—	1	125
Total psychiatrist services	604 613	636 236	358 280	129 023	172 254	42 968	19 628	4 220	1 967 222

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
GP mental health specific services									
GP mental health care	520 403	434 383	290 904	138 410	111 352	28 783	19 020	6 688	1 549 943
Focussed psychological strategies	13 238	10 693	6 037	1 115	3 261	249	345	226	35 164
Family group therapy	6 696	6 144	1 000	274	560	161	85	16	14 936
Total GP mental health specific services	540 337	451 220	297 941	139 799	115 173	29 193	19 450	6 930	1 600 043
Psychologist services									
Psychological therapy — clinical psychologists	298 137	226 729	111 728	145 385	77 824	28 968	14 297	1 767	904 835
Focussed psychological strategies — psychologists	517 849	550 951	315 067	76 491	59 519	23 591	25 367	4 963	1 573 798
Enhanced primary care — psychologists	2 705	1 858	1 413	267	178	88	68	14	6 591
Assessment and treatment of pervasive developmental disorder	1 180	2 196	399	348	244	101	87	20	4 575
Total psychologist services	819 871	781 734	428 607	222 491	137 765	52 748	39 819	6 764	2 489 799
Other allied health services									
Focussed psychological strategies — occupational therapist	9 207	7 689	3 373	1 951	2 956	519	182	10	25 887
Focussed psychological strategies — social worker	42 707	41 722	17 111	9 107	7 860	2 451	449	133	121 540
Enhanced Primary Care — mental health worker (f)	1 059	742	298	39	169	13	—	2	2 322
Total allied health services	52 973	50 153	20 782	11 097	10 985	2 983	631	145	149 749
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	85.9	118.6	82.4	58.5	106.9	85.9	56.4	19.0	90.9
GP mental health specific services	76.7	84.1	68.5	63.4	71.4	58.4	55.9	31.3	73.9

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Psychologist services	116.4	145.7	98.5	100.9	85.5	105.4	114.5	30.5	115.0
Other allied health services	7.5	9.3	4.8	5.0	6.8	6.0	1.8	0.7	6.9
<i>2009-10</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	34 265	26 289	17 780	8 249	7 264	1 902	1 385	366	97 511
Patient attendances (d)	543 765	577 090	338 197	124 506	160 934	36 999	17 554	3 822	1 802 867
Group psychotherapy	22 013	16 144	2 504	669	563	3 190	135	21	45 239
Interview with non-patient	4 238	3 093	2 613	428	593	131	59	18	11 173
Telepsychiatry	733	117	697	29	107	8	19	9	1 719
Case conferencing	302	884	93	93	36	21	5	–	1 434
Electroconvulsive therapy (e)	5 715	6 320	6 642	2 217	1 565	720	123	24	23 326
Assessment and treatment of pervasive developmental disorder	50	69	68	np	16	np	–	–	212
Total psychiatrist services	611 081	630 006	368 594	136 206	171 078	42 976	19 280	4 260	1 983 481
GP mental health specific services									
GP mental health care	581 755	343 420	492 773	154 864	127 135	32 634	8 789	20 543	1 761 913
Focussed psychological strategies	13 609	9 101	6 078	1 289	3 135	451	285	318	34 266
Family group therapy	6 080	895	5 833	244	516	92	13	97	13 770
Total GP mental health specific services	601 444	353 416	504 684	156 397	130 786	33 177	9 087	20 958	1 809 949
Psychologist services									
Psychological therapy — clinical psychologists	343 733	277 745	146 601	168 215	97 566	33 247	17 445	2 617	1 087 169
Focussed psychological strategies — psychologists	614 418	640 812	390 393	93 016	68 990	27 300	28 131	6 143	1 869 203

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Enhanced primary care — psychologists	2 968	1 834	1 322	358	239	95	58	28	6 902
Assessment and treatment of pervasive developmental disorder	1 863	3 323	1 170	555	441	93	117	155	7 717
Total psychologist services (h)	962 998	923 714	539 486	262 144	167 236	60 735	45 751	8 959	2 971 023
Other allied health services									
Focussed psychological strategies — occupational therapist	13 062	9 474	np	3 940	2 267	1 075	259	np	34 194
Focussed psychological strategies — social worker	51 896	58 436	24 164	11 255	10 964	4 001	1 073	292	162 081
Enhanced Primary Care — mental health worker (f)	np	np	680	120	78	8	np	7	2 669
Total allied health services (h)	65 889	68 753	28 960	13 351	15 273	5 084	1 336	307	198 953
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	85.0	114.6	82.4	60.0	104.7	85.0	54.3	18.7	89.5
GP mental health specific services	83.6	64.3	112.8	68.9	80.0	65.6	25.6	92.0	81.7
Psychologist services	133.9	168.1	120.6	115.5	102.4	120.2	128.9	39.3	134.1
Other allied health services	9.2	12.5	6.5	5.9	9.3	10.1	3.8	1.3	9.0
<i>2010-11</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	35 803	27 131	19 866	8 591	7 099	1 741	1 582	312	102 125
Patient attendances (d)	557 867	576 962	344 504	124 555	154 924	35 592	18 856	3 945	1 817 205
Group psychotherapy	22 572	15 306	2 411	557	400	2 818	242	68	44 374

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Interview with non-patient	5 953	3 915	4 219	475	668	152	173	16	15 571
Telepsychiatry	941	149	1 184	127	182	18	14	18	2 633
Case conferencing	517	956	209	145	160	22	10	7	2 026
Electroconvulsive therapy (e)	12 621	13 809	15 951	4 404	4 350	2 268	275	72	53 750
Assessment and treatment of pervasive developmental disorder	55	69	54	3	12	4	1	—	198
Total psychiatrist services	636 329	638 297	388 398	138 857	167 795	42 615	21 153	4 438	2 037 882
GP mental health specific services									
GP mental health care	676 154	579 248	397 898	175 073	147 956	38 433	24 211	8 728	2 047 701
Focussed psychological strategies	17 504	10 485	8 606	1 512	3 332	716	424	326	42 905
Family group therapy	5 626	4 755	769	212	603	147	95	15	12 222
Total GP mental health specific services	699 284	594 488	407 273	176 797	151 891	39 296	24 730	9 069	2 102 828
Psychologist services									
Psychological therapy — clinical psychologists	399 144	333 786	184 361	175 818	116 009	35 023	23 066	3 043	1 270 250
Focussed psychological strategies — psychologists	694 950	693 592	445 505	111 650	73 850	36 235	28 534	6 933	2 091 249
Enhanced primary care — psychologists	2 844	1 889	1 312	430	217	125	61	9	6 887
Assessment and treatment of pervasive developmental disorder	2 065	3 626	1 367	726	414	39	144	64	8 445
Total psychologist services (h)	1 099 029	1 032 894	632 552	288 627	190 492	71 422	51 805	10 049	3 376 870
Other allied health services									
Focussed psychological strategies — occupational therapist	18 101	10 304	3 672	2 584	5 407	939	350	9	41 366

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Focussed psychological strategies — social worker	57 507	71 410	26 016	12 796	12 061	4 478	1 464	259	185 991
Enhanced Primary Care — mental health worker (f)	1 222	1 143	744	341	141	12	4	3	3 610
Total allied health services (h)	76 832	82 857	30 434	15 721	17 609	5 429	1 818	272	230 972
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	87.5	114.3	85.4	59.9	101.7	83.7	58.4	19.3	90.7
GP mental health specific services	96.2	106.4	89.5	76.3	92.0	77.2	68.3	39.5	93.6
Psychologist services	151.1	184.9	139.1	124.6	115.4	140.2	143.1	43.7	150.2
Other allied health services	10.6	14.8	6.7	6.8	10.7	10.7	5.0	1.2	10.3
<i>2011-12</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	37 346	29 634	21 864	9 406	7 124	1 651	1 536	290	108 877
Patient attendances (d)	561 520	590 523	368 265	124 548	154 032	33 233	17 079	3 465	1 852 665
Group psychotherapy	26 936	14 018	3 005	580	254	1 470	208	105	46 576
Interview with non-patient	6 079	5 614	5 411	374	895	150	174	24	18 721
Telepsychiatry	872	148	1 122	55	47	28	21	8	2 301
Case conferencing	966	1 716	378	161	159	20	15	6	3 421
Electroconvulsive therapy (e)	5 350	7 020	8 094	2 366	2 004	980	139	33	25 986
Assessment and treatment of pervasive developmental disorder	68	78	61	16	np	np	np	np	230
Total psychiatrist services	639 137	648 751	408 200	137 511	164 522	37 536	19 182	3 938	2 058 777
GP mental health specific services									
GP mental health care	699 492	605 877	417 905	167 758	150 998	39 415	25 166	9 506	2 116 117

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Focussed psychological strategies	15 866	10 090	7 387	1 428	2 709	817	266	129	38 692
Family group therapy	5 217	4 321	712	137	661	125	58	7	11 238
Electroconvulsive therapy (i)	6 964	6 987	8 406	2 753	2 094	1 084	163	32	28 483
Total GP mental health specific services	727 541	627 275	434 410	172 076	156 462	41 441	25 653	9 674	2 194 532
Clinical psychologist services									
Total clinical psychologist services	428 948	365 900	214 421	174 908	127 577	35 887	27 315	3 133	1 378 089
Other psychologist services									
Focussed psychological strategies — psychologists	677 689	673 360	442 712	111 347	76 946	36 903	24 859	7 086	2 050 902
Enhanced primary care — psychologists	4 119	2 770	1 920	578	410	104	85	42	10 028
Assessment and treatment of pervasive developmental disorder	2 642	4 659	1 660	789	509	90	132	113	10 594
Total other psychologist services (h)	684 502	680 798	446 365	112 717	77 865	37 097	25 076	7 277	2 071 697
Other allied health services									
Focussed psychological strategies — occupational therapist	17 266	10 666	4 116	2 354	6 168	770	275	32	41 647
Focussed psychological strategies — social worker	55 398	73 476	26 691	11 812	12 393	4 085	1 709	269	185 833
Enhanced Primary Care — mental health worker (f)	1 128	1 246	659	328	np	np	np	np	3 614
Total allied health services (h)	73 801	85 465	31 466	14 495	18 800	4 863	1 991	301	231 182
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	88.2	116.4	90.4	57.6	100.0	73.4	51.7	16.9	91.6

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
GP mental health specific services	100.4	112.5	96.3	72.1	95.1	81.0	69.2	41.6	97.6
Clinical psychologist services	59.2	65.6	47.5	73.3	77.6	70.1	73.7	13.5	61.3
Other psychologist services	94.4	122.1	98.9	47.2	47.3	72.5	67.6	31.3	92.1
Other allied health services	10.2	15.3	7.0	6.1	11.4	9.5	5.4	1.3	10.3
<i>2012-13</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	40 822	31 180	24 188	8 944	7 362	2 019	1 443	226	116 335
Patient attendances (d)	577 986	595 569	401 566	127 066	156 869	35 329	15 793	2 392	1 914 411
Group psychotherapy	26 746	11 591	2 224	208	281	1 942	226	np	43 319
Interview with non-patient	8 112	7 283	8 467	453	1 043	186	174	26	25 790
Telepsychiatry	698	233	1 292	60	31	np	35	np	2 365
Case conferencing	1 256	1 844	427	217	367	26	15	np	4 162
Electroconvulsive therapy (e)	6 326	8 070	8 906	2 788	1 972	921	177	np	29 241
Assessment and treatment of pervasive developmental disorder	89	60	140	np	np	np	np	np	298
Total psychiatrist services	662 042	655 834	447 217	139 745	167 927	40 426	17 870	2 834	2 136 042
GP mental health specific services									
GP mental health care	773 175	672 556	467 101	178 659	156 920	42 226	29 846	11 046	2 333 319
Focussed psychological strategies	13 650	8 818	6 595	1 255	1 825	350	153	np	32 724
Family group therapy	4 977	4 298	1 187	166	717	129	73	7	11 569
Electroconvulsive therapy (i)	7 857	8 313	8 494	3 212	1 990	809	228	np	30 983
Total GP mental health specific services	799 662	693 990	483 378	183 292	161 453	43 514	30 302	11 173	2 408 612

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Clinical psychologist services									
Total clinical psychologist services	483 570	427 987	244 465	182 566	139 446	45 195	30 079	3 302	1 558 063
Other psychologist services									
Focussed psychological strategies — psychologists	734 906	696 416	456 268	112 805	86 778	34 537	27 443	6 878	2 158 106
Enhanced primary care — psychologists	4 219	2 442	1 802	641	219	75	251	np	9 670
Assessment and treatment of pervasive developmental disorder	2 371	5 278	1 602	942	611	114	137	69	11 145
Total other psychologist services (h)	741 606	704 157	459 732	114 388	87 608	34 730	27 871	6 969	2 179 161
Other allied health services									
Focussed psychological strategies — occupational therapist	19 827	12 263	4 988	2 538	7 205	828	434	np	48 123
Focussed psychological strategies — social worker	54 615	80 110	30 181	13 386	17 178	4 289	2 160	258	202 280
Enhanced Primary Care — mental health worker (f)	938	1 364	1 598	381	204	np	np	np	4 513
Total allied health services (h)	75 385	93 793	36 864	16 325	24 590	5 130	2 610	302	255 129
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	90.1	115.5	97.0	56.5	101.0	78.9	47.1	12.0	93.3
GP mental health specific services	108.8	122.2	104.8	74.1	97.1	84.9	79.8	47.2	105.2
Clinical psychologist services	65.8	75.4	53.0	73.8	83.9	88.2	79.2	13.9	68.0
Other psychologist services	100.9	124.0	99.7	46.3	52.7	67.8	73.4	29.4	95.1
Other allied health services	10.3	16.5	8.0	6.6	14.8	10.0	6.9	1.3	11.1

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>2013-14</i>									
<i>Number of services</i>									
Psychiatrist services									
Initial consultations new patient (c)	42 782	33 646	27 072	9 800	6 944	2 082	1 379	252	123 955
Patient attendances (d)	594 535	615 641	432 841	131 440	148 278	36 828	16 030	2 451	1 978 044
Group psychotherapy	27 858	10 742	2 321	269	277	2 448	423	86	44 424
Interview with non-patient	11 995	7 875	10 844	580	1 166	174	222	39	32 895
Telepsychiatry	503	203	949	73	55	12	29	29	1 853
Case conferencing	2 146	2 287	541	241	238	31	37	10	5 532
Electroconvulsive therapy (e)	6 201	8 040	9 131	3 150	2 305	779	212	29	29 847
Assessment and treatment of pervasive developmental disorder	96	50	142	1	8	np	np	1	298
Total psychiatrist services	686 117	678 483	483 841	145 554	159 271	42 354	18 331	2 897	2 216 848
GP mental health specific services									
GP mental health care	848 618	740 984	522 286	206 941	167 467	45 243	33 549	11 524	2 576 612
Focussed psychological strategies	13 033	9 465	6 124	867	1 407	346	170	43	31 455
Family group therapy	5 295	3 894	1 318	123	634	94	71	5	11 434
Electroconvulsive therapy (i)	8 404	8 436	9 080	2 906	2 279	852	315	48	32 320
Total GP mental health specific services	875 350	762 778	538 809	210 837	171 787	46 535	34 105	11 620	2 651 821
Clinical psychologist services	525 968	472 076	290 515	184 529	155 356	49 396	30 870	3 309	1 712 018
Total clinical psychologist services	525 968	472 076	290 515	184 529	155 356	49 396	30 870	3 309	1 712 018
Other psychologist services									
Focussed psychological strategies — psychologists	710 516	681 143	486 731	122 368	91 384	33 715	29 157	6 821	2 161 834

TABLE 12A.17

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aust
Enhanced primary care — psychologists	6 454	4 912	3 437	974	495	117	92	88	16 568
Assessment and treatment of pervasive developmental disorder	2 660	5 099	1 828	800	700	163	102	37	11 390
Total other psychologist services (h)	719 800	691 210	492 217	124 144	92 585	34 007	29 355	6 958	2 190 276
Other allied health services									
Focussed psychological strategies — occupational therapist	19 406	13 370	6 200	2 903	6 027	752	580	52	49 290
Focussed psychological strategies — social worker	55 617	88 854	37 035	14 648	18 348	3 405	2 441	575	220 923
Enhanced Primary Care — mental health worker (f)	1 699	1 719	2 332	154	217	66	8	12	6 208
Total allied health services (h)	76 951	103 963	45 755	17 720	24 592	4 223	3 028	639	276 870
<i>Rate per 1000 people (g)</i>									
Psychiatrist services	91.9	117.2	103.1	57.1	95.0	82.4	47.7	11.9	95.1
GP mental health specific services	117.3	131.7	114.9	82.7	102.4	90.5	88.8	47.9	113.7
Clinical psychologist services	70.5	81.5	61.9	72.3	92.6	96.1	80.4	13.6	73.4
Other psychologist services	96.4	119.4	104.9	48.7	55.2	66.2	76.4	28.7	93.9
Other allied health services	10.3	18.0	9.8	6.9	14.7	8.2	7.9	2.6	11.9

(a) State and territory is based on the postcode of the mailing address of the patient as recorded by Medicare Australia. Provider type is based on the MBS item numbers claimed.

(b) A listing of the MBS items associated with each of the categories is available in the Medicare Benefits Schedule and General practice data source sections of the *Mental Health Services in Australia* (various issues), (<http://mhsa.aihw.gov.au/home/>).

(c) Includes consultations in consulting room, hospital and home visits.

(d) Includes attendances in consulting room, hospital and other locations.

(e) Data for electroconvulsive therapy may include services provided by medical practitioners other than psychiatrists.

Table 12A.17 **Mental health care specific MBS items processed (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
(f) Mental health workers include psychologists, mental health nurses, occupational therapists, social workers and Aboriginal health workers.									
(g) Crude rates based on the preliminary Australian estimated resident population as at 31 December mid-point of financial year.									
(h) Totals for psychologist/other psychologist and other allied health services include specific services for Aboriginal and Torres Strait Islander Australians that were introduced on 1 November 2008.									
(i) This item is for the initiation of management of anaesthesia for electroconvulsive therapy and includes data for services provided by medical practitioners other than GPs.									

– Nil or rounded to zero. **np** Not published.

Source: AIHW (various issues) *Mental Health Services in Australia* (various years) (available at <http://mhsa.aihw.gov.au/home/>).

TABLE 12A.18

Table 12A.18 **GP mental health-related encounters (general and mental health specific) (a)**

	2006–07	2007–08	2008–09	2009–10	2010–11	2011–12	2012–13	2013–14
<i>Total GP encounters that are mental health-related (per cent)</i>	10.4	10.8	11.7	11.4	11.7	12.1	12.3	12.8
<i>Estimated number of mental health-related encounters (b)</i>	10 713 000	11 862 000	13 202 000	13 283 000	13 931 000	14 956 000	15 842 000	17 131 180
<i>Lower 95% confidence limit</i>	10 261 000	11 280 000	12 661 000	12 714 000	13 353 000	14 250 000	15 187 000	16 355 184
<i>Upper 95% confidence limit</i>	11 165 000	12 375 000	13 678 000	13 881 000	14 426 000	15 614 000	16 474 000	17 907 070
<i>Estimated number of mental health-related encounters per 1000 population (b), (c)</i>	519.4	564.4	614.8	607.5	628.3	664.3	691.6	735.0
<i>Lower 95% confidence limit</i>	497.5	536.7	589.6	581.5	602.2	633.0	663.0	701.0
<i>Upper 95% confidence limit</i>	541.3	588.9	636.9	634.8	650.6	693.6	719.2	768.0

(a) The confidence intervals show that the difference between some of the years is not statistically significant.

(b) The estimated number of encounters is based on the proportion of encounters in the BEACH survey of general practice activity that are mental health-related, multiplied by the total number of Medicare services for non-Referred (GP) Attendances (excluding practice nurse items) as reported by the Department of Human Services (see Mental Health Services in Australia for more details).

(c) Crude rate is based on the Australian estimated resident population as at 31 December of the reference year.

Source: AIHW (2015) *Mental Health Services in Australia* (available at <http://mhsa.aihw.gov.au/home/>).

Table 12A.19 GP mental health-related encounters (general and mental health specific), by patient demographics, 2013-14

	<i>Per cent of total mental health-related encounters (a)</i>	<i>Rate (per 100 encounters)</i>	<i>95% LCL</i>	<i>95% UCL</i>	<i>Encounters per 1000 population (b)</i>
<i>Age group</i>					
Less than 15 years	3.9	4.5	4.0	5.0	153.1
15–24 years	8.0	13.3	12.3	14.3	436.7
25–34 years	12.9	16.2	14.8	17.6	645.6
35–44 years	16.9	19.3	17.9	20.8	896.3
45–54 years	17.3	17.4	16.3	18.5	955.3
55–64 years	15.4	13.7	12.8	14.7	989.5
65 years or over	25.6	10.1	9.5	10.8	1 288.4
<i>Sex</i>					
Male	41.3	12.3	11.5	13.2	607.8
Female	58.7	13.3	12.7	13.8	858.0
<i>Indigenous status (b), (c)</i>					
Aboriginal and Torres Strait Islander	2.0	15.8	13.1	18.6	605.7
Non-Indigenous	98.0	12.9	12.3	13.5	647.7
<i>Remoteness area of usual residence</i>					
Major cities	71.0	13.2	12.5	14.0	734.3
Inner regional	20.4	12.7	12.0	13.5	817.0
Outer regional	8.1	12.9	11.7	14.2	663.4
Remote and very remote	0.5	9.3	7.5	11.1	162.7
Total	100.0	12.8	12.3	13.4	739.3

LCL—lower confidence limit; UCL—upper confidence limit.

- (a) The percentages shown do not include those encounters for which the demographic information was missing or not reported.
- (b) Estimated encounters are crude rates, with the exception of Indigenous status, which is an age standardised rate (directly age-standardised using the June 2001 estimated resident population — Australian standard population).
- (c) Information on this variable was missing or not reported for more than 5 per cent of encounters.

Source: AIHW (2015) *Mental Health Services in Australia* (available at <http://mhsa.aihw.gov.au/home/>).

TABLE 12A.20

Table 12A.20 **The 10 most frequent GP managed mental health-related problems, by gender, 2013-14**

	<i>Per cent of total mental health-related encounters (a)</i>	<i>Rate (per 100 encounters)</i>	<i>95% LCL</i>	<i>95% UCL</i>
<i>Males</i>				
Depression	27.2	3.6	3.3	3.9
Anxiety	13.6	1.8	1.6	2.0
Sleep disturbance	11.6	1.5	1.4	1.7
Acute stress reaction	3.7	0.5	0.4	0.6
Tobacco abuse	5.8	0.8	0.6	0.9
Drug abuse	7.6	1.0	0.3	1.7
Dementia	3.7	0.5	0.4	0.6
Schizophrenia	5.5	0.7	0.6	0.8
Affective psychosis	2.1	0.3	0.2	0.4
Alcohol abuse	4.3	0.6	0.5	0.7
Other	14.8	2.0	1.8	2.2
Total (a)	100.0	13.2	12.3	14.2
<i>Females</i>				
Depression	34.6	4.9	4.6	5.1
Anxiety	18.5	2.6	2.4	2.8
Sleep disturbance	11.1	1.6	1.4	1.7
Acute stress reaction	7.0	1.0	0.9	1.1
Tobacco abuse	4.1	0.6	0.5	0.7
Drug abuse	2.6	0.4	0.2	0.5
Dementia	4.5	0.6	0.5	0.8
Schizophrenia	2.1	0.3	0.2	0.3
Affective psychosis	2.8	0.4	0.3	0.4
Alcohol abuse	1.3	0.2	0.1	0.2
Other	11.5	1.6	1.5	1.8
Total (a)	100.0	14.0	13.4	14.6
<i>Total</i>				
Depression	31.5	4.3	4.1	4.5
Anxiety	16.5	2.2	2.1	2.4
Sleep disturbance	11.3	1.5	1.4	1.7
Acute stress reaction	5.6	0.8	0.7	0.9
Tobacco abuse	4.8	0.7	0.6	0.7

Table 12A.20 **The 10 most frequent GP managed mental health-related problems, by gender, 2013-14**

	<i>Per cent of total mental health-related encounters (a)</i>	<i>Rate (per 100 encounters)</i>	<i>95% LCL</i>	<i>95% UCL</i>
Drug abuse	4.7	0.6	0.3	1.0
Dementia	4.2	0.6	0.4	0.7
Schizophrenia	3.5	0.5	0.4	0.5
Affective psychosis	2.6	0.3	0.3	0.4
Alcohol abuse	2.5	0.3	0.3	0.4
Other	12.9	1.8	1.6	1.9
Total (a)	100.0	13.7	13.0	14.3

LCL—lower confidence limit; UCL—upper confidence limit.

(a) Totals may not add due to rounding.

Source: AIHW (2015) *Mental Health Services in Australia* (available at <http://mhsa.aihw.gov.au/home/>).

TABLE 12A.21

Table 12A.21 **Mental health patient days (a), (b), (c)**

	NSW (d), (e)	Vic	Qld (f)	WA (g)	SA	Tas	ACT (h)	NT (h)	Aust
<i>Patient days</i>									
Admitted patient — acute units									
2005-06	468 925	325 855	216 029	167 257	117 148	30 681	15 342	11 266	1 352 503
2006-07	502 521	328 817	216 505	165 365	120 755	28 219	16 419	11 854	1 390 455
2007-08	501 388	322 087	222 006	183 741	119 808	30 924	18 539	10 990	1 409 483
2008-09	525 512	334 711	224 395	181 426	115 412	31 291	19 884	11 517	1 444 148
2009-10	531 649	332 677	226 762	182 647	114 605	29 615	21 484	10 877	1 450 316
2010-11	536 310	345 369	228 406	177 733	117 123	29 249	22 941	11 518	1 468 649
2011-12	541 039	343 809	230 274	188 644	115 761	32 148	23 163	10 489	1 485 327
2012-13	555 369	342 192	227 282	192 445	109 927	28 749	26 097	12 943	1 495 004
2013-14	565 865	349 444	230 097	195 461	111 515	27 773	23 068	11 832	1 515 055
Admitted patient — non-acute units									
2005-06	256 893	55 745	225 242	44 800	90 200	9 074	681 954
2006-07	252 391	56 837	222 783	50 751	84 637	9 482	676 881
2007-08	279 349	63 428	219 026	36 838	77 836	7 128	683 605
2008-09	265 820	54 667	215 715	38 357	65 509	9 125	649 193
2009-10	285 494	53 712	213 343	40 061	59 746	8 531	660 887
2010-11	287 011	54 293	216 365	51 600	56 073	9 779	675 121
2011-12	286 731	51 032	209 993	47 013	46 036	10 011	650 816
2012-13	290 218	52 982	218 517	44 201	43 626	7 843	657 387
2013-14	288 676	50 796	190 666	40 745	44 084	9 618	624 585
24-hour staffed community residential									
2005-06	73 112	321 675	..	11 380	8 635	34 155	13 981	..	462 938
2006-07	73 773	338 377	..	12 006	9 232	34 697	14 023	..	482 108
2007-08	42 051	352 741	..	14 888	15 277	27 194	13 599	1 737	467 487
2008-09	37 375	344 623	..	24 725	20 649	28 727	14 262	3 550	473 911

TABLE 12A.21

Table 12A.21 **Mental health patient days (a), (b), (c)**

	NSW (d), (e)	Vic	Qld (f)	WA (g)	SA	Tas	ACT (h)	NT (h)	Aust
2009-10	35 355	351 719	..	33 008	20 187	30 172	15 416	3 841	489 698
2010-11	34 503	353 996	..	17 605	22 529	29 958	14 961	4 144	477 696
2011-12	38 742	363 985	..	30 073	34 397	27 333	15 367	4 828	514 725
2012-13	38 328	364 505	..	30 459	40 158	22 777	16 045	5 508	517 780
2013-14	36 426	366 237	..	29 229	43 332	27 993	17 155	7 056	527 428
<i>Patient days per 1000 people</i>									
Admitted patient — acute units									
2005-06	69.8	64.9	54.5	82.4	75.8	62.9	46.0	54.3	66.6
2006-07	74.1	64.4	53.4	79.6	77.3	57.4	48.5	56.2	67.4
2007-08	72.8	61.9	53.4	86.1	75.9	62.4	53.9	50.7	67.1
2008-09	75.1	63.0	52.5	82.1	72.2	62.4	56.6	51.8	67.2
2009-10	74.9	61.4	51.9	80.7	70.8	58.5	60.0	47.8	66.3
2010-11	74.7	62.8	51.5	76.6	71.7	57.3	62.9	50.0	66.2
2011-12	74.7	61.7	51.0	79.0	70.4	62.8	62.5	45.1	66.1
2012-13	75.6	60.2	49.3	77.8	66.1	56.1	68.8	54.6	65.3
2013-14	75.8	60.3	49.1	76.6	66.5	54.0	60.0	48.8	65.0
Admitted patient — non-acute units									
2005-06	38.2	11.1	56.8	22.1	58.4	18.6	33.6
2006-07	37.2	11.1	54.9	24.4	54.2	19.3	32.8
2007-08	40.6	12.2	52.7	17.3	49.3	14.4	32.5
2008-09	38.0	10.3	50.5	17.4	41.0	18.2	30.2
2009-10	40.2	9.9	48.8	17.7	36.9	16.8	30.2
2010-11	40.0	9.9	48.8	22.3	34.3	19.2	30.4
2011-12	39.6	9.2	46.5	19.7	28.0	19.6	28.9
2012-13	39.5	9.3	47.4	17.9	26.2	15.3	28.7
2013-14	38.7	8.8	40.6	16.0	26.3	18.7	26.8

Table 12A.21 **Mental health patient days (a), (b), (c)**

	NSW (d), (e)	Vic	Qld (f)	WA (g)	SA	Tas	ACT (h)	NT (h)	Aust
24-hour staffed community residential									
2005-06	10.9	64.0	..	5.6	5.6	70.0	41.9	..	22.8
2006-07	10.9	66.3	..	5.8	5.9	70.6	41.4	..	23.4
2007-08	6.1	67.8	..	7.0	9.7	54.8	39.5	8.0	22.2
2008-09	5.3	64.9	..	11.2	12.9	57.3	40.6	16.0	22.1
2009-10	5.0	64.9	..	14.6	12.5	59.6	43.1	16.9	22.4
2010-11	4.8	64.4	..	7.6	13.8	58.7	41.0	18.0	21.5
2011-12	5.3	65.3	..	12.6	20.9	53.4	41.5	20.8	22.9
2012-13	5.2	64.2	..	12.3	24.2	44.4	42.3	23.3	22.6
2013-14	4.9	63.2	..	11.5	25.8	54.5	44.7	29.1	22.6

- (a) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of patient day estimates.
- (b) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (c) Hospital patient days include those provided in services funded by government, but managed and operated by private and non-government entities.
- (d) Caution is required when interpreting NSW data. Seven residential mental health services in 2006–07 were reclassified as non-acute older person specialised hospital services in 2007–08, reflecting a change in function of those units.
- (e) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.
- (f) Queensland does not fund community residential services; however, it funds a number of campus based and non-campus based extended treatment services. Data from these services are included as non-acute units.
- (g) Caution is required when interpreting WA data. Several residential services that reported as 24-hour staffed services in 2009-10 transitioned to a non-24-hour staffed model of care as of 1 July 2010. In addition, a review of services resulted in the reclassification of beds between the acute and non-acute categories for the 2010–11 collection, to more accurately reflect the function of these services.
- (h) The ACT and the NT do not have non-acute hospital units.
.. Not applicable.

Source: AIHW (unpublished) MHE NMDS; table 12A.99.

Table 12A.22 Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)

ICD-10		Public	Public	Total	Per cent
		acute hospitals	psychiatric hospitals		
		no.	no.	no.	%
<i>2005-06</i>					
F00-F03	Dementia	609	188	797	0.9
F04-F09	Other organic mental disorders	599	146	745	0.8
F10	Mental and behavioural disorders due to use of alcohol	1 623	542	2 165	2.4
F11-F19	Mental and behavioural disorders due to other psychoactive substances use	3 464	878	4 342	4.9
F20	Schizophrenia	17 402	3 231	20 633	23.1
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 505	260	1 765	2.0
F22	Persistent delusional disorders	787	163	950	1.1
F23	Acute and transient psychotic disorders	1 309	217	1 526	1.7
F25	Schizoaffective disorders	5 078	1 028	6 106	6.8
F30	Manic episode	449	71	520	0.6
F31	Bipolar affective disorders	7 331	1 157	8 488	9.5
F32	Depressive episode	10 844	1 068	11 912	13.3
F33	Recurrent depressive disorders	3 761	251	4 012	4.5
F34	Persistent mood (affective) disorders	910	109	1 019	1.1
F38, F39	Other and unspecified mood (affective) disorders	143	41	184	0.2
F40	Phobic anxiety disorders	62	14	76	0.1
F41	Other anxiety disorders	994	57	1 051	1.2
F42	Obsessive-compulsive disorders	239	22	261	0.3
F43	Reaction to severe stress and adjustment disorders	7 232	1 402	8 634	9.7
F44	Dissociative (conversion) disorders	124	13	137	0.2
F45, F48	Somatoform and other neurotic disorders	79	10	89	0.1
F50	Eating disorders	604	15	619	0.7
F51-F59	Other behavioural syndromes associated with physiological disturbances and physical factors	169	24	193	0.2
F60	Specific personality disorders	3 642	542	4 184	4.7
F61-F69	Disorders of adult personality and behaviour	189	45	234	0.3
F70-F79	Mental retardation	139	53	192	0.2

TABLE 12A.22

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F80–F89	Disorders of psychological development	168	31	199	0.2
F90	Hyperkinetic disorders	114	11	125	0.1
F91	Conduct disorders	291	53	344	0.4
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	170	61	231	0.3
F99	Mental disorder not otherwise specified	251	22	273	0.3
G30	Alzheimer's disease	509	134	643	0.7
	Other factors related to mental and behavioural disorders and substance use (b)	224	357	581	0.7
	Other specified mental health-related principal diagnosis (c)	209	17	226	0.3
	Other (d)	4 796	1 022	5 818	6.5
	Total	76 019	13 255	89 274	100.0
<i>2006-07</i>					
F00–F03	Dementia	557	178	735	0.8
F04–F09	Other organic mental disorders	569	133	702	0.8
F10	Mental and behavioural disorders due to use of alcohol	1 980	621	2 601	2.8
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	3 606	981	4 587	5.0
F20	Schizophrenia	17 610	3 014	20 624	22.3
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 456	248	1 704	1.8
F22	Persistent delusional disorders	776	130	906	1.0
F23	Acute and transient psychotic disorders	1 395	211	1 606	1.7
F25	Schizoaffective disorders	5 359	1 021	6 380	6.9
F30	Manic episode	559	69	628	0.7
F31	Bipolar affective disorders	7 935	1 089	9 024	9.8
F32	Depressive episode	11 103	1 065	12 168	13.2
F33	Recurrent depressive disorders	3 701	314	4 015	4.3
F34	Persistent mood (affective) disorders	998	118	1 116	1.2
F38, F39	Other and unspecified mood (affective) disorders	133	30	163	0.2
F40	Phobic anxiety disorders	54	6	60	0.1
F41	Other anxiety disorders	1 160	102	1 262	1.4

TABLE 12A.22

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F42	Obsessive-compulsive disorders	226	24	250	0.3
F43	Reaction to severe stress and adjustment disorders	8 141	1 274	9 415	10.2
F44	Dissociative (conversion) disorders	116	8	124	0.1
F45, F48	Somatoform and other neurotic disorders	81	8	89	0.1
F50	Eating disorders	575	7	582	0.6
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	193	12	205	0.2
F60	Specific personality disorders	3 744	531	4 275	4.6
F61–F69	Disorders of adult personality and behaviour	163	33	196	0.2
F70–F79	Mental retardation	156	44	200	0.2
F80–F89	Disorders of psychological development	175	31	206	0.2
F90	Hyperkinetic disorders	112	9	121	0.1
F91	Conduct disorders	298	32	330	0.4
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	190	58	248	0.3
F99	Mental disorder not otherwise specified	267	86	353	0.4
G30	Alzheimer's disease	497	85	582	0.6
	Other factors related to mental and behavioural disorders and substance use (b)	218	324	542	0.6
	Other specified mental health-related principal diagnosis (c)	235	36	271	0.3
	Other (d)	5 400	839	6 239	6.7
	Total	79 738	12 771	92 509	100.0
<i>2007-08</i>					
F00–F03	Dementia	592	221	813	0.9
F04–F09	Other organic mental disorders	596	172	768	0.8
F10	Mental and behavioural disorders due to use of alcohol	2 128	690	2 818	3.1
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	3 155	779	3 934	4.3
F20	Schizophrenia	17 250	2 834	20 084	21.9
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 609	260	1 869	2.0

Table 12A.22 Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F22	Persistent delusional disorders	817	136	953	1.0
F23	Acute and transient psychotic disorders	1 432	168	1 600	1.7
F25	Schizoaffective disorders	5 354	949	6 303	6.9
F30	Manic episode	532	60	592	0.6
F31	Bipolar affective disorders	7 628	1 157	8 785	9.6
F32	Depressive episode	11 051	1 121	12 172	13.3
F33	Recurrent depressive disorders	2 997	554	3 551	3.9
F34	Persistent mood (affective) disorders	938	116	1 054	1.2
F38, F39	Other and unspecified mood (affective) disorders	145	25	170	0.2
F40	Phobic anxiety disorders	79	11	90	0.1
F41	Other anxiety disorders	1 089	99	1 188	1.3
F42	Obsessive-compulsive disorders	236	19	255	0.3
F43	Reaction to severe stress and adjustment disorders	8 501	1 098	9 599	10.5
F44	Dissociative (conversion) disorders	112	11	123	0.1
F45, F48	Somatoform and other neurotic disorders	106	8	114	0.1
F50	Eating disorders	523	6	529	0.6
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	155	9	164	0.2
F60	Specific personality disorders	3 834	614	4 448	4.9
F61–F69	Disorders of adult personality and behav	197	73	270	0.3
F70–F79	Mental retardation	147	56	203	0.2
F80–F89	Disorders of psychological development	199	42	241	0.3
F90	Hyperkinetic disorders	106	17	123	0.1
F91	Conduct disorders	262	29	291	0.3
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	172	58	230	0.3
F99	Mental disorder not otherwise specified	167	101	268	0.3
G30	Alzheimer's disease	491	150	641	0.7
	Other factors related to mental and behavioural disorders and substance use (b)	191	247	438	0.5

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
	Other specified mental health-related principal diagnosis (c)	296	10	306	0.3
	Other (d)	5 832	823	6 655	7.3
	Total	78 919	12 723	91 642	100.0
<i>2008-09</i>					
F00–F03	Dementia	565	163	728	0.7
F04–F09	Other organic mental disorders	600	101	701	0.7
F10	Mental and behavioural disorders due to use of alcohol	2 365	572	2 937	3.0
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	2 827	558	3 385	3.4
F20	Schizophrenia	18 127	2 270	20 397	20.7
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 966	174	2 140	2.2
F22	Persistent delusional disorders	803	108	911	0.9
F23	Acute and transient psychotic disorders	1 338	137	1 475	1.5
F25	Schizoaffective disorders	6 239	733	6 972	7.1
F30	Manic episode	577	51	628	0.6
F31	Bipolar affective disorders	8 622	1 080	9 702	9.9
F32	Depressive episode	14 406	1 105	15 511	15.8
F33	Recurrent depressive disorders	3 433	342	3 775	3.8
F34	Persistent mood (affective) disorders	821	93	914	0.9
F38, F39	Other and unspecified mood (affective) disorders	117	24	141	0.1
F40	Phobic anxiety disorders	65	7	72	0.1
F41	Other anxiety disorders	1 386	107	1 493	1.5
F42	Obsessive-compulsive disorders	210	15	225	0.2
F43	Reaction to severe stress and adjustment disorders	8 863	931	9 794	10.0
F44	Dissociative (conversion) disorders	108	7	115	0.1
F45, F48	Somatoform and other neurotic disorders	73	10	83	0.1
F50	Eating disorders	635	6	641	0.7
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	180	8	188	0.2
F60	Specific personality disorders	3 979	550	4 529	4.6

TABLE 12A.22

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F61–F69	Disorders of adult personality and behaviour	211	58	269	0.3
F70–F79	Mental retardation	190	np	190	0.2
F80–F89	Disorders of psychological development	236	28	264	0.3
F90	Hyperkinetic disorders	85	–	85	0.1
F91	Conduct disorders	311	np	311	0.3
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	381	25	406	0.4
F99	Mental disorder not otherwise specified	189	64	253	0.3
G30	Alzheimer's disease	452	100	552	0.6
	Other factors related to mental and behavioural disorders and substance use (b)	235	np	235	0.2
	Other specified mental health-related principal diagnosis (c)	349	11	360	0.4
	Other (d)	6 853	1 047	7 900	8.0
	Total	87 797	10 562	98 359	100.0
<i>2009-10</i>					
F00–F03	Dementia	534	126	660	0.7
F04–F09	Other organic mental disorders	645	119	764	0.8
F10	Mental and behavioural disorders due to use of alcohol	2 235	560	2 795	3.1
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	2 626	530	3 156	3.4
F20	Schizophrenia	17 155	2 436	19 591	21.4
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 707	221	1 928	2.1
F22	Persistent delusional disorders	770	79	849	0.9
F23	Acute and transient psychotic disorders	1 303	145	1 448	1.6
F25	Schizoaffective disorders	5 376	750	6 126	6.7
F30	Manic episode	511	51	562	0.6
F31	Bipolar affective disorders	7 726	976	8 702	9.5
F32	Depressive episode	11 932	1 139	13 071	14.3
F33	Recurrent depressive disorders	2 631	348	2 979	3.3
F34	Persistent mood (affective) disorders	790	72	862	0.9

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F38, F39	Other and unspecified mood (affective) disorders	131	20	151	0.2
F40	Phobic anxiety disorders	71	10	81	0.1
F41	Other anxiety disorders	1 442	131	1 573	1.7
F42	Obsessive-compulsive disorders	230	23	253	0.3
F43	Reaction to severe stress and adjustment disorders	8 528	964	9 492	10.4
F44	Dissociative (conversion) disorders	128	13	141	0.2
F45, F48	Somatoform and other neurotic disorders	69	7	76	0.1
F50	Eating disorders	576	9	585	0.6
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	158	10	168	0.2
F60	Specific personality disorders	3 599	578	4 177	4.6
F61–F69	Disorders of adult personality and behaviour	171	31	202	0.2
F70–F79	Mental retardation	144	51	195	0.2
F80–F89	Disorders of psychological development	243	38	281	0.3
F90	Hyperkinetic disorders	80	19	99	0.1
F91	Conduct disorders	331	49	380	0.4
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	352	21	373	0.4
F99	Mental disorder not otherwise specified	199	81	280	0.3
G30	Alzheimer's disease	518	88	606	0.7
	Other factors related to mental and behavioural disorders and substance use (b)	227	232	459	0.5
	Other specified mental health-related principal diagnosis (c)	364	7	371	0.4
	Other (d)	7 004	1 063	8 067	8.8
	Total	80 506	10 997	91 503	100.0
<i>2010-11</i>					
F00–F03	Dementia	443	61	504	0.5
F04–F09	Other organic mental disorders	618	90	708	0.7
F10	Mental and behavioural disorders due to use of alcohol	2 318	487	2 805	2.9

Table 12A.22 Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	3 517	600	4 117	4.3
F20	Schizophrenia	18 164	2 137	20 301	21.1
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 978	202	2 180	2.3
F22	Persistent delusional disorders	802	97	899	0.9
F23	Acute and transient psychotic disorders	1 318	99	1 417	1.5
F25	Schizoaffective disorders	6 031	792	6 823	7.1
F30	Manic episode	625	47	672	0.7
F31	Bipolar affective disorders	8 147	896	9 043	9.4
F32	Depressive episode	11 874	917	12 791	13.3
F33	Recurrent depressive disorders	2 625	170	2 795	2.9
F34	Persistent mood (affective) disorders	752	69	821	0.9
F38, F39	Other and unspecified mood (affective) disorders	165	13	178	0.2
F40	Phobic anxiety disorders	72	9	81	0.1
F41	Other anxiety disorders	1 612	67	1 679	1.7
F42	Obsessive-compulsive disorders	249	10	259	0.3
F43	Reaction to severe stress and adjustment disorders	9 446	928	10 374	10.8
F44	Dissociative (conversion) disorders	149	4	153	0.2
F45, F48	Somatoform and other neurotic disorders	96	2	98	0.1
F50	Eating disorders	616	11	627	0.7
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	174	16	190	0.2
F60	Specific personality disorders	4 146	420	4 566	4.7
F61–F69	Disorders of adult personality and behaviour	162	23	185	0.2
F70–F79	Mental retardation	177	30	207	0.2
F80–F89	Disorders of psychological development	243	23	266	0.3
F90	Hyperkinetic disorders	75	3	78	0.1
F91	Conduct disorders	396	10	406	0.4
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	393	8	401	0.4

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F99	Mental disorder not otherwise specified	352	–	352	0.4
G30	Alzheimer's disease	511	51	562	0.6
	Other factors related to mental and behavioural disorders and substance use (b)	199	70	269	0.3
	Other specified mental health-related principal diagnosis (c)	271	3	274	0.3
	Other (d)	7 953	1 196	9 149	9.5
	Total	86 669	9 561	96 230	100.0
<i>2011-12</i>					
F00–F03	Dementia	446	72	518	0.5
F04–F09	Other organic mental disorders	584	93	677	0.7
F10	Mental and behavioural disorders due to use of alcohol	2 219	391	2 610	2.6
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	4 337	610	4 947	5.0
F20	Schizophrenia	18 190	1 873	20 063	20.1
F21, F24, F28, F29	Schizotypal and other delusional disorders	2 184	232	2 416	2.4
F22	Persistent delusional disorders	856	54	910	0.9
F23	Acute and transient psychotic disorders	1 378	133	1 511	1.5
F25	Schizoaffective disorders	6 137	705	6 842	6.9
F30	Manic episode	627	52	679	0.7
F31	Bipolar affective disorders	8 349	772	9 121	9.1
F32	Depressive episode	12 586	875	13 461	13.5
F33	Recurrent depressive disorders	2 509	135	2 644	2.6
F34	Persistent mood (affective) disorders	781	41	822	0.8
F38, F39	Other and unspecified mood (affective) disorders	256	14	270	0.3
F40	Phobic anxiety disorders	75	2	77	0.1
F41	Other anxiety disorders	1 792	96	1 888	1.9
F42	Obsessive-compulsive disorders	306	18	324	0.3
F43	Reaction to severe stress and adjustment disorders	10 180	911	11 091	11.1
F44	Dissociative (conversion) disorders	195	13	208	0.2
F45, F48	Somatoform and other neurotic disorders	83	7	90	0.1

TABLE 12A.22

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F50	Eating disorders	691	9	700	0.7
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	182	12	194	0.2
F60	Specific personality disorders	4 476	457	4 933	4.9
F61–F69	Disorders of adult personality and behaviour	176	27	203	0.2
F70–F79	Mental retardation	168	47	215	0.2
F80–F89	Disorders of psychological development	242	24	266	0.3
F90	Hyperkinetic disorders	87	16	103	0.1
F91	Conduct disorders	369	36	405	0.4
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	386	30	416	0.4
F99	Mental disorder not otherwise specified	245	–	245	0.2
G30	Alzheimer's disease	509	58	567	0.6
	Other factors related to mental and behavioural disorders and substance use (b)	198	438	636	0.6
	Other specified mental health-related principal diagnosis (c)	266	4	270	0.3
	Other (d)	8 211	1 260	9 471	9.5
	Total	90 276	9 517	99 793	100.0
<i>2012-13</i>					
F00–F03	Dementia	421	103	524	0.5
F04–F09	Other organic mental disorders	634	112	746	0.7
F10	Mental and behavioural disorders due to use of alcohol	2 201	397	2 598	2.5
F11–F19	Mental and behavioural disorders due to other psychoactive substances use	5 046	638	5 684	5.5
F20	Schizophrenia	18 198	1 815	20 013	19.3
F21, F24, F28, F29	Schizotypal and other delusional disorders	2 275	199	2 474	2.4
F22	Persistent delusional disorders	818	79	897	0.9
F23	Acute and transient psychotic disorders	1 363	125	1 488	1.4
F25	Schizoaffective disorders	6 328	632	6 960	6.7
F30	Manic episode	629	41	670	0.6

TABLE 12A.22

Table 12A.22 **Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)**

<i>ICD-10</i>		<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
F31	Bipolar affective disorders	8 518	753	9 271	8.9
F32	Depressive episode	13 361	781	14 142	13.6
F33	Recurrent depressive disorders	2 608	140	2 748	2.7
F34	Persistent mood (affective) disorders	888	35	923	0.9
F38, F39	Other and unspecified mood (affective) disorders	214	11	225	0.2
F40	Phobic anxiety disorders	81	5	86	0.1
F41	Other anxiety disorders	1 930	108	2 038	2.0
F42	Obsessive-compulsive disorders	325	12	337	0.3
F43	Reaction to severe stress and adjustment disorders	10 417	837	11 254	10.9
F44	Dissociative (conversion) disorders	134	9	143	0.1
F45, F48	Somatoform and other neurotic disorders	88	7	95	0.1
F50	Eating disorders	766	12	778	0.8
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	188	7	195	0.2
F60	Specific personality disorders	4 764	456	5 220	5.0
F61–F69	Disorders of adult personality and behaviour	190	29	219	0.2
F70–F79	Mental retardation	143	42	185	0.2
F80–F89	Disorders of psychological development	239	29	268	0.3
F90	Hyperkinetic disorders	61	20	81	0.1
F91	Conduct disorders	258	25	283	0.3
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	339	19	358	0.3
F99	Mental disorder not otherwise specified	160	644	804	0.8
G30	Alzheimer's disease	556	73	629	0.6
	Other factors related to mental and behavioural disorders and substance use (b)	162	433	595	0.6
	Other specified mental health-related principal diagnosis (c)	806	7	813	0.8
	Other (d)	8 493	1 438	9 931	9.6
	Total	93 602	10 073	103 675	100.0

Table 12A.22 Admitted patient mental health-related separations with specialised psychiatric care, by principal diagnosis in ICD-10-AM and hospital type (a)

<i>ICD-10</i>	<i>Public acute hospitals</i>	<i>Public psychiatric hospitals</i>	<i>Total</i>	<i>Per cent</i>
---------------	---------------------------------------	---	--------------	-----------------

- (a) Admitted patient separations refers to those non-ambulatory separations when a patient undergoes a hospital's formal admission process, completes an episode of care and 'separates' from the hospital, excluding ambulatory-equivalent separations. Separations for which care type was reported as Newborn with no qualified days and records for Hospital boarders and Posthumous organ procurement have been excluded.
- (b) Includes ICD-10-AM codes Z00.4, Z03.2, Z04.6, Z09.3, Z13.3, Z54.3, Z61.9, Z63.1, Z63.8, Z63.9, Z65.8, Z65.9, Z71.4, Z71.5 and Z76.0.
- (c) Includes separations for which the principal diagnosis was any other mental health-related principal diagnosis.
- (d) Includes all other codes not included as a mental health principal diagnosis.

– Nil or rounded to zero. **np** Not published.

Source: AIHW (various issues) *Mental Health Services in Australia* (various years), (available at <http://mhsa.aihw.gov.au/home/>).

Table 12A.23 Ambulatory-equivalent public mental health-related separations with specialised psychiatric care, by principal diagnosis, 2012-13 (a)

<i>ICD-10-AM code</i>		<i>Number of separations</i>	<i>Proportion of total separations</i>
F00–F09	Organic, including symptomatic, mental disorders	26	0.3
F10–F19	Mental and behavioural disorders due to psychoactive substance use	336	3.3
F20–F29	Schizophrenia, schizotypal and delusional disorders	604	6.0
F30–F39	Mood (affective) disorders	2 384	23.7
F40–F48	Neurotic, stress-related and somatoform disorders	2 878	28.6
F50–F59	Behavioural syndromes associated with physiological disturbances and physical factors	1 425	14.2
F60–F69	Disorders of adult personality and behaviour	206	2.0
F70–F79	Mental retardation	60	0.6
F80–F89	Disorders of psychological development	225	2.2
F90–F98	Behavioural and emotional disorders with onset usually occurring in childhood and adolescence	1 082	10.8
F99	Mental disorder not otherwise specified	4	–
G30	Alzheimer's disease	24	0.2
	Other factors related to mental and behavioural disorders and substance use (b)	113	1.1
	Other specified mental health-related principal diagnosis (c)	81	0.8
	Other (d)	609	6.1
Total		10 057	100.0

(a) Includes separations for both public acute and public psychiatric hospitals.

(b) Includes ICD-10-AM codes Z00.4, Z03.2, Z04.6, Z09.3, Z13.3, Z54.3, Z61.9, Z63.1, Z63.8, Z63.9, Z65.8, Z65.9, Z71.4, Z71.5 and Z76.0.

(c) Includes separations for which the principal diagnosis was any other mental health-related principal diagnosis.

(d) Includes all other codes not included as a mental health principal diagnosis.

– Nil or rounded to zero.

Source: AIHW (2015) *Mental Health Services in Australia* (available at <http://mhsa.aihw.gov.au/home/>).

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>2005-06</i>									
<i>Number</i>									
<i>Males</i>									
Less than 15 years	39 242	61 978	65 976	27 955	23 046	2 652	8 294	1 534	230 677
15–24	135 686	152 875	83 386	30 460	25 441	2 531	28 628	3 871	462 878
25–34	252 587	252 055	108 586	47 572	34 707	4 812	32 443	6 435	739 197
35–44	199 198	194 510	91 381	45 952	32 112	4 062	16 903	4 290	588 408
45–54	113 329	119 193	57 663	32 580	22 076	4 822	12 055	2 162	363 880
55–64	51 652	65 399	33 349	21 487	9 102	1 782	4 657	1 212	188 640
65 years and over	29 325	106 367	26 531	22 786	10 204	5 496	5 092	794	206 595
Total males (a)	821 019	952 377	466 872	228 792	156 688	26 157	108 072	20 298	2 780 275
<i>Females</i>									
Less than 15 years	30 780	38 115	45 103	18 043	13 925	2 195	9 272	649	158 082
15–24	112 548	150 119	79 990	38 489	19 770	4 416	30 477	3 038	438 847
25–34	129 122	153 943	80 377	44 052	21 971	4 023	19 210	4 221	456 919
35–44	121 075	160 153	77 948	44 759	25 206	3 916	14 329	3 616	451 002
45–54	92 416	129 707	64 160	45 469	19 741	4 136	11 232	2 817	369 678
55–64	57 219	74 678	36 751	24 617	12 383	3 048	6 025	1 228	215 949
65 years and over	46 767	174 060	41 180	48 247	22 129	9 058	11 660	488	353 589
Total females (a)	589 927	880 775	425 509	263 676	135 125	30 792	102 205	16 057	2 444 066
<i>People</i>									
Less than 15 years	70 129	100 093	111 085	45 998	37 020	4 864	17 599	2 184	388 972
15–24	248 456	303 005	163 378	68 949	45 224	6 949	59 160	6 909	902 030
25–34	382 257	405 998	188 965	91 624	56 678	8 847	51 733	10 656	1 196 758
35–44	320 939	354 663	169 330	90 711	57 321	7 989	31 307	7 906	1 040 166
45–54	206 402	248 900	121 823	78 049	41 817	8 993	23 311	4 979	734 274

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
55–64	109 218	140 079	70 101	46 103	21 487	4 831	10 691	2 440	404 950
65 years and over	75 010	271 201	67 638	70 777	32 275	14 672	14 202	1 268	547 043
Total (b)	1 832 177	1 833 205	892 393	492 468	302 400	65 576	210 833	36 356	5 665 408
<i>Rate (per 1,000 population) (c)</i>									
Males									
Less than 15 years	57.4	124.0	155.3	132.9	157.0	53.3	260.4	57.8	111.2
15–24	288.7	425.5	287.0	202.1	235.9	77.1	1 037.5	231.8	318.0
25–34	523.7	687.2	382.7	329.6	341.7	168.8	1 215.8	355.3	509.0
35–44	401.5	513.6	308.8	295.2	283.4	119.8	684.2	244.7	388.1
45–54	245.9	346.9	208.6	223.3	201.6	137.5	531.0	146.5	258.3
55–64	141.1	244.0	148.1	192.2	103.2	61.1	275.0	122.4	169.1
65 years and over	72.1	353.6	118.0	208.7	99.1	173.5	362.9	157.9	172.8
Total males (a)	246.3	378.9	232.4	223.8	208.2	111.2	624.3	185.0	274.1
Females									
Less than 15 years	47.4	80.5	111.9	91.6	99.2	46.7	301.5	26.1	80.4
15–24	249.6	435.0	286.1	273.5	192.8	138.9	1 164.8	196.3	315.1
25–34	265.5	417.6	283.4	314.7	222.2	135.6	713.6	238.6	314.7
35–44	241.3	414.3	258.0	292.1	222.7	111.5	565.2	222.3	294.1
45–54	197.9	370.4	230.1	315.2	176.1	115.5	463.4	207.5	259.2
55–64	157.2	274.2	167.9	231.8	136.3	104.2	348.1	156.3	195.1
65 years and over	92.6	463.9	157.3	373.8	168.4	232.0	675.5	110.4	241.7
Total females (a)	173.7	336.1	210.6	259.5	171.8	120.5	594.4	158.6	235.4
Total people									
Less than 15 years	52.6	102.8	134.1	113.0	128.9	50.3	281.1	42.4	96.3
15–24	269.8	430.2	286.5	236.6	215.0	107.5	1 100.5	214.7	316.7
25–34	394.6	552.1	333.1	322.2	282.7	152.1	965.1	297.7	412.1

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
35–44	321.7	463.4	283.1	293.7	253.1	115.8	625.4	233.9	341.1
45–54	222.5	358.8	219.4	269.0	188.7	126.9	496.6	175.7	259.0
55–64	149.6	259.2	157.9	211.5	120.0	82.7	312.2	137.4	182.2
65 years and over	82.3	401.2	139.0	297.1	137.7	207.5	453.8	134.2	205.8
Total (b)	265.1	357.3	221.5	242.2	195.6	130.5	616.3	170.8	274.9
<i>2006-07</i>									
<i>Number</i>									
Males									
Less than 15 years	52 850	65 142	68 238	29 023	26 869	6 118	8 058	1 715	258 013
15–24	157 769	146 075	99 033	35 453	26 836	4 085	26 355	3 735	499 341
25–34	293 437	255 661	136 745	52 831	48 005	6 654	31 352	6 857	831 542
35–44	242 766	200 969	110 867	50 402	44 058	6 020	18 745	5 352	679 179
45–54	147 155	125 412	68 829	35 713	29 942	6 840	11 414	1 893	427 198
55–64	70 202	69 302	37 575	23 399	12 528	2 566	4 170	1 148	220 890
65 years and over	38 374	88 736	31 958	22 163	9 776	6 580	3 974	679	202 240
Total males (a)	1 003 086	955 935	553 343	249 098	198 083	38 926	104 893	21 384	3 124 748
Females									
Less than 15 years	34 800	42 273	49 801	17 356	17 002	4 062	7 953	992	174 239
15–24	127 370	150 159	94 250	44 259	24 824	7 897	28 382	2 767	479 908
25–34	145 183	156 335	92 550	46 035	27 152	5 230	19 528	4 533	496 546
35–44	153 131	161 996	96 595	50 486	34 278	6 774	16 953	3 732	523 945
45–54	121 441	131 390	74 283	48 786	27 710	6 065	11 262	2 825	423 762
55–64	71 887	77 097	43 412	28 175	16 460	3 867	5 556	1 185	247 639
65 years and over	68 461	152 440	46 652	46 821	21 430	15 554	11 044	362	362 764
Total females (a)	722 696	874 196	497 571	282 028	168 894	49 474	102 205	16 401	2 713 465
People									

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Less than 15 years	87 685	107 415	118 065	46 379	43 871	10 183	16 055	2 707	432 360
15–24	285 537	296 287	193 287	79 712	51 660	12 014	54 772	6 502	979 771
25–34	439 120	412 062	229 296	98 868	75 157	11 886	50 910	11 390	1 328 689
35–44	396 346	362 993	207 463	100 888	78 348	12 830	35 718	9 084	1 203 670
45–54	269 194	256 802	143 112	84 499	57 653	12 973	22 694	4 718	851 645
55–64	142 214	146 399	80 987	51 574	28 989	6 485	9 726	2 333	468 707
65 years and over	106 985	241 176	78 610	68 984	31 206	22 166	15 018	1 041	565 186
Total (b)	1 828 468	1 830 278	1 050 960	535 809	382 304	93 186	207 487	37 785	5 966 277
<i>Rate (per 1,000 population) (c)</i>									
Males									
Less than 15 years	77.3	129.7	158.8	136.2	183.0	123.3	253.0	64.7	123.8
15–24	331.2	397.0	332.9	229.5	244.3	123.0	955.1	215.2	336.3
25–34	610.1	695.6	477.2	362.0	475.1	237.6	1 174.9	378.4	571.4
35–44	490.4	526.6	369.1	317.5	388.6	179.5	758.8	303.3	445.2
45–54	315.9	359.3	244.4	240.6	270.4	193.2	502.8	126.1	298.9
55–64	187.2	251.3	161.3	201.5	137.9	85.0	246.3	111.5	192.3
65 years and over	92.1	286.4	137.0	195.1	92.9	202.6	283.2	126.1	164.2
Total males (a)	299.2	375.2	271.2	238.1	262.6	165.0	605.3	188.8	304.5
Females									
Less than 15 years	53.5	88.7	122.2	86.9	121.2	87.0	258.6	39.6	88.1
15–24	277.5	424.8	328.1	306.8	237.4	246.6	1 084.7	174.6	337.3
25–34	300.0	426.1	325.5	328.1	276.3	179.2	725.5	254.5	343.0
35–44	304.8	414.6	315.0	325.5	302.9	194.3	668.7	226.5	339.1
45–54	255.9	368.4	259.7	331.3	244.5	166.7	464.6	202.4	291.7
55–64	191.4	272.9	190.4	253.8	174.7	127.6	321.0	142.6	215.7
65 years and over	133.2	398.1	172.9	351.7	160.4	391.0	639.8	76.5	242.5

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Total females (a)	209.9	330.0	241.6	271.7	214.8	191.9	594.5	155.2	258.1
Total people									
Less than 15 years	65.7	109.8	141.1	112.3	152.8	105.7	256.4	52.5	106.4
15–24	305.3	410.7	330.6	266.8	240.9	184.2	1 018.9	195.8	337.0
25–34	455.1	561.0	401.7	345.4	377.1	207.8	949.8	317.0	457.7
35–44	397.4	470.0	341.8	321.4	345.9	187.5	713.6	266.3	392.0
45–54	286.3	363.9	252.1	285.8	257.3	180.7	483.5	162.9	295.5
55–64	189.5	262.2	175.7	227.0	156.7	107.2	284.0	125.4	204.1
65 years and over	114.9	348.1	156.3	279.6	130.7	306.8	479.9	102.9	207.2
Total (b)	269.7	353.3	256.7	257.9	249.3	189.2	602.9	172.3	288.0
<i>2007-08</i>									
<i>Number</i>									
Males									
Less than 15 years	54 762	54 125	76 331	29 163	29 505	9 447	8 265	1 640	263 238
15–24	184 734	137 121	108 312	36 359	29 943	7 412	24 591	3 215	531 687
25–34	355 111	236 320	153 452	56 300	56 261	11 232	27 680	7 053	903 409
35–44	292 683	197 867	127 742	51 256	51 794	10 167	17 279	4 889	753 677
45–54	183 155	126 146	81 201	37 727	37 971	10 928	10 690	2 409	490 227
55–64	83 938	67 908	42 359	25 594	15 663	4 931	4 259	909	245 561
65 years and over	45 786	82 281	35 607	24 218	11 745	8 410	5 444	528	214 019
Total males (a)	1 200 743	906 012	625 063	260 826	232 893	62 527	98 692	20 646	3 407 402
Females									
Less than 15 years	36 288	36 896	52 758	16 990	16 432	7 796	10 379	778	178 317
15–24	132 106	144 876	100 645	46 955	27 868	11 066	29 435	3 007	495 958
25–34	163 717	141 706	101 403	46 049	33 118	8 750	17 649	4 138	516 530
35–44	174 214	158 411	106 223	56 335	44 022	11 435	16 781	3 644	571 065

TABLE 12A.24

Table 12A.24 Community mental health service contacts, by sex and age group

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
45–54	132 986	128 081	80 389	48 451	34 139	11 326	12 871	2 556	450 799
55–64	70 774	78 566	44 263	30 097	20 837	6 149	6 496	1 005	258 187
65 years and over	74 591	139 767	51 659	43 517	23 763	19 503	13 565	368	366 733
Total females (a)	785 095	830 400	537 415	288 596	200 195	76 035	108 200	15 500	2 841 436
People									
Less than 15 years	91 158	91 021	129 090	46 156	45 937	17 244	18 646	2 418	441 670
15–24	317 087	281 997	208 957	83 315	57 812	18 478	54 093	6 222	1 027 961
25–34	519 221	378 026	254 855	102 350	89 379	19 982	45 451	11 191	1 420 455
35–44	467 790	356 307	233 965	107 592	95 845	21 603	34 102	8 533	1 325 737
45–54	316 282	254 232	161 590	86 178	72 135	22 255	23 573	4 965	941 210
55–64	154 799	146 484	86 622	55 693	36 500	11 081	10 772	1 914	503 865
65 years and over	120 459	222 048	87 266	67 735	35 508	27 914	19 031	896	580 857
Total (b)	2 072 440	1 736 456	1 162 557	554 558	456 942	147 701	207 467	36 146	6 374 267
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	80.1	106.5	173.4	134.1	199.9	189.5	255.5	61.2	125.0
15–24	378.3	361.5	353.3	228.7	269.3	222.5	879.9	181.2	349.0
25–34	730.6	631.2	520.6	370.7	553.2	403.9	1 002.3	377.2	609.2
35–44	590.4	513.6	415.7	315.2	457.8	307.7	683.8	276.6	489.3
45–54	387.5	355.1	281.8	248.4	339.2	305.8	465.1	157.3	337.1
55–64	217.7	238.9	176.0	212.4	167.7	158.2	236.9	84.8	207.3
65 years and over	107.0	258.1	147.5	205.9	109.2	251.8	357.1	91.3	168.8
Total males (a)	353.1	346.3	297.8	241.8	305.1	265.7	558.8	175.8	324.9
Females									
Less than 15 years	55.8	76.4	126.5	83.3	116.3	166.0	331.4	30.8	89.2
15–24	283.6	403.3	341.4	317.6	263.3	347.2	1 122.1	184.9	342.5

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
25–34	335.3	380.6	348.1	318.9	333.8	302.8	641.3	225.7	351.3
35–44	345.2	401.1	339.7	356.4	389.0	330.3	650.8	218.3	365.9
45–54	275.6	352.2	273.6	322.3	297.9	307.3	525.9	178.9	304.4
55–64	182.3	268.5	187.1	259.9	214.4	195.8	346.6	114.4	217.1
65 years and over	142.3	357.3	186.1	318.1	175.2	482.3	729.0	72.7	239.8
Total females (a)	225.8	307.3	255.0	270.5	252.1	293.6	619.6	141.3	265.6
Total people									
Less than 15 years	68.3	91.8	150.6	109.5	159.0	178.1	292.9	46.4	107.6
15–24	332.3	381.8	347.5	271.6	266.3	283.4	998.4	182.9	345.9
25–34	532.9	506.2	434.9	345.4	444.8	352.4	824.3	302.2	480.9
35–44	467.6	456.7	377.4	335.5	423.5	319.2	667.9	248.3	427.5
45–54	331.1	353.6	277.7	285.2	318.4	306.6	496.7	167.7	320.7
55–64	200.1	253.9	181.5	235.7	191.5	177.1	293.3	98.1	212.3
65 years and over	126.5	312.7	168.2	266.2	146.0	378.0	562.2	82.6	207.6
Total (b)	289.8	327.1	276.7	256.6	279.4	280.9	591.8	158.6	295.7
2008-09									
Number									
Males									
Less than 15 years	53 539	57 020	69 564	34 115	33 837	9 406	8 128	1 975	267 584
15–24	171 329	133 507	84 433	38 255	35 906	10 491	25 270	4 347	503 538
25–34	313 446	216 375	125 107	60 557	59 071	13 937	27 686	7 559	823 738
35–44	282 427	193 192	105 837	58 506	59 530	14 136	20 277	4 831	738 736
45–54	186 573	125 183	68 080	41 871	42 059	12 907	10 206	2 541	489 420
55–64	84 909	72 207	35 777	25 053	18 046	8 227	5 549	1 236	251 004
65 years and over	58 257	79 146	29 029	26 172	13 837	8 911	6 658	455	222 465
Total males (a)	1 156 291	876 648	517 871	285 039	262 412	78 015	103 779	22 955	3 303 010

TABLE 12A.24

Table 12A.24 Community mental health service contacts, by sex and age group

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Females									
Less than 15 years	37 897	37 270	48 266	21 148	19 012	7 034	10 011	1 028	181 666
15–24	136 950	142 510	83 923	53 375	33 094	13 786	31 795	3 405	498 838
25–34	153 624	142 334	81 922	51 410	41 034	10 557	22 738	4 362	507 981
35–44	172 520	155 971	83 097	59 040	51 147	14 705	18 509	3 805	558 794
45–54	137 339	125 192	64 417	52 783	39 505	12 354	13 842	2 274	447 706
55–64	74 183	78 068	35 652	32 153	24 338	7 722	7 771	1 265	261 152
65 years and over	89 167	131 117	43 626	49 024	26 211	20 078	13 861	221	373 305
Total females (a)	805 354	812 501	441 009	319 368	234 382	86 247	118 527	16 371	2 833 759
People									
Less than 15 years	91 569	94 290	117 847	55 269	52 849	16 440	18 141	3 005	449 410
15–24	308 462	276 021	168 356	91 631	69 000	24 277	57 215	7 752	1 002 714
25–34	467 566	358 709	207 029	111 989	100 105	24 494	50 567	11 921	1 332 380
35–44	455 922	349 275	188 938	117 553	110 687	28 912	38 911	8 636	1 298 834
45–54	324 932	250 377	132 497	94 658	81 568	25 305	24 133	4 815	938 285
55–64	159 347	150 275	71 430	57 209	42 386	15 949	13 338	2 501	512 435
65 years and over	147 707	210 324	72 655	75 203	40 059	28 989	20 657	676	596 270
Total (b)	2 051 579	1 689 328	958 921	609 276	525 217	173 788	223 328	39 328	6 270 765
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	77.9	110.8	154.6	152.6	227.7	187.8	247.3	72.9	125.3
15–24	345.0	344.6	268.3	234.2	320.5	312.7	901.8	240.7	324.1
25–34	633.6	563.0	411.2	376.7	568.7	497.8	969.5	395.6	540.6
35–44	569.4	496.7	337.7	350.9	528.3	432.5	792.8	271.2	475.3
45–54	389.5	346.4	231.3	268.6	372.2	358.5	439.3	163.3	331.0
55–64	215.0	246.9	144.6	200.9	188.7	257.7	299.3	111.0	206.3

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
65 years and over	132.3	241.1	116.1	214.4	125.6	258.4	419.9	73.8	170.2
Total males (a)	336.2	330.2	241.4	256.3	340.6	329.3	583.8	189.6	309.9
Females									
Less than 15 years	58.0	76.2	113.1	100.7	133.7	148.6	316.0	40.5	89.7
15–24	289.0	387.7	278.3	352.0	309.4	431.3	1 213.5	207.6	337.9
25–34	309.5	373.9	273.8	343.0	406.0	365.8	812.2	231.5	338.0
35–44	341.5	391.6	261.9	367.8	454.4	426.9	709.3	225.5	355.6
45–54	280.7	338.3	214.3	344.0	342.3	332.4	562.2	156.6	297.4
55–64	185.8	258.6	145.9	266.9	243.8	238.0	402.2	136.1	212.9
65 years and over	166.6	327.9	152.7	347.9	190.0	485.9	718.9	41.0	238.4
Total females (a)	227.9	296.4	205.1	292.4	293.9	333.1	673.0	144.4	260.7
Total people									
Less than 15 years	68.3	93.9	134.4	127.4	181.7	168.7	281.1	57.2	108.0
15–24	317.9	365.6	273.2	290.9	315.1	370.6	1 055.2	224.9	330.9
25–34	471.8	468.9	343.1	360.5	488.4	430.8	894.2	314.1	440.2
35–44	455.3	443.6	299.6	359.2	491.4	430.7	753.1	248.9	415.5
45–54	335.6	342.3	222.7	306.1	357.1	345.8	504.3	160.1	314.5
55–64	200.7	252.8	145.3	233.3	216.9	247.8	352.3	122.4	209.7
65 years and over	151.4	288.9	135.6	286.0	161.4	382.4	587.9	58.5	207.5
Total (b)	294.8	313.6	223.5	277.1	335.5	351.5	632.5	167.4	291.9
2009-10									
Number									
Males									
Less than 15 years	55 617	58 865	64 050	36 263	36 458	11 539	10 502	2 894	276 188
15–24	194 198	136 613	75 322	43 713	37 083	14 725	23 932	4 080	529 666
25–34	360 216	212 696	112 225	68 442	59 549	20 832	30 156	6 456	870 572

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
35–44	321 520	198 512	96 721	64 902	61 150	17 384	21 995	5 110	787 294
45–54	203 206	128 415	62 759	47 702	46 587	15 444	12 722	2 666	519 501
55–64	95 362	72 427	34 715	28 721	21 436	8 458	7 146	1 171	269 436
65 years and over	66 302	81 070	27 779	31 519	13 287	10 247	11 291	401	241 896
Total males (a)	1 300 584	888 610	473 593	321 343	275 600	98 681	117 749	22 779	3 498 939
Females									
Less than 15 years	42 034	38 740	43 742	25 000	18 535	9 827	11 446	1 187	190 511
15–24	138 723	153 599	78 342	58 934	32 183	16 731	36 918	3 382	518 812
25–34	156 345	146 349	73 952	55 207	44 709	14 281	25 806	4 465	521 114
35–44	183 051	160 410	76 764	63 702	52 955	19 542	20 050	3 716	580 190
45–54	144 038	134 412	59 620	59 271	41 185	16 258	14 457	1 984	471 225
55–64	88 349	80 891	34 718	35 638	25 366	10 308	10 017	1 131	286 418
65 years and over	95 084	132 732	42 704	53 999	25 146	19 118	20 162	336	389 281
Total females (a)	849 771	847 150	409 855	351 908	240 123	106 109	138 868	16 205	2 959 989
People									
Less than 15 years	97 709	97 605	107 792	61 263	54 993	21 423	21 948	4 081	466 814
15–24	333 043	290 216	153 672	102 649	69 267	31 571	60 938	7 462	1 048 818
25–34	516 863	359 201	186 179	123 674	104 258	35 213	56 025	10 921	1 392 334
35–44	505 271	358 974	173 485	128 624	114 176	37 026	42 091	8 826	1 368 473
45–54	347 565	262 865	122 379	106 975	87 781	31 772	27 213	4 650	991 200
55–64	184 322	153 318	69 433	64 362	46 803	18 801	17 163	2 302	556 504
65 years and over	161 548	213 802	70 483	85 522	38 453	29 400	31 453	737	631 398
Total (b)	2 242 034	1 736 010	883 458	680 134	543 348	212 599	257 497	38 984	6 594 064
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	80.4	113.1	139.4	159.3	244.5	230.1	314.0	106.0	127.9

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
15–24	377.6	336.5	229.9	256.0	323.8	430.7	842.1	216.5	328.0
25–34	699.3	526.1	354.3	403.0	556.5	739.1	1 016.4	323.2	547.2
35–44	643.8	504.7	304.2	385.2	545.4	535.3	849.8	281.4	502.1
45–54	419.7	350.5	209.2	300.5	408.5	428.2	541.9	168.5	346.6
55–64	236.6	241.6	137.1	223.9	219.2	260.1	375.8	101.6	216.4
65 years and over	146.0	239.5	106.6	248.6	117.6	287.2	683.5	61.0	178.9
Total males (a)	369.9	325.7	215.1	281.2	352.6	419.0	661.2	182.1	320.5
Females									
Less than 15 years	63.9	78.5	100.5	116.7	129.7	206.9	357.5	46.3	93.0
15–24	285.5	404.7	250.8	375.6	295.9	518.1	1 394.4	201.1	341.5
25–34	303.9	367.5	236.6	349.7	429.4	488.6	886.7	226.9	332.9
35–44	358.6	397.8	238.3	389.8	471.3	569.1	759.8	214.6	364.9
45–54	290.4	357.3	194.1	378.0	354.1	434.8	584.1	134.3	308.1
55–64	215.8	260.4	138.1	285.7	248.8	311.1	504.8	115.3	227.1
65 years and over	173.5	323.7	144.2	372.0	178.6	452.6	1 008.8	58.8	242.1
Total females (a)	235.5	302.5	185.6	312.9	298.4	416.6	780.0	140.4	266.4
Total people									
Less than 15 years	72.4	96.3	120.5	138.6	188.3	219.4	335.3	77.1	110.9
15–24	333.0	369.5	240.1	313.3	310.2	474.9	1 110.0	209.2	334.7
25–34	502.0	447.6	295.8	377.4	493.8	613.4	953.3	275.4	441.1
35–44	500.3	450.6	271.1	387.6	508.6	554.2	805.2	248.8	433.4
45–54	354.6	354.0	201.6	339.0	381.1	432.5	564.3	152.0	327.3
55–64	226.9	251.2	137.6	254.4	234.3	286.4	441.7	107.9	222.0
65 years and over	161.2	285.6	126.6	314.5	151.5	377.3	861.6	60.0	213.3
Total (b)	312.1	315.7	197.5	299.7	332.4	420.7	725.4	171.0	297.7

2010-11

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Number</i>									
Males									
Less than 15 years	53 036	57 824	72 813	42 776	32 021	7 000	8 900	2 339	276 709
15–24	206 312	147 891	92 363	54 328	38 495	9 058	27 770	4 397	580 614
25–34	374 096	211 602	128 045	74 158	63 636	14 058	26 754	7 325	899 674
35–44	351 095	204 707	111 891	70 991	65 026	13 116	20 986	5 535	843 347
45–54	214 607	133 645	73 858	51 986	47 926	11 299	14 007	2 788	550 116
55–64	103 602	73 148	38 367	31 831	23 731	5 281	5 971	1 255	283 186
65 years and over	67 449	86 616	31 531	31 509	14 623	8 192	8 124	418	248 462
Total males (a)	1 378 280	915 441	548 876	357 783	285 478	68 048	112 834	24 061	3 690 801
Females									
Less than 15 years	42 780	35 815	51 300	28 703	20 137	6 107	10 502	827	196 171
15–24	161 084	169 999	96 151	70 234	34 661	11 384	39 911	3 040	586 464
25–34	173 977	149 064	82 701	60 239	45 294	9 033	21 600	5 224	547 132
35–44	202 688	171 229	88 749	67 465	54 097	12 340	21 573	4 208	622 349
45–54	158 044	136 234	68 181	65 891	41 782	12 316	13 965	2 163	498 576
55–64	93 863	84 995	41 885	38 616	26 114	8 260	10 058	1 291	305 082
65 years and over	101 540	138 036	45 582	54 456	28 308	15 615	11 783	404	395 724
Total females (a)	938 018	885 380	474 560	385 808	250 423	75 100	129 900	17 160	3 156 349
People									
Less than 15 years	95 881	93 665	124 113	71 479	52 158	13 132	19 402	3 166	472 996
15–24	367 518	317 934	188 540	124 570	73 160	20 501	67 706	7 437	1 167 366
25–34	548 366	360 809	210 754	134 403	108 930	23 157	48 373	12 549	1 447 341
35–44	554 048	376 073	200 640	138 490	119 130	25 645	42 569	9 743	1 466 338
45–54	373 577	269 912	142 067	117 887	89 718	23 660	27 974	4 951	1 049 746
55–64	197 940	158 143	80 256	70 453	49 849	13 570	16 029	2 546	588 786

TABLE 12A.24

Table 12A.24 Community mental health service contacts, by sex and age group

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
65 years and over	169 546	224 652	77 113	85 985	42 937	23 838	19 907	822	644 800
Total (b)	2 408 488	1 994 752	1 023 502	752 186	560 498	150 689	242 857	41 221	7 174 193
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	76.2	112.1	160.5	187.9	216.6	141.5	263.3	86.7	128.6
15–24	418.9	375.4	290.8	319.6	339.4	264.7	926.7	225.6	369.5
25–34	734.7	521.2	409.1	420.3	590.1	481.0	875.6	346.0	564.4
35–44	707.1	525.3	355.8	414.8	586.6	402.0	789.3	298.3	540.4
45–54	440.9	365.3	246.8	320.8	421.1	309.1	591.5	172.8	365.7
55–64	252.2	241.3	151.4	243.2	240.1	156.3	311.2	104.4	224.3
65 years and over	143.9	250.4	118.7	243.7	127.0	218.5	474.7	62.2	179.3
Total males (a)	395.2	337.3	252.2	307.8	366.3	283.1	611.8	186.5	339.1
Females									
Less than 15 years	65.0	73.2	119.3	131.6	142.3	132.0	327.5	32.8	96.1
15–24	342.8	454.1	313.4	440.8	322.3	354.9	1 403.9	184.0	392.3
25–34	342.9	371.5	266.4	363.7	430.9	305.1	717.6	266.3	348.7
35–44	398.7	425.5	276.8	403.0	488.5	362.0	801.5	242.0	391.9
45–54	319.0	361.3	223.0	410.8	361.1	330.3	559.4	145.9	325.5
55–64	225.1	269.1	165.1	295.7	254.3	244.6	501.0	128.0	237.6
65 years and over	181.4	329.9	150.9	361.1	199.0	360.7	567.1	70.0	240.8
Total females (a)	262.2	316.1	216.1	334.4	311.9	290.3	708.3	148.2	284.6
Total people									
Less than 15 years	70.8	93.2	140.4	160.3	180.3	137.2	294.6	60.6	112.8
15–24	381.9	413.8	301.9	378.3	331.1	309.2	1 159.5	206.5	380.7
25–34	539.5	447.0	338.1	392.9	511.6	393.6	797.5	307.7	457.6
35–44	551.3	474.7	316.0	409.1	537.6	384.4	795.6	271.1	465.7

TABLE 12A.24

Table 12A.24 Community mental health service contacts, by sex and age group

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
45–54	380.4	363.3	234.8	365.6	390.9	320.4	575.0	159.9	345.8
55–64	239.1	255.5	158.3	269.4	247.4	200.9	408.3	115.2	231.2
65 years and over	164.8	294.0	135.8	307.0	166.8	295.1	525.3	65.8	212.8
Total (b)	341.4	362.1	234.3	325.4	354.8	301.9	659.9	168.1	326.8
2011-12 (d), (e)									
Number									
Males									
Less than 15 years	55 345	na	81 536	42 604	27 559	8 921	7 537	3 442	226 944
15–24	216 554	na	112 892	53 638	41 286	7 245	32 648	5 441	469 704
25–34	409 178	na	153 979	72 842	65 433	9 027	30 842	7 567	748 868
35–44	381 886	na	135 033	67 868	68 603	7 331	26 509	6 124	693 354
45–54	235 841	na	87 201	51 029	49 904	5 818	14 634	2 998	447 425
55–64	109 607	na	46 872	31 779	25 077	2 377	7 259	1 481	224 452
65 years and over	74 692	na	37 292	32 460	19 286	4 627	7 209	431	175 997
Total males (a)	1 499 608	na	654 897	352 432	297 165	45 374	127 087	27 491	3 004 054
Females									
Less than 15 years	46 038	na	56 586	32 588	20 968	6 887	8 283	1 769	173 119
15–24	180 415	na	117 464	77 430	38 808	10 642	39 588	4 503	468 850
25–34	182 640	na	100 772	62 852	45 980	5 749	24 701	5 940	428 634
35–44	194 485	na	105 091	65 798	56 571	7 198	21 515	5 121	455 779
45–54	160 016	na	81 160	62 452	47 170	6 325	16 430	2 350	375 903
55–64	95 997	na	47 803	38 979	28 823	5 553	8 560	1 458	227 173
65 years and over	115 156	na	50 243	51 280	33 037	6 905	12 501	301	269 423
Total females (a)	980 470	na	559 217	391 569	271 368	49 284	132 237	21 445	2 405 590
People									
Less than 15 years	101 432	na	138 122	75 193	48 527	15 830	15 820	5 211	400 135

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
15–24	397 089	na	230 408	131 068	80 107	17 903	72 236	9 944	938 755
25–34	592 121	na	254 751	135 697	111 452	14 779	55 555	13 507	1 177 862
35–44	576 860	na	240 124	133 668	125 201	14 543	48 024	11 245	1 149 665
45–54	396 524	na	168 363	113 487	97 099	12 148	31 070	5 348	824 039
55–64	206 078	na	94 714	70 762	53 900	7 951	15 819	2 939	452 163
65 years and over	190 403	na	87 535	83 740	52 330	11 545	19 710	732	445 995
Total (b)	2 573 242	na	1 214 208	752 419	593 178	98 778	259 346	48 936	5 540 107
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	78.5	na	176.5	181.8	184.9	180.3	216.8	126.4	136.6
15–24	440.0	na	353.5	311.3	365.8	212.6	1 097.0	284.6	398.2
25–34	794.3	na	483.1	388.3	596.9	306.7	975.6	349.2	617.0
35–44	763.9	na	424.9	387.0	620.6	225.7	977.9	330.6	586.7
45–54	485.4	na	289.7	310.5	439.4	161.3	613.7	185.1	392.2
55–64	263.5	na	183.0	237.3	252.0	69.6	378.5	121.3	231.1
65 years and over	152.2	na	133.3	238.1	160.5	117.4	394.8	59.4	161.2
Total males (a)	426.9	na	296.3	294.3	377.6	189.4	670.5	211.6	363.1
Females									
Less than 15 years	69.1	na	129.3	144.8	147.0	149.1	251.6	69.5	109.8
15–24	384.5	na	380.9	478.1	362.1	335.0	1 402.8	276.0	417.5
25–34	355.4	na	318.6	359.4	431.2	193.3	794.0	298.6	359.4
35–44	380.6	na	325.2	385.3	513.4	214.4	788.6	294.3	381.9
45–54	323.0	na	263.5	384.6	409.1	171.2	658.9	157.5	324.7
55–64	227.9	na	186.8	292.2	279.9	163.2	423.7	140.8	232.3
65 years and over	199.5	na	159.8	326.4	225.8	154.4	574.0	48.8	212.5
Total females (a)	272.0	na	251.7	332.0	334.9	196.6	709.6	183.2	287.6

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Total people									
Less than 15 years	74.0	na	153.5	163.7	166.3	165.5	233.7	98.9	123.6
15–24	413.0	na	367.1	392.1	364.1	271.9	1 245.9	280.6	407.7
25–34	575.4	na	401.2	374.4	515.3	249.8	885.7	325.0	489.5
35–44	570.6	na	374.6	386.1	567.2	220.2	882.9	313.0	484.0
45–54	404.1	na	276.5	347.4	424.3	166.4	636.9	171.9	358.5
55–64	246.2	na	185.0	264.7	266.2	116.7	401.7	130.2	232.0
65 years and over	178.3	na	147.3	285.4	196.4	137.2	492.3	54.5	189.0
Total (b)	362.0	na	274.1	316.8	371.8	200.9	690.4	198.0	333.2
2012-13 (d), (e)									
Number									
Males									
Less than 15 years	54 886	na	101 393	40 319	32 198	6 017	4 694	4 444	243 951
15–24	252 340	na	140 155	57 807	42 579	4 780	32 991	6 750	537 402
25–34	427 303	na	179 352	78 313	67 864	5 547	31 878	10 433	800 690
35–44	390 035	na	167 225	70 617	71 895	4 614	24 683	8 337	737 406
45–54	256 661	na	110 105	51 691	54 143	3 711	17 538	3 360	497 209
55–64	120 294	na	54 708	31 928	27 278	1 548	6 794	1 616	244 166
65 years and over	94 464	na	47 018	33 355	21 324	3 659	7 452	709	207 981
Total males (a)	1 623 182	na	800 010	364 182	317 288	29 911	126 526	35 659	3 296 758
Females									
Less than 15 years	66 002	na	82 575	37 485	23 693	6 265	9 562	2 993	228 575
15–24	239 714	na	159 312	93 177	44 801	8 942	41 531	5 850	593 327
25–34	215 308	na	119 338	69 032	49 182	4 210	26 061	7 491	490 622
35–44	224 573	na	125 231	70 924	57 947	4 520	23 613	6 412	513 220
45–54	189 919	na	95 677	62 987	52 638	3 733	18 052	3 317	426 323

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
55–64	111 697	na	59 243	39 149	29 635	3 628	10 281	1 976	255 609
65 years and over	142 960	na	64 849	57 774	39 404	5 985	11 662	405	323 039
Total females (a)	1 202 344	na	706 340	430 817	297 348	37 348	141 357	28 452	2 844 006
People									
Less than 15 years	121 001	na	183 971	77 804	55 891	12 287	14 256	7 437	472 647
15–24	492 382	na	299 506	150 990	87 435	13 734	74 522	12 600	1 131 169
25–34	642 807	na	298 692	147 349	117 125	9 802	57 942	17 924	1 291 641
35–44	614 826	na	292 473	141 545	129 922	9 148	48 296	14 749	1 250 959
45–54	447 357	na	205 951	114 680	106 817	7 458	35 590	6 677	924 530
55–64	232 217	na	113 969	71 082	56 932	5 182	17 075	3 592	500 049
65 years and over	239 217	na	111 867	91 129	60 770	9 652	19 114	1 114	532 863
Total (b)	2 924 684	na	1 506 598	795 202	639 715	67 680	267 887	64 111	6 265 877
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	77.3	..	215.7	166.6	213.9	..	130.9	161.5	144.7
15–24	510.1	..	432.7	328.2	378.9	..	1 116.4	350.7	451.6
25–34	816.8	..	546.6	388.0	607.3	..	971.1	466.8	640.5
35–44	775.5	..	520.7	391.3	653.2	..	889.1	444.7	617.9
45–54	528.4	..	361.5	308.2	476.4	..	723.8	205.7	433.1
55–64	286.3	..	211.3	234.0	272.2	..	351.2	129.9	248.6
65 years and over	184.8	..	160.0	232.0	170.9	..	384.9	89.8	182.1
Total males (a)	456.6	..	355.9	293.4	400.7	..	649.7	269.9	391.8
Females									
Less than 15 years	98.2	..	185.5	161.4	164.9	..	281.9	116.3	143.0
15–24	507.8	..	507.3	562.6	418.7	..	1 488.8	361.3	523.1
25–34	411.6	..	366.9	370.1	452.5	..	805.8	365.9	400.1

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
35–44	437.3	..	383.4	406.7	528.2	..	847.2	365.5	426.7
45–54	383.1	..	307.3	381.0	456.7	..	717.6	221.4	366.1
55–64	260.9	..	227.7	287.2	285.3	..	503.4	183.3	257.1
65 years and over	239.9	..	197.8	352.1	261.6	..	506.7	61.1	245.8
Total females (a)	330.9	..	312.6	356.0	364.3	..	747.6	239.3	335.8
Total people									
Less than 15 years	87.5	..	201.0	164.1	190.0	..	204.3	139.7	143.9
15–24	509.3	..	469.5	441.8	398.6	..	1 297.2	355.5	486.6
25–34	614.4	..	457.2	379.4	531.3	..	889.1	418.5	521.6
35–44	604.9	..	451.5	398.9	591.2	..	868.1	406.4	522.0
45–54	455.8	..	334.4	344.4	466.6	..	720.6	213.2	399.8
55–64	273.7	..	219.5	260.6	279.0	..	429.4	154.7	253.0
65 years and over	216.1	..	179.9	296.0	220.7	..	451.1	76.7	216.9
Total (b)	406.8	..	334.2	324.6	398.2	..	698.5	255.1	371.1
2013-14 (f), (g)									
Number									
Males									
Less than 15 years	48 440	42 652	99 002	41 639	28 635	6 559	5 158	4 856	276 941
15–24	296 551	139 876	153 097	63 076	45 543	9 808	33 863	7 533	749 347
25–34	454 441	175 610	183 249	85 720	66 068	13 173	32 500	11 742	1 022 503
35–44	459 504	191 273	187 016	73 838	71 737	13 846	29 251	9 045	1 035 510
45–54	289 002	128 748	117 433	52 889	54 398	12 808	19 787	4 208	679 273
55–64	132 928	65 651	61 860	32 779	29 575	7 434	8 710	1 716	340 653
65 years and over	88 687	85 871	50 266	29 636	22 314	6 634	6 683	974	291 065
Total males (a)	1 797 476	883 077	851 967	379 732	318 305	71 034	136 406	40 076	4 478 073
Females									

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Less than 15 years	63 318	40 830	87 878	40 280	23 177	7 922	12 989	3 438	279 832
15–24	265 279	187 694	173 423	103 558	48 406	14 183	46 767	8 539	847 849
25–34	236 614	141 781	126 423	72 847	51 522	10 962	29 200	8 409	677 758
35–44	245 042	153 300	132 448	75 205	55 584	13 733	25 439	7 051	707 802
45–54	202 139	117 491	102 779	66 512	54 615	12 865	20 234	3 885	580 520
55–64	121 977	76 158	62 534	43 391	30 144	7 468	11 587	2 080	355 339
65 years and over	139 272	137 905	68 192	54 611	40 072	11 697	13 854	812	466 415
Total females (a)	1 288 584	914 991	753 696	456 765	303 527	79 595	160 729	34 215	3 992 102
People									
Less than 15 years	111 761	83 504	186 880	81 922	51 812	14 481	18 147	8 294	556 801
15–24	562 031	327 852	326 624	166 634	93 976	23 991	80 630	16 072	1 597 810
25–34	691 392	317 459	309 685	158 571	117 617	24 135	61 701	20 152	1 700 712
35–44	704 665	344 736	319 464	149 046	127 354	27 579	54 690	16 096	1 743 630
45–54	491 727	246 261	220 270	119 403	109 028	25 673	40 021	8 093	1 260 476
55–64	255 045	141 809	124 407	76 170	59 719	14 902	20 297	3 796	696 145
65 years and over	229 764	223 776	118 458	84 256	62 395	18 331	20 537	1 786	759 303
Total (b)	3 179 782	1 898 524	1 605 851	844 650	662 195	150 701	297 136	74 292	8 713 131
<i>Rate (per 1000 population) (c)</i>									
Males									
Less than 15 years	67.0	78.5	207.3	167.9	188.6	133.8	140.9	174.8	122.7
15–24	593.4	353.8	468.0	352.2	405.7	289.2	1 185.7	384.9	469.5
25–34	852.2	401.1	549.9	402.1	581.4	451.4	983.6	480.5	595.0
35–44	909.8	477.7	582.3	400.1	658.9	441.1	1 040.4	468.1	647.5
45–54	595.0	343.2	382.3	310.1	479.2	363.5	815.0	255.4	444.4
55–64	311.8	206.8	235.9	235.7	291.4	214.5	444.9	135.0	259.3
65 years and over	167.5	219.8	164.1	197.4	173.0	155.6	329.9	115.8	184.6

TABLE 12A.24

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Total males (a)	485.5	308.7	364.9	295.6	383.2	277.4	716.3	311.4	386.4
Females									
Less than 15 years	92.7	79.2	194.4	169.2	160.6	172.9	373.6	131.2	130.8
15–24	557.8	496.6	549.3	621.0	454.3	455.3	1 716.5	525.6	558.8
25–34	441.8	323.0	379.1	371.6	465.6	369.3	889.3	398.4	399.0
35–44	474.1	373.1	403.5	424.2	509.7	423.4	905.7	397.5	436.7
45–54	406.2	302.9	326.6	395.5	473.3	353.2	803.2	257.0	371.9
55–64	278.8	229.4	234.8	310.9	285.5	213.8	558.9	187.2	263.6
65 years and over	226.8	299.2	199.9	319.9	259.2	246.9	575.0	114.4	256.3
Total females (a)	342.7	312.9	320.5	363.5	358.7	308.7	833.0	298.5	341.0
Total people									
Less than 15 years	79.5	78.8	201.0	168.5	174.9	152.7	254.3	153.6	126.6
15–24	576.2	423.9	508.1	481.8	429.5	368.7	1 444.9	448.7	513.2
25–34	646.9	362.1	464.5	387.5	524.4	410.0	936.6	442.5	497.7
35–44	689.6	424.9	492.0	411.9	584.4	432.1	973.1	434.3	541.5
45–54	500.0	322.7	354.2	352.5	476.3	358.3	809.0	256.2	408.0
55–64	295.3	218.3	235.4	273.4	288.4	214.2	503.5	159.3	261.5
65 years and over	200.9	262.7	183.0	262.6	220.0	203.7	463.0	115.1	223.5
Total (b)	426.1	328.2	342.7	332.4	394.9	293.3	775.0	305.3	374.0

(a) Includes service contacts for which age group was not reported.

(b) Includes service contacts for which sex and/or age group was not reported.

(c) Total rates for males, females and all were directly age-standardised for data up to 2012-13. Age rates are crude rates, as detailed in Technical information—see Technical notes section of *Mental Health Services in Australia* online.

Table 12A.24 **Community mental health service contacts, by sex and age group**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
--	------------	------------	------------	-----------	-----------	------------	------------	-----------	-------------

- (d) Data were not available for Victoria in 2011–12 or 2012-13 due to service level collection gaps resulting from protected industrial action during this period. Victoria required that data for 2011–12 and 2012-13 be excluded from all totals, with no proxy data to be included for Victoria when calculating national totals. Industrial action in Tasmania in 2011–12 and 2012-13 has affected the quality and quantity of Tasmania's data (see the online data source of the Community mental health care section).
- (e) Totals include only those jurisdictions that provided data. Rates for 2011-12 and 2012-13 were calculated using a methodology which accounts for missing data, as detailed in the online technical information. Comparisons between jurisdictions and over time should be made with caution.
- (f) SA transitioned to a new hospital-based system during 2013–14 which impacted on activity data reporting for a small number of hospital-based services.
- (g) Tasmania transitioned to a new clinical information system in 2013–14 which impacted on activity data reporting.

na Not available. .. Not applicable. – Nil or rounded to zero.

Source: AIHW (various issues) *Mental Health Services in Australia* (various years), (available at <http://mhsa.aihw.gov.au/home/>).

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
<i>2005-06</i>										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	2.0	3.7	..	np	3.6	18.5	np	..	1.9
Non-Indigenous	per 10 000 people	0.6	1.6	..	0.9	0.9	15.4	1.8	..	1.1
Rate ratio (f)		3.3	2.3	..	0.8	4.0	1.2	1.2	..	1.7
Total	per 10 000 people	0.6	1.6	..	0.9	0.9	15.2	1.8	..	1.2
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	659.2	750.3	378.6	346.3	393.3	61.7	1 170.8	183.3	458.0
Non-Indigenous (h)	per 1 000 people	161.1	360.5	223.3	226.7	182.4	99.0	398.9	169.0	234.5
Rate ratio (f)		4.1	2.1	1.7	1.5	2.2	0.6	2.9	1.1	2.0
Total	per 1 000 people	272.7	364.9	225.1	242.6	195.7	134.4	632.2	175.3	278.9
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separations	no.	1 709	316	1 007	623	328	np	np	np	4 478
Separation rate (d)	per 1 000 people	13.6	10.9	8.2	9.3	12.2	np	np	np	10.4
Non-Indigenous (h)										
Separations	no.	36 704	25 380	25 438	10 976	9 990	np	np	np	109 139
Separation rate (d)	per 1 000 people	5.5	5.0	6.6	5.6	6.4	np	np	np	5.7
Rate ratio (f)		2.5	2.2	1.2	1.7	1.9	np	np	np	1.8
<i>2006-07</i>										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Aboriginal and Torres Strait Islander (e)	per 10 000 people	1.8	10.3	..	np	0.8	15.4	1.6	np	1.8
Non-Indigenous	per 10 000 people	0.6	1.9	..	0.9	0.8	12.8	2.1	0.5	1.2
Rate ratio (f)		3.0	5.4	..	np	1.0	1.2	0.8	np	1.5
Total	per 10 000 people	0.6	2.0	..	0.9	0.8	15.1	2.4	0.4	1.2
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	777.8	810.7	517.1	331.8	464.0	135.8	791.7	176.1	536.1
Non-Indigenous (h)	per 1 000 people	197.8	352.8	254.9	246.7	224.7	160.3	518.4	165.2	259.6
Rate ratio (f)		3.9	2.3	2.0	1.3	2.1	0.8	1.5	1.1	2.1
Total	per 1 000 people	269.4	358.6	259.1	258.0	244.9	189.6	613.2	179.1	289.2
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	15.1	12.6	10.1	8.4	13.5	np	np	7.0	11.3
Non-Indigenous (h)										
Separation rate (d)	per 1 000 people	5.6	5.2	6.2	5.6	6.8	np	np	3.3	5.7
Rate ratio (f)		2.7	2.4	1.6	1.5	2.0	np	np	2.1	1.8
<i>2007-08</i>										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	..	np	np	np	np	np	1.9

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Non-Indigenous (h)	per 10 000 people	np	np	..	np	np	np	np	np	1.4
Rate ratio (f)		np	np	..	np	np	np	np	np	1.4
Total	per 10 000 people	0.4	2.9	..	1.1	1.2	18.3	2.2	0.2	1.5
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	1 013.9	787.3	608.0	370.0	627.8	142.6	883.2	169.1	644.3
Non-Indigenous (h)	per 1 000 people	230.4	327.6	273.0	249.4	259.6	230.3	514.2	135.8	271.7
Rate ratio (f)		4.4	2.4	2.2	1.5	2.4	0.6	1.7	1.2	2.4
Total	per 1 000 people	301.1	334.0	279.5	259.7	289.5	297.9	602.8	166.9	303.3
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	14.1	11.9	9.3	8.3	11.3	np	np	5.9	10.5
Non-Indigenous										
Separation rate (d)	per 1 000 people	5.6	5.5	5.9	6.0	6.0	np	np	3.4	5.7
Rate ratio (f)		2.5	2.2	1.6	1.4	1.9	np	np	1.7	1.8
<i>2008-09</i>										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	..	np	np	np	np	np	1.7
Non-Indigenous	per 10 000 people	np	np	..	np	np	np	np	np	1.5

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Rate ratio (f)		np	np	..	np	np	np	np	np	1.1
Total	per 10 000 people	0.3	3.3	..	1.1	1.5	19.3	1.3	2.2	1.6
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	952.9	819.5	497.1	432.7	806.2	210.3	946.2	187.6	623.2
Non-Indigenous	per 1 000 people	192.8	313.1	217.9	265.0	287.8	301.3	540.1	134.7	250.6
Rate ratio (f)		4.9	2.6	2.3	1.6	2.8	0.7	1.8	1.4	2.5
Total	per 1 000 people	293.0	317.9	224.3	275.8	328.7	346.3	636.1	176.7	292.0
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	10.6
Non-Indigenous (h)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	6.0
Rate ratio (f)		np	np	np	np	np	np	np	np	1.8
2009-10										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	np	np	np	np	np	np	2.5

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Non-Indigenous	per 10 000 people	np	np	np	np	np	np	np	np	1.7
Rate ratio (f)		np	np	np	np	np	np	np	np	1.5
Total	per 10 000 people	0.3	4.1	..	1.0	1.4	18.3	1.6	3.6	1.8
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	1 199.2	797.2	466.0	495.0	789.8	1 010.5	1 485.3	216.2	730.9
Non-Indigenous (h)	per 1 000 people	235.6	313.9	195.8	285.8	292.3	381.0	642.3	141.3	266.1
Rate ratio (f)		5.1	2.5	2.4	1.7	2.7	2.7	2.3	1.5	2.7
Total	per 1 000 people	315.7	320.3	202.3	300.4	335.7	419.8	719.5	171.1	301.6
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	10.6
Non-Indigenous (h)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	5.9
Rate ratio (f)		np	np	np	np	np	np	np	np	1.8
2010-11										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	np	np	np	np	np	np	2.6
Non-Indigenous	per 10 000 people	np	np	np	np	np	np	np	np	1.8
Rate ratio (f)		np	np	np	np	np	np	np	np	1.4

TABLE 12A.25

Table 12A.25 Specialised mental health care reported, by Indigenous status

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Total	per 10 000 people	0.3	4.5	..	1.0	2.3	14.9	2.1	3.7	1.9
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	1 240.3	724.7	557.4	603.8	811.3	244.2	1 449.9	242.1	755.5
Non-Indigenous (h)	per 1 000 people	255.0	318.4	222.1	307.7	298.9	252.3	588.3	142.8	277.9
Rate ratio (f)		4.9	2.3	2.5	2.0	2.7	1.0	2.5	1.7	2.7
Total	per 1 000 people	335.4	363.0	230.7	324.3	343.3	295.3	665.7	179.0	323.6
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	12.9
Non-Indigenous (h)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	5.8
Rate ratio (f)		np	np	np	np	np	np	np	np	2.2
2011-12 (m), (n)										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	np	np	np	np	np	np	5.0
Non-Indigenous	per 10 000 people	np	np	np	np	np	np	np	np	2.4
Rate ratio (f)		np	np	np	np	np	np	np	np	2.1
Total	per 10 000 people	0.4	5.0	..	1.2	7.4	18.5	1.8	5.2	2.5
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d), (m), (n)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	1 438.4	np	701.3	636.2	935.8	166.5	1 640.1	267.8	877.2

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Non-Indigenous (h)	per 1 000 people	265.9	np	257.0	300.0	311.3	164.2	619.3	175.5	276.8
Rate ratio (f)		5.4	np	2.7	2.1	3.0	1.0	2.6	1.5	3.2
Total	per 1 000 people	354.4	np	268.7	314.6	360.2	192.9	698.8	210.3	327.1
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	13.9
Non-Indigenous (h)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	6.1
Rate ratio (f)		np	np	np	np	np	np	np	np	2.3
2012-13 (m), (n)										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	np	np	np	np	np	np	4.1
Non-Indigenous	per 10 000 people	np	np	np	np	np	np	np	np	2.8
Rate ratio (f)		np	np	np	np	np	np	np	np	1.5
Total	per 10 000 people	0.4	5.3	..	1.1	10.3	21.4	1.8	4.3	2.9
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d), (m), (n)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	1 546.7	na	889.0	704.3	928.8	118.9	1 710.4	374.5	991.1
Non-Indigenous (h)	per 1 000 people	302.5	na	311.7	300.8	335.0	133.5	652.4	212.9	309.9
Rate ratio (f)		5.1	na	2.9	2.3	2.8	0.9	2.6	1.8	3.2

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Total	per 1 000 people	397.5	na	326.9	320.7	384.9	132.1	708.8	267.9	363.4
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	12.7
Non-Indigenous (h)										
Separation rate (d)	per 1 000 people	np	np	np	np	np	np	np	np	6.3
Rate ratio (f)		np	np	np	np	np	np	np	np	2.0
2013-14 (m), (n)										
<i>Episodes of community-based residential mental health care (a), (b), (c)</i>										
Rate per 10 000 people (d)										
Aboriginal and Torres Strait Islander (e)	per 10 000 people	np	np	np	np	np	np	np	np	4.8
Non-Indigenous	per 10 000 people	np	np	np	np	np	np	np	np	2.9
Rate ratio (f)		np	np	np	np	np	np	np	np	1.7
Total	per 10 000 people	0.3	6.4	..	1.2	10.6	14.6	1.5	4.7	3.0
<i>Community-based ambulatory mental health service contacts (g)</i>										
Rate per 1000 people (d), (m), (n)										
Aboriginal and Torres Strait Islander (e)	per 1 000 people	1 729.3	709.0	914.7	845.4	890.8	284.5	2 242.9	422.2	1 066.3

TABLE 12A.25

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
Non-Indigenous (h)	per 1 000 people	339.9	283.9	327.5	311.6	336.1	295.4	726.7	244.4	324.3
Rate ratio (f)		5.1	2.5	2.8	2.7	2.7	1.0	3.1	1.7	3.3
Total	per 1 000 people	426.1	328.2	342.7	332.4	394.9	293.3	775.0	305.3	374.0
<i>Admitted patient mental health-related separations with specialised psychiatric care (i), (j), (k), (l)</i>										
Aboriginal and Torres Strait Islander (e)										
Separation rate (d)	per 1 000 people	na	na	na	na	na	na	na	na	na
Non-Indigenous (h)		na	na	na	na	na	na	na	na	na
Separation rate (d)	per 1 000 people	na	na	na	na	na	na	na	na	na
Rate ratio (f)		na	na	na	na	na	na	na	na	na

- (a) Data for episodes of community residential care should be interpreted with caution due to the varying quality and completeness of Indigenous identification across jurisdictions.
- (b) Queensland does not have any government-operated residential mental health services. Tasmanian information contains data for government-funded residential units operated by the non-government sector in that state, being the only jurisdiction providing this level of reporting. The NT did not have any community residential units in 2005-06.
- (c) For NSW, Confused and Disturbed Elderly (CADE) residential mental health services were reclassified as admitted patient hospital services from 1 July 2007. All data relating to these services have been reclassified from 2007-08 onwards. Comparison of NSW data over time therefore should be approached with caution.
- (d) The rates were directly aged standardised against the Australian Estimated Resident Population as at 30 June 2001.
- (e) Includes patients identified as being either of Aboriginal but not Torres Strait Islander origin, Torres Strait Islander but not Aboriginal origin, Aboriginal and Torres Strait Islander origin and patients identified as of Aboriginal or Torres Strait Islander origin. For admitted patient care, separations from private hospitals are also included.
- (f) The rate ratio is equal to the service use (episodes, contacts or separations) rate for Indigenous Australians divided by the service use rate for non-Indigenous Australians.
- (g) Data for community mental health service contacts should be interpreted with caution. Across jurisdictions, the data quality and completeness of Indigenous identification varies or is unknown. See *Mental Health Services in Australia* (<http://mhsa.aihw.gov.au/home>) for further information.
- (h) Includes data for people where Indigenous status was missing or not reported.

Table 12A.25 **Specialised mental health care reported, by Indigenous status**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
(i)	Admitted patient separations refers to those non-ambulatory separations when a patient undergoes a hospital's formal admission process, completes an episode of care and 'separates' from the hospital, excluding ambulatory-equivalent separations. Separations for which care type was reported as Newborn with no qualified days and records for Hospital boarders and Posthumous organ procurement have been excluded. Comprises separations with and without mental health-related principal diagnoses but with specialised psychiatric care.									
(j)	Interpretation of differences between jurisdictions needs to be undertaken with care as they may reflect different service delivery and admission practices and/or differences in the types of establishments categorised as hospitals.									
(k)	Includes only public hospital separations for the NT.									
(l)	Indigenous status data for NSW, Victoria, Queensland, WA, SA and the NT public hospitals are considered to be of acceptable quality for analytical purposes. Indigenous identification is likely to be incomplete and to vary among jurisdictions. Total includes data for these jurisdictions only.									
(m)	Data were not available for Victoria in 2011–12 and 2012-13 due to service level collection gaps resulting from protected industrial action during this period. Victoria required that data for 2011–12 and 2012-13 be excluded from all totals, with no proxy data to be included for Victoria when calculating national totals. Industrial action in Tasmania in 2011–12 and 2012-13 has affected the quality and quantity of Tasmania's data (see the Mental Health Services in Australia online data source of the Community mental health care section).									
(n)	Totals include only those jurisdictions that provided data. Rates were calculated using a methodology which accounts for missing data, as detailed in the online technical information. Comparisons between jurisdictions and over time should be made with caution.									

na Not available. .. Not applicable. – Nil or rounded to zero. **np** Not published.

Source: AIHW (various issues) *Mental Health Services in Australia* (various years), (available at <http://mhsa.aihw.gov.au/home/>).

Table 12A.26 Available beds in State and Territory governments' specialised mental health services (a), (b), (c), (d)

	NSW (e), (f)	Vic	Qld (g)	WA (h)	SA	Tas (i), (j)	ACT (j)	NT (j)	Aust
<i>No. of beds</i>									
Psychiatric hospitals									
2005-06	1 072	116	375	245	455	2 263
2006-07	1 060	134	375	254	388	2 211
2007-08	1 024	154	376	245	357	2 156
2008-09	911	154	375	246	343	2 029
2009-10	967	150	375	243	267	2 002
2010-11	1 064	152	375	246	247	2 083
2011-12	902	150	345	246	230	1 873
2012-13	887	152	345	242	205	1 831
2013-14	854	152	335	226	194	1 761
Acute hospitals with psychiatric units or wards									
2005-06	1 151	1 048	1 014	403	188	125	50	32	4 011
2006-07	1 227	1 050	1 022	415	247	126	70	34	4 191
2007-08	1 400	1 062	1 033	425	243	128	70	34	4 395
2008-09	1 542	1 064	1 029	432	233	130	63	34	4 527
2009-10	1 558	1 082	1 033	452	246	128	63	34	4 597
2010-11	1 586	1 104	1 044	454	252	127	65	33	4 666
2011-12	1 747	1 091	1 057	463	250	131	65	32	4 836
2012-13	1 768	1 092	1 110	479	247	131	70	40	4 937
2013-14	1 838	1 122	1 066	499	268	127	70	41	5 030
Community-based residential units									
2005-06	440	1 319	..	80	43	174	80	10	2 146
2006-07	437	1 359	..	85	63	176	75	5	2 200
2007-08	251	1 404	..	130	71	176	77	5	2 114
2008-09	196	1 456	..	178	99	165	83	13	2 190
2009-10	195	1 430	..	260	89	169	83	13	2 239
2010-11	185	1 448	..	283	97	170	83	15	2 281
2011-12	171	1 476	..	303	138	162	82	15	2 347
2012-13	158	1 495	..	298	137	156	95	16	2 356
2013-14	156	1 536	..	315	147	166	95	22	2 437
<i>Proportion of all beds in different settings (%)</i>									
Psychiatric hospitals									
2005-06	40.3	4.7	27.0	33.7	66.3	26.9
2006-07	38.9	5.3	26.8	33.7	55.6	25.7
2007-08	38.3	5.9	26.7	30.6	53.2	24.9
2008-09	34.4	5.8	26.7	28.7	50.8	23.2
2009-10	35.6	5.6	26.6	25.4	44.3	22.7
2010-11	37.5	5.6	26.4	25.0	41.4	23.1
2011-12	32.0	5.5	24.6	24.3	37.2	20.7

Table 12A.26 Available beds in State and Territory governments' specialised mental health services (a), (b), (c), (d)

	<i>NSW (e), (f)</i>	<i>Vic</i>	<i>Qld (g)</i>	<i>WA (h)</i>	<i>SA</i>	<i>Tas (i), (j)</i>	<i>ACT (j)</i>	<i>NT (j)</i>	<i>Aust</i>
2012-13	31.5	5.5	23.7	23.7	34.8	20.1
2013-14	30.0	5.4	23.9	21.7	31.9	19.1
Acute hospitals with psychiatric units or wards									
2005-06	43.2	42.2	73.0	55.4	27.4	41.8	38.5	76.2	47.6
2006-07	45.0	41.3	73.2	55.0	35.4	41.7	48.3	87.2	48.7
2007-08	52.3	40.5	73.3	53.1	36.2	42.1	47.6	87.2	50.7
2008-09	58.2	39.8	73.3	50.5	34.5	44.1	43.2	72.3	51.8
2009-10	57.3	40.6	73.4	47.3	40.9	43.1	43.2	72.3	52.0
2010-11	55.9	40.8	73.6	46.2	42.3	42.8	43.9	68.9	51.7
2011-12	61.9	40.2	75.4	45.8	40.5	44.7	44.2	68.1	53.4
2012-13	62.9	39.9	76.3	47.0	41.8	45.6	42.4	71.4	54.1
2013-14	64.5	39.9	76.1	48.0	44.0	43.3	42.4	65.1	54.5
Community-based residential units									
2005-06	16.5	53.1	..	11.0	6.3	58.2	61.5	23.8	25.5
2006-07	16.0	53.4	..	11.3	9.0	58.3	51.7	12.8	25.6
2007-08	9.4	53.6	..	16.3	10.6	57.9	52.4	12.8	24.4
2008-09	7.4	54.5	..	20.8	14.7	55.9	56.8	27.7	25.0
2009-10	7.2	53.7	..	27.2	14.8	56.9	56.8	27.7	25.3
2010-11	6.5	53.6	..	28.8	16.3	57.2	56.1	31.1	25.3
2011-12	6.1	54.3	..	29.9	22.3	55.3	55.8	31.9	25.9
2012-13	5.6	54.6	..	29.3	23.3	54.4	57.6	28.6	25.8
2013-14	5.5	54.7	..	30.3	24.1	56.7	57.6	34.9	26.4
<i>Beds per 100 000 people</i>									
Psychiatric hospitals									
2005-06	16.0	2.3	9.5	12.1	29.5	11.1
2006-07	15.6	2.6	9.2	12.2	24.9	10.7
2007-08	14.9	3.0	9.0	11.5	22.6	10.3
2008-09	13.0	2.9	8.8	11.1	21.5	9.4
2009-10	13.6	2.8	8.6	10.7	16.5	9.2
2010-11	14.8	2.8	8.5	10.6	15.1	9.4
2011-12	12.4	2.7	7.6	10.3	14.0	8.3
2012-13	12.1	2.7	7.5	9.8	12.4	8.0
2013-14	11.4	2.6	7.1	8.9	11.6	7.6
Acute hospitals with psychiatric units or wards									
2005-06	17.1	20.9	25.6	19.9	12.2	25.6	15.0	15.4	19.7
2006-07	18.1	20.6	25.2	20.0	15.8	25.6	20.7	16.1	20.3
2007-08	20.3	20.4	24.8	19.9	15.4	25.8	20.3	15.7	20.9
2008-09	22.0	20.0	24.1	19.6	14.6	25.9	17.9	15.3	21.1
2009-10	21.9	20.0	23.7	20.0	15.2	25.3	17.6	14.9	21.0
2010-11	22.1	20.1	23.5	19.6	15.4	24.9	17.8	14.5	21.0

Table 12A.26 Available beds in State and Territory governments' specialised mental health services (a), (b), (c), (d)

	NSW (e), (f)	Vic	Qld (g)	WA (h)	SA	Tas (i), (j)	ACT (j)	NT (j)	Aust
2011-12	24.1	19.6	23.4	19.4	15.2	25.6	17.5	13.8	21.5
2012-13	24.1	19.2	24.1	19.4	14.8	25.6	18.4	16.9	21.6
2013-14	24.6	19.4	22.7	19.6	16.0	24.7	18.2	16.9	21.6
Community-based residential units									
2005-06	6.5	26.3	..	3.9	2.8	35.6	24.0	4.8	10.6
2006-07	6.4	26.6	..	4.1	4.0	35.8	22.2	2.4	10.7
2007-08	3.6	27.0	..	6.1	4.5	35.5	22.4	2.3	10.1
2008-09	2.8	27.4	..	8.1	6.2	32.9	23.6	5.8	10.2
2009-10	2.7	26.4	..	11.5	5.5	33.3	23.2	5.7	10.2
2010-11	2.6	26.3	..	12.2	6.0	33.3	22.8	6.5	10.3
2011-12	2.4	26.5	..	12.7	8.4	31.7	22.1	6.5	10.4
2012-13	2.1	26.3	..	12.1	8.3	30.4	25.0	6.8	10.3
2013-14	2.1	26.5	..	12.3	8.8	32.3	24.7	9.1	10.4
Total									
2005-06	39.6	49.4	35.0	35.9	44.4	61.3	39.0	20.3	41.5
2006-07	40.1	49.8	34.4	36.3	44.7	61.4	42.9	18.5	41.7
2007-08	38.9	50.4	33.9	37.5	42.5	61.3	42.7	18.0	41.2
2008-09	37.8	50.3	32.8	38.8	42.2	58.8	41.6	21.1	40.7
2009-10	38.3	49.1	32.2	42.2	37.2	58.6	40.8	20.6	40.4
2010-11	39.5	49.2	32.0	42.4	36.5	58.2	40.6	21.0	40.7
2011-12	38.9	48.7	31.1	42.4	37.6	57.3	39.7	20.2	40.3
2012-13	38.3	48.2	31.6	41.2	35.5	56.0	43.5	23.6	39.8
2013-14	38.1	48.5	29.9	40.8	36.3	57.0	43.0	26.0	39.6

- (a) Bed numbers represent the average number of beds which are immediately available for use by an admitted patient or resident within the establishment. See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the bed estimates. Available beds are counted as the average of monthly available bed numbers. Available beds counts exclude beds in wards that were closed for any reason (except weekend closures for beds/wards staffed and available on weekdays only).
- (b) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (c) Hospital bed can include government funded beds managed and operated by private and non-government entities.
- (d) Community-based residential beds data include 24-hour and non-24-hour staffed units.
- (e) Caution is required when interpreting NSW data. Seven residential mental health services in 2006–07 were reclassified as non-acute older person specialised hospital services in 2007–08, reflecting a change in function of those units.
- (f) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.

Table 12A.26 Available beds in State and Territory governments' specialised mental health services (a), (b), (c), (d)

	<i>NSW</i> (e), (f)	<i>Vic</i>	<i>Qld</i> (g)	<i>WA</i> (h)	<i>SA</i>	<i>Tas</i> (i), (j)	<i>ACT</i> (j)	<i>NT</i> (j)	<i>Aust</i>
(g)	Queensland does not fund community residential services, however, it funds a number of campus based and non-campus based extended treatment services. These services are reported either as wards of public acute hospitals or beds in public psychiatric hospitals. Furthermore, limiting the classification of all inpatient beds to either co-located or standalone results in the reporting of some psychogeriatric beds co-located with nursing homes being reported as 'standalone' which results in the reporting of these beds as psychiatric hospital beds in this report. In 2005-06, there was temporary closure of acute beds in one Queensland hospital and some transitional extended treatment beds were permanently closed. In addition, Queensland did not change its method for counting beds until 2007-08 (see 2011 Report for details of previous method).								
(h)	Beds numbers in WA include publicly funded mental health beds in private hospitals for all years. Bed numbers in WA include emergency department observation beds in one hospital for all years prior to 2010-11.								
(i)	In Tasmania, for 2005-06, non-government organisations' residential beds funded by government were included for the first time in the publicly funded community residential facilities category.								
(j)	Tasmania, the ACT and the NT do not have public psychiatric hospitals. .. Not applicable.								

Source: AIHW (unpublished) MHE NMDS; table 12A.99.

TABLE 12A.27

Table 12A.27 Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people) (a), (b), (c)

	NSW (d)	Vic	Qld (e)	WA	SA	Tas	ACT	NT	Aust
2005-06									
Medical									
Consultant psychiatrist	5.5	4.6	4.3	4.9	5.1	5.0	3.2	3.7	4.9
Psychiatry registrar	4.7	4.7	5.0	4.4	5.7	3.0	5.1	3.1	4.8
Other medical officers	0.8	1.9	0.8	2.9	1.6	0.8	1.3	2.7	1.4
Total	11.0	11.2	10.1	12.2	12.4	8.8	9.7	9.5	11.0
Nursing									
Registered nursing	53.1	53.7	47.8	63.2	56.9	60.4	37.8	40.6	53.3
Non-registered	8.3	12.0	7.6	7.9	14.5	10.3	7.0	3.9	9.5
Total	61.4	65.7	55.4	71.1	71.4	70.7	44.8	44.4	62.8
Allied health									
Occupation therapist	3.2	4.4	3.6	5.9	3.0	1.8	2.1	0.5	3.8
Social worker	5.3	7.9	6.9	8.4	12.5	4.0	7.4	2.4	7.1
Psychologist	8.5	7.8	7.7	7.1	5.4	5.6	21.9	5.8	7.9
Other allied health staff	5.1	2.2	2.9	5.7	3.6	5.9	4.1	8.2	3.9
Total	22.1	22.3	21.1	27.2	24.6	17.3	35.5	16.9	22.7
Other personal care	1.8	5.1	4.7	4.4	0.9	27.7	8.9	2.4	4.1
Total	96.2	104.3	91.4	114.9	109.3	124.5	98.9	73.2	100.6
2006-07									
Medical									
Consultant psychiatrist	5.5	4.6	4.6	4.8	5.8	4.5	3.8	3.9	5.0
Psychiatry registrar	5.4	4.5	5.6	4.8	6.1	2.8	4.5	4.0	5.1
Other medical officers	0.7	1.6	0.8	3.4	1.6	1.0	0.5	2.2	1.3
Total	11.6	10.7	11.1	12.9	13.5	8.4	8.8	10.1	11.4
Nursing									
Registered nursing	54.4	52.0	50.1	61.6	61.1	65.0	41.7	41.8	54.1
Non-registered	8.2	14.1	7.5	8.7	13.8	10.6	8.4	4.5	10.0
Total	62.5	66.1	57.6	70.3	74.9	75.6	50.1	46.3	64.1
Allied health									
Occupation therapist	3.3	4.7	3.5	6.3	3.6	3.0	1.8	0.5	3.9
Social worker	5.2	8.2	7.0	9.5	12.7	6.1	5.9	3.4	7.3
Psychologist	8.3	8.3	8.1	8.1	5.1	5.5	17.9	5.9	8.1
Other allied health staff	5.4	1.7	2.9	5.3	3.9	5.8	2.0	5.7	3.8
Total	22.2	22.9	21.6	29.3	25.3	20.3	27.7	15.4	23.2
Other personal care	2.4	4.2	5.0	4.3	1.5	29.2	8.5	3.3	4.2
Total	98.7	103.9	95.3	116.8	115.2	133.5	95.1	75.1	102.9
2007-08									
Medical									
Consultant psychiatrist	5.6	4.3	5.7	4.8	6.1	5.4	4.3	4.0	5.2

TABLE 12A.27

Table 12A.27 **Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people) (a), (b), (c)**

	<i>NSW (d)</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Psychiatry registrar	5.4	4.8	5.6	4.8	6.6	3.2	4.8	4.1	5.2
Other medical officers	1.2	2.2	0.9	3.6	1.1	0.2	0.4	2.3	1.6
Total	12.1	11.3	12.1	13.2	13.8	8.9	9.5	10.4	12.0
Nursing									
Registered nursing	54.4	51.0	52.2	61.1	63.8	59.2	40.6	42.7	54.3
Non-registered	8.1	14.4	8.3	8.8	15.0	11.0	8.6	4.2	10.3
Total	62.5	65.4	60.4	70.0	78.7	70.2	49.2	46.9	64.6
Allied health									
Occupation therapist	3.3	4.5	3.9	6.8	4.4	2.2	2.2	0.8	4.1
Social worker	4.7	8.4	7.7	9.5	14.3	5.6	6.8	3.8	7.4
Psychologist	8.7	7.5	9.5	7.4	6.9	5.0	14.2	5.8	8.3
Other allied health staff	5.6	2.2	3.1	5.7	4.4	5.9	0.2	6.1	4.1
Total	22.3	22.6	24.2	29.5	30.0	18.7	23.4	16.5	23.9
Other personal care	1.0	4.7	4.8	6.2	1.5	31.2	9.4	3.6	4.1
Total	97.9	104.0	101.5	118.8	124.0	129.0	91.4	77.3	104.6
<i>2008-09</i>									
Medical									
Consultant psychiatrist	6.2	4.8	5.7	5.3	6.1	4.6	5.8	5.8	5.6
Psychiatry registrar	5.8	5.1	5.9	4.7	7.1	2.8	4.5	4.3	5.5
Other medical officers	0.9	1.5	0.5	3.5	0.2	2.7	0.8	2.7	1.3
Total	12.9	11.4	12.1	13.5	13.4	10.1	11.1	12.8	12.4
Nursing									
Registered nursing	56.0	50.9	52.7	61.9	62.3	60.2	37.4	46.6	54.9
Non-registered	8.2	15.4	8.4	9.3	15.3	11.0	9.3	2.4	10.7
Total	64.2	66.2	61.1	71.2	77.6	71.2	46.6	48.9	65.5
Allied health									
Occupation therapist	3.9	4.6	4.5	6.8	4.2	3.3	4.2	1.0	4.5
Social worker	5.7	8.8	7.8	9.2	14.7	5.9	5.9	5.2	7.9
Psychologist	8.3	7.8	9.3	7.9	5.6	4.6	13.5	4.5	8.1
Other allied health staff	4.0	1.7	3.7	6.2	3.0	4.5	0.4	5.0	3.5
Total	22.0	22.9	25.3	30.0	27.5	18.3	24.0	15.7	24.0
Other personal care	0.8	4.8	4.1	6.0	4.7	30.2	7.7	9.4	4.2
Total	99.9	105.5	102.7	120.7	123.2	129.9	89.4	86.9	106.1
<i>2009-10</i>									
Medical									
Consultant psychiatrist	6.1	5.4	5.6	5.6	6.2	5.4	6.8	5.8	5.8
Psychiatry registrar	6.6	4.7	5.8	4.5	6.8	2.8	5.2	4.6	5.6
Other medical officers	0.4	1.7	0.7	3.5	0.4	2.2	0.5	2.3	1.1
Total	13.1	11.9	12.1	13.7	13.4	10.4	12.5	12.7	12.6

TABLE 12A.27

Table 12A.27 **Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people) (a), (b), (c)**

	<i>NSW (d)</i>	<i>Vic Qld (e)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Nursing									
Registered nursing	59.3	50.3	50.0	60.5	62.9	58.3	40.9	45.5	55.1
Non-registered	7.3	15.3	7.5	9.7	14.6	11.1	8.5	3.3	10.2
Total	66.6	65.6	57.5	70.3	77.6	69.4	49.4	48.8	65.3
Allied health									
Occupation therapist	4.1	4.8	4.1	6.4	4.7	2.9	3.2	1.4	4.5
Social worker	6.0	8.6	8.6	8.8	15.0	6.2	6.0	4.2	8.1
Psychologist	8.4	7.6	8.7	7.2	5.6	4.7	14.4	4.7	7.9
Other allied health staff	4.1	1.6	3.6	6.4	2.5	6.6	0.4	5.7	3.5
Total	22.5	22.5	25.1	28.8	27.8	20.4	24.0	16.0	24.0
Other personal care	0.9	4.6	4.8	7.4	5.9	34.3	7.6	8.7	4.6
Total	103.1	104.6	99.4	120.1	124.7	134.5	93.5	86.2	106.5
<i>2010-11</i>									
Medical									
Consultant psychiatrist	6.3	5.5	6.0	6.0	7.6	6.4	6.0	4.5	6.1
Psychiatry registrar	6.8	5.0	5.8	4.3	5.8	2.5	5.1	5.6	5.7
Other medical officers	0.5	1.5	1.0	3.8	0.3	1.9	0.2	2.1	1.2
Total	13.6	12.0	12.9	14.1	13.7	10.7	11.3	12.2	13.0
Nursing									
Registered nursing	61.0	51.7	53.2	59.6	65.7	59.1	40.4	45.1	56.8
Non-registered	6.6	15.3	7.3	9.7	14.9	10.5	7.6	3.3	9.9
Total	67.6	67.0	60.5	69.3	80.6	69.6	48.0	48.3	66.7
Allied health									
Occupation therapist	4.2	5.2	4.5	6.4	4.7	3.0	3.3	1.2	4.7
Social worker	6.4	8.7	8.9	9.0	14.6	6.4	6.1	5.9	8.3
Psychologist	8.8	7.7	9.2	7.4	5.8	4.3	16.8	7.2	8.3
Other allied health staff	4.7	1.9	3.9	6.1	2.5	7.2	0.2	4.2	3.8
Total	24.1	23.4	26.6	28.9	27.6	20.9	26.4	18.5	25.1
Other personal care	0.6	4.5	5.3	10.7	7.2	33.2	7.1	10.1	5.0
Total	105.9	106.9	105.3	123.0	129.1	134.5	92.8	89.1	109.8
<i>2011-12</i>									
Medical									
Consultant psychiatrist	5.9	5.2	6.2	6.2	7.1	5.8	8.2	6.7	6.0
Psychiatry registrar	6.0	5.3	6.6	4.8	6.2	3.0	4.6	6.3	5.8
Other medical officers	1.3	1.7	0.8	3.6	0.3	1.1	0.2	0.4	1.4
Total	13.2	12.3	13.6	14.6	13.6	10.0	13.0	13.4	13.2
Nursing									
Registered nursing	63.0	52.0	56.0	60.5	63.1	56.8	38.7	47.7	57.9
Non-registered	7.5	15.6	8.0	10.0	13.4	9.2	7.6	3.7	10.3

TABLE 12A.27

Table 12A.27 **Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people) (a), (b), (c)**

	<i>NSW (d)</i>	<i>Vic Qld (e)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Total	70.5	67.5	64.0	70.5	76.5	66.0	46.4	68.2
Allied health								
Occupation therapist	3.8	5.6	4.5	5.9	4.5	2.5	3.6	4.6
Social worker	6.2	8.6	9.2	9.0	14.8	6.1	6.8	8.4
Psychologist	8.7	7.6	9.7	7.4	5.9	3.9	16.1	8.3
Other allied health staff	5.3	1.9	3.9	7.2	0.9	6.0	0.2	4.6
Total	24.1	23.6	27.4	29.6	26.0	18.5	26.6	25.2
Other personal care	0.6	4.4	5.0	11.5	7.4	31.6	7.0	5.0
Total	108.4	107.8	110.0	126.3	123.6	126.0	93.1	111.6
<i>2012-13</i>								
Medical								
Consultant psychiatrist	5.8	4.9	6.1	6.1	7.2	5.5	8.4	5.8
Psychiatry registrar	7.3	5.8	6.7	3.9	6.3	4.1	4.8	6.2
Other medical officers	0.9	1.7	0.7	5.2	0.4	1.2	0.2	1.5
Total	14.0	12.4	13.5	15.1	13.9	10.7	13.4	13.5
Nursing								
Registered nursing	63.6	51.4	54.8	62.4	61.7	55.3	39.6	57.8
Non-registered	6.4	14.2	8.0	9.8	12.3	9.2	7.9	9.5
Total	69.9	65.6	62.8	72.1	74.0	64.5	47.6	67.3
Allied health								
Occupation therapist	4.0	5.3	4.6	6.6	4.6	2.6	3.5	4.7
Social worker	6.6	8.1	8.9	10.0	14.9	5.5	6.5	8.4
Psychologist	8.4	7.3	9.4	7.4	5.9	5.7	16.4	8.1
Other allied health staff	5.4	2.2	5.6	5.6	1.0	5.8	0.2	4.3
Total	24.4	22.9	28.5	29.6	26.5	19.6	26.5	25.4
Other personal care	0.6	4.1	5.2	10.1	7.0	27.7	11.4	4.8
Total	108.9	105.0	109.9	127.1	121.4	122.5	98.9	111.1
<i>2013-14</i>								
Medical								
Consultant psychiatrist	6.1	4.9	5.9	6.1	7.7	6.7	9.4	5.9
Psychiatry registrar	7.0	6.3	6.9	3.9	6.8	3.7	4.9	6.4
Other medical officers	0.5	1.5	0.7	4.7	0.2	1.3	0.2	1.3
Total	13.7	12.7	13.6	14.7	14.8	11.7	14.5	13.6
Nursing								
Registered nursing	64.6	52.7	51.9	61.6	62.1	55.7	41.1	57.9
Non-registered	5.5	14.3	7.7	9.7	12.6	9.2	9.0	9.2
Total	70.1	67.0	59.6	71.3	74.7	64.9	50.2	67.1
Allied health								
Occupation therapist	4.1	5.2	4.5	6.1	4.9	3.0	3.7	4.7

Table 12A.27 Full time equivalent (FTE) direct care staff employed in specialised mental health services by staff type (per 100 000 people) (a), (b), (c)

	<i>NSW (d)</i>	<i>Vic Qld (e)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Social worker	6.6	8.9	8.5	9.1	16.2	8.3	6.0	7.3	8.6
Psychologist	8.3	7.3	9.1	7.0	4.9	4.7	15.6	5.5	7.9
Other allied health staff	5.7	2.4	3.7	6.4	1.5	3.3	0.2	5.8	4.1
Total	24.8	23.8	25.9	28.7	27.5	19.4	25.4	19.3	25.2
Other personal care	0.5	4.1	3.8	10.1	7.4	30.1	11.6	20.1	4.6
Total	109.0	107.6	102.9	124.9	124.3	126.2	101.7	111.7	110.5

- (a) Professional categories are defined by profession rather than role. See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of staffing estimates.
- (b) Total FTE figures presented in this table can differ from those in table 12A.28. A new organisational overhead setting for reporting FTE was implemented from the 2012-13 collection period, which may result in decreased FTE in the other service setting categories for some jurisdictions. Time series analyses should be approached with caution. Care and consumer worker FTE has been included in service setting reporting since the 2012-13 collection period. These categories are not included in staff type FTE data. Comparisons between these tables should be made with caution.
- (c) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (d) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.
- (e) Queensland implemented a new method to calculate FTE from the 2009-10 data. The new method is associated with the reduction in reported FTE so caution should be exercised when conducting time series analysis.

Source: AIHW (unpublished) MHE NMDS; table 12A.99.

TABLE 12A.28

Table 12A.28 **FTE direct care staff employed in specialised mental health services, by service setting (per 100 000 people) (a), (b), (c)**

	<i>NSW (d), (e)</i>	<i>Vic Qld (f), (g)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
2005-06									
Admitted patient	53.5	37.0	55.5	62.5	62.9	48.8	25.4	30.6	50.6
Ambulatory	38.7	44.1	35.9	49.5	45.0	40.9	50.0	40.6	41.3
Residential	4.0	22.9	..	2.9	1.4	34.6	19.2	1.9	8.6
Total	96.2	103.9	91.4	114.9	109.3	124.3	94.5	73.2	100.4
2006-07									
Admitted patient	55.6	37.3	54.7	63.9	67.4	58.6	28.2	32.3	52.0
Ambulatory	38.8	44.7	40.6	49.7	46.4	40.2	50.5	41.3	42.5
Residential	4.3	21.9	..	3.3	1.4	32.3	16.4	1.4	8.3
Total	98.7	103.9	95.3	116.8	115.2	131.1	95.1	75.0	102.9
2007-08									
Admitted patient	55.8	37.5	57.0	63.9	70.1	56.5	28.3	31.9	52.7
Ambulatory	39.9	44.2	44.5	49.4	50.9	38.9	49.0	43.8	43.8
Residential	2.3	22.2	..	5.5	3.0	31.6	14.0	1.7	8.0
Total	97.9	104.0	101.5	118.8	124.0	126.9	91.4	77.4	104.6
2008-09									
Admitted patient	57.9	38.6	55.8	64.8	67.1	56.6	26.4	38.0	53.4
Ambulatory	40.1	44.6	46.9	49.6	51.2	40.9	48.7	42.8	44.6
Residential	1.8	22.2	..	6.4	5.0	30.3	14.2	6.1	8.1
Total	99.9	105.5	102.7	120.7	123.2	127.8	89.4	86.9	106.0
2009-10									
Admitted patient	59.8	38.5	51.8	63.8	64.2	57.6	28.5	36.6	52.9
Ambulatory	41.7	44.6	47.6	49.4	55.5	42.2	50.1	43.1	45.6
Residential	1.6	21.5	..	6.9	5.0	32.9	14.9	6.4	8.0
Total	103.1	104.6	99.4	120.1	124.7	132.6	93.5	86.1	106.4
2010-11									
Admitted patient	61.2	39.4	53.6	64.1	62.5	58.3	29.8	38.0	53.8
Ambulatory	43.4	46.2	51.6	50.9	60.6	42.3	48.9	44.2	47.9
Residential	1.2	21.3	..	8.1	6.0	31.6	14.1	6.8	8.0
Total	105.9	106.9	105.3	123.0	129.1	132.2	92.7	89.1	109.7
2011-12									
Admitted patient	64.9	39.9	56.2	65.7	55.5	54.9	26.9	37.2	55.2
Ambulatory	42.2	46.9	53.9	51.8	58.2	40.1	51.6	49.2	48.1
Residential	1.3	21.0	..	8.8	9.9	26.8	14.1	6.6	8.2
Total	108.4	107.8	110.0	126.2	123.6	121.8	92.6	93.0	111.5
2012-13									
Admitted patient	61.5	38.6	55.2	65.7	51.2	47.4	29.4	48.6	53.3
Ambulatory	39.4	46.6	51.7	49.8	59.8	42.9	51.9	49.5	46.7
Residential	0.9	20.3	..	7.6	9.8	28.5	17.6	7.8	7.9
Total	101.8	105.6	106.9	123.1	120.7	118.8	98.8	105.8	107.8

Table 12A.28 FTE direct care staff employed in specialised mental health services, by service setting (per 100 000 people) (a), (b), (c)

	<i>NSW (d), (e)</i>	<i>Vic</i>	<i>Qld (f), (g)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
2013-14									
Admitted patient	62.2	40.2	51.4	64.2	52.6	52.9	34.1	48.0	53.3
Ambulatory	39.3	47.2	47.6	49.1	61.0	41.4	49.4	55.3	45.9
Residential	0.8	20.7	..	8.4	10.4	29.0	18.2	8.5	8.1
Total	102.2	108.1	99.0	121.7	123.9	123.3	101.7	111.8	107.3

(a) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of staffing estimates.

(b) Total FTE figures in this table can differ from those in table 12A.27. A new organisational overhead setting for reporting FTE was implemented from the 2012–13 collection period, which may result in decreased FTE in the other service setting categories for some jurisdictions. Time series analyses should be approached with caution. Care and consumer worker FTE has been included in service setting reporting since the 2012–13 collection period. These categories are not included in staff type FTE data. Comparisons between these tables should be made with caution.

(c) Due to the ongoing validation of NMDS, data could differ from previous reports.

(d) Caution is required when interpreting NSW data. Seven residential mental health services in 2006–07 were reclassified as non-acute older person specialised hospital services in 2007–08, reflecting a change in function of those units.

(e) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.

(f) The apparent absence of community residential services in Queensland reflects Queensland's preference to describe such facilities as 'extended inpatient care'.

(g) Queensland implemented a new method to calculate FTE from the 2009–10 data. The new method is associated with the reduction in reported FTE so caution should be exercised when conducting time series analysis.

.. Not applicable.

Source: AIHW (unpublished) MHE NMDS.

Table 12A.29 Targeted Community Care (Mental Health) program participants (number)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>2012-13</i>									
Personal Helpers and Mentors	4 325	3 319	3 173	1 410	1 781	458	266	334	15 066
Family Mental Health Support Services	9 693	12 190	11 701	13 323	8 989	3 337	5 484	3 689	68 406
Mental Health Respite: Carer Support	11 760	8 915	4 179	3 003	4 254	1 163	333	714	34 321
<i>2013-14</i>									
Personal Helpers and Mentors	5 256	4 118	3 919	1 880	2 069	570	326	401	18 539
Family Mental Health Support Services	18 151	11 283	21 211	22 249	10 731	4 206	4 774	6 059	98 664
Mental Health Respite: Carer Support	7 675	12 894	8 364	4 123	4 581	1 262	629	939	40 467

Source: DSS (various issues) *Targeted Community Care (Mental Health) Program – Summary Data (various years)*, (available at www.dss.gov.au/our-responsibilities/mental-health/publications-articles).

Table 12A.30 Admitted patient mental health-related separations without specialised psychiatric care, by principal diagnosis in ICD-10-AM groupings, 2012-13

ICD-10		Public acute and public psychiatric hospital	Per cent
		no.	%
F00–F03	Dementia	4 077	4.8
F04–F09	Other organic mental disorders	9 465	11.1
F10	Mental and behavioural disorders due to use of alcohol	16 623	19.4
F11–F19	Mental and behavioural disorders due to other psychoactive substance use	5 981	7.0
F20	Schizophrenia	4 157	4.9
F21, F24, F28, F29	Schizotypal and other delusional disorders	1 034	1.2
F22	Persistent delusional disorders	532	0.6
F23	Acute and transient psychotic disorders	814	1.0
F25	Schizoaffective disorders	1 628	1.9
F30	Manic episode	329	0.4
F31	Bipolar affective disorders	2 750	3.2
F32	Depressive episode	10 170	11.9
F33	Recurrent depressive disorders	1 634	1.9
F34	Persistent mood (affective) disorders	119	0.1
F38–F39	Other and unspecified mood (affective) disorders	68	0.1
F40	Phobic anxiety disorders	36	–
F41	Other anxiety disorders	4 639	5.4
F42	Obsessive-compulsive disorders	51	0.1
F43	Reaction to severe stress and adjustment disorders	5 659	6.6
F44	Dissociative (conversion) disorders	1 301	1.5
F45, F48	Somatoform and other neurotic disorders	398	0.5
F50	Eating disorders	1 373	1.6
F51–F59	Other behavioural syndromes associated with physiological disturbances and physical factors	737	0.9
F60	Specific personality disorders	1 182	1.4
F61–F69	Disorders of adult personality and behaviour	100	0.1
F70–F79	Mental retardation	146	0.2
F80–F89	Disorders of psychological development	550	0.6
F90	Hyperkinetic disorders	23	–

Table 12A.30 **Admitted patient mental health-related separations without specialised psychiatric care, by principal diagnosis in ICD-10-AM groupings, 2012-13**

<i>ICD-10</i>		<i>Public acute and public psychiatric hospital</i>	<i>Per cent</i>
F91	Conduct disorders	502	0.6
F92–F98	Other and unspecified disorders with onset in childhood or adolescence	349	0.4
F99	Mental disorder not otherwise specified	202	0.2
G30	Alzheimer's disease	2 493	2.9
	Other factors related to mental and behavioural disorders and substance use (a)	354	0.4
	Other specified mental health-related principal diagnosis (b)	6 081	7.1
	Total	85 557	100.0

(a) Includes ICD-10-AM codes Z00.4, Z03.2, Z04.6, Z09.3, Z13.3, Z54.3, Z63.1, Z63.8, Z63.9, Z65.8, Z65.9, Z71.4, Z71.5 and Z76.0.

(b) Includes separations for which the principal diagnosis was any other mental health-related principal diagnosis as listed in the online technical information.

– Nil or rounded to zero.

Source: AIHW (2014) *Mental Health Services in Australia*, (available at <http://mhsa.aihw.gov.au/home/>).

Table 12A.31 Ambulatory-equivalent mental health-related separations without specialised psychiatric care, by principal diagnosis, 2012-13 (a)

<i>ICD-10-AM code</i>		<i>Number of separations</i>	<i>Proportion of total separations</i>
F00–F09	Organic, including symptomatic, mental disorders	432	2.4
F10–F19	Mental and behavioural disorders due to psychoactive substance use	7 859	42.9
F20–F29	Schizophrenia, schizotypal and delusional disorders	1 104	6.0
F30–F39	Mood (affective) disorders	1 734	9.5
F40–F48	Neurotic, stress-related and somatoform disorders	4 174	22.8
F50–F59	Behavioural syndromes associated with physiological disturbances and physical factors	116	0.6
F60–F69	Disorders of adult personality and behaviour	559	3.1
F70–F79	Mental retardation	25	0.1
F80–F89	Disorders of psychological development	55	0.3
F90–F98	Behavioural and emotional disorders with onset usually occurring in childhood and adolescence	213	1.2
F99	Mental disorder not otherwise specified	56	0.3
G30	Alzheimer's disease	63	0.3
	Other factors related to mental and behavioural disorders and substance use (b)	313	1.7
	Other specified mental health-related principal diagnosis (c)	1 617	8.8
Total		18 320	100.0

(a) Includes separations for both public acute and public psychiatric hospitals.

(b) Includes ICD-10-AM codes Z00.4, Z03.2, Z04.6, Z09.3, Z13.3, Z54.3, Z61.9, Z63.1, Z63.8, Z63.9, Z65.8, Z65.9, Z71.4, Z71.5 and Z76.0.

(c) Includes separations for which the principal diagnosis was any other mental health-related principal diagnosis.

Source: AIHW (2015) *Mental Health Services in Australia*, (available at <http://mhsa.aihw.gov.au/home/>).

TABLE 12A.32

Table 12A.32 **Mental health-related emergency department occasions of service in public hospitals, by episode end status, 2012–13 (a)**

	<i>NSW</i>	<i>Vic (b)</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Episode end status (number)									
Admitted to this hospital (c)	22 582	12 553	13 196	7 549	5 743	1 819	1 198	2 823	67 463
Non-admitted patient emergency department service episode completed (d)	35 107	26 782	34 104	12 419	10 022	2 667	2 374	3 638	127 113
Referred to another hospital for admission	1 152	1 951	953	892	849	43	115	6	5 961
Did not wait to be attended by a health care professional (b)	1 068	..	337	121	38	11	14	20	1 609
Left at own risk (e)	2 294	1 228	1 589	387	269	44	50	250	6 111
Principal diagnosis (f)									
Organic, including symptomatic, mental disorders	3 976	2 376	3 664	1 911	1 158	395	312	134	13 926
Mental and behavioural disorders due to psychoactive substance use	14 097	10 975	13 121	5 810	4 340	1 039	891	4 215	54 488
Schizophrenia, schizotypal and delusional disorders	7 081	5 922	5 805	1 604	2 270	680	689	756	24 807
Mood (affective) disorders	9 167	8 122	5 923	2 757	1 606	1 021	752	407	29 755
Neurotic, stress-related and somatoform disorders	17 766	10 975	13 147	7 767	5 968	876	825	1 076	58 400
Behavioural syndromes associated with physiological disturbances and physical factors	483	437	595	96	174	32	23	9	1 849
Disorders of adult personality and behaviour	2 314	1 695	1 207	357	378	234	141	18	6 344

TABLE 12A.32

Table 12A.32 **Mental health-related emergency department occasions of service in public hospitals, by episode end status, 2012–13 (a)**

	NSW	Vic (b)	Qld	WA	SA	Tas	ACT	NT	Aust
Mental retardation	9	56	..	1	3	–	–	–	69
Disorders of psychological development	48	–	..	7	45	2	9	–	111
Behavioural and emotional disorders with onset usually occurring in childhood and adolescence	1 393	960	923	612	810	73	26	122	4 919
Unspecified mental disorder	8 693	997	5 798	450	197	253	83	–	16 471
Total (g)	65 027	42 515	50 183	21 372	16 949	4 605	3 751	6 737	211 139
Rates (per 10 000 population) (h)	88.4	74.8	108.8	86.2	102.0	89.9	99.2	282.4	92.1

(a) Emergency department occasions of service included are those that had a principal diagnosis that fell within the Mental and behavioural disorders chapter (Chapter 5) of ICD-10-AM (codes F00–F99) or the equivalent ICD-9-CM codes.

(b) Victoria does not record a diagnosis for occasions of service with an episode end status of *Did not wait to be attended by a health care professional*.

(c) Includes admissions to beds or units within the emergency department.

(d) Patient departed without being admitted or referred to another hospital.

(e) Patient left at own risk after being attended by a health care professional but before the non-admitted patient emergency department occasion of service was completed.

(f) NSW and SA used a combination of ICD-9-CM and ICD-10-AM.

(g) The number of occasions of service may not sum to the total due to missing or not reported data. Also included in the total are 6 occasions of service with an episode end status of *Died in emergency department as a non-admitted patient* and 1 occasion of service with an episode end status of *Dead on arrival, not treated in emergency department*.

(h) Crude rate is based on the preliminary Australian estimated resident population as at 31 December 2012 and is expressed per 10 000 population as detailed in the online technical information.

.. Not applicable. – Nil or rounded to zero.

Source: AIHW (2015) *Mental Health Services in Australia*, (available at <http://mhsa.aihw.gov.au/services/emergency-departments/>).

TABLE 12A.33

Table 12A.33 **New clients as a proportion of total clients under the care of State or Territory specialised public mental health services (a), (b)**

	<i>Unit</i>	<i>NSW (c), (d)</i>	<i>Vic (e)</i>	<i>Qld (f)</i>	<i>WA (g)</i>	<i>SA (h)</i>	<i>Tas (i)</i>	<i>ACT</i>	<i>NT (j)</i>	<i>Aust</i>
<i>2009-10</i>										
New clients	no.	46 323	22 717	32 301	17 059	13 206	1 369	3 210	2 652	138 837
Total clients	no.	116 276	61 129	73 903	41 928	31 186	6 209	7 661	5 552	343 844
Proportion of total clients who are new	%	39.8	37.2	43.7	40.7	42.3	22.0	41.9	47.8	40.4
<i>2010-11</i>										
New clients	no.	48 506	22 695	34 440	18 749	13 302	1 691	3 305	2 815	145 503
Total clients	no.	119 380	61 686	78 129	44 980	31 689	7 845	8 093	5 834	357 636
Proportion of total clients who are new	%	40.6	36.8	44.1	41.7	42.0	21.6	40.8	48.3	40.7
<i>2011-12</i>										
New clients	no.	49 590	na	36 655	19 772	14 557	1 204	3 470	3 263	128 511
Total clients	no.	123 341	na	82 179	47 296	34 092	6 390	8 412	6 607	308 317
Proportion of total clients who are new	%	40.2	na	44.6	41.8	42.7	18.8	41.3	49.4	41.7
<i>2012-13</i>										
New clients	no.	51 651	na	39 807	21 448	15 693	3 880	3 751	3 453	139 683
Total clients	no.	129 183	na	86 469	50 267	35 992	6 678	9 046	7 212	324 847
Proportion of total clients who are new	%	40.0	na	46.0	42.7	43.6	58.1	41.5	47.9	43.0
<i>2013-14</i>										
New clients	no.	54 171	23 880	40 429	22 790	15 903	3 817	3 949	3 402	168 341
Total clients	no.	133 513	64 978	89 119	53 166	37 168	8 440	9 825	7 381	403 590
Proportion of total clients who are new	%	40.6	36.8	45.4	42.9	42.8	45.2	40.2	46.1	41.7

(a) Clients in receipt of services include all people who received one or more community service contacts or had one or more days of inpatient or residential care in the data period.

(b) A new client is defined as a consumer who has not been seen in the five years preceding the first contact with a State or Territory specialised public mental health service in the data period.

Table 12A.33 New clients as a proportion of total clients under the care of State or Territory specialised public mental health services (a), (b)

	<i>Unit</i>	<i>NSW (c), (d)</i>	<i>Vic (e)</i>	<i>Qld (f)</i>	<i>WA (g)</i>	<i>SA (h)</i>	<i>Tas (i)</i>	<i>ACT</i>	<i>NT (j)</i>	<i>Aust</i>
(c)	NSW has implemented a Statewide Unique Patient Identifier (SUPI) for mental health care. The identification of prior contacts for mental health (MH) clients is dependent upon the SUPI, both in coverage (all clients having a SUPI) and in the resolution of possible duplicates. There are differences in the completeness of coverage between the Local Health Districts/Networks and over time. The average SUPI coverage at a State level for 2012-13 is 99.9 per cent. The numbers provided are a distinct count of individuals using the SUPI (majority) and a count of individuals at the facility level for a small percentage of clients without a SUPI in the reporting period (which may include some duplicates of those who attended multiple facilities).									
(d)	For NSW, residential clients are not included because their data are manually collected without SUPI assigned, thus making the unique counts of the residential clients together with the inpatient and ambulatory clients not possible. The client base of the NSW MH residential is very small which will have minimal effect on the final result (for example, total residential MH clients in 2010-11 is 185 with 59 potential new clients, 243 total residential MH clients with 130 potential new clients in 2011-12 and 237 total residential MH clients with 131 potential new clients in 2012-13).									
(e)	Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. The total only includes those jurisdictions that have provided data.									
(f)	For Qld, a linkage program is utilised to link between admitted and community activity and patients.									
(g)	For WA, the matching of mental health community contacts to inpatient episodes is done from 2012-13 between two separate data systems and requires the use of record linkage to be able to identify the same person in both systems. There are delays associated in the use of record linkage and these delays can result in not getting a match between a community contact and a separation when there should be one. The number of unique consumers (both total and new) could be over estimated as a result. Data before 2012-13 are based on data submitted for the National Minimum Data Set (NMDS) and have not been revised.									
(h)	For SA, the new client (numerator) count is not unique: it is an aggregation of three separate databases with no linkage between them. Similarly, the total client (denominator) count is not unique: it is an aggregation of three separate databases with no linkage between them. However, the impact on the result should be minimal due to populations being relatively stable within the three respective catchments.									
(i)	For Tasmania, information for years before 2012-13 were extracted from three different data sources and linked together with a Statistical Linkage Key (SLK) for each individual present in the extracts for the reporting period. While every attempt has been made to reduce any duplication of identified clients, using an SLK will lead to some duplication and can wrongly identify clients as new clients. Industrial action in Tasmania has limited the available data quality and quantity of data for 2011-12 and 2012-13. Tasmania has been progressively implementing a state-wide patient identification system. Data for 2012-13 is considered to be the first collection period with this system fully implemented. It is likely that an improved patient identification system will increase the percentage of post-discharge community care reported by Tasmania. Therefore, Tasmanian data is not comparable across years.									
(j)	For the NT, for 2009-10, the count of all clients will not be exactly the same as provided in other reported collections due to non-availability of 'snapshot' or archived annual data sets.									
	na Not available.									

Table 12A.33 **New clients as a proportion of total clients under the care of State or Territory specialised public mental health services (a), (b)**

<i>Unit</i>	<i>NSW (c), (d)</i>	<i>Vic (e)</i>	<i>Qld (f)</i>	<i>WA (g)</i>	<i>SA (h)</i>	<i>Tas (i)</i>	<i>ACT</i>	<i>NT (j)</i>	<i>Aust</i>
-------------	---------------------	----------------	----------------	---------------	---------------	----------------	------------	---------------	-------------

Source: AIHW (unpublished) derived from data provided by State and Territory governments.

TABLE 12A.34

Table 12A.34 **New clients as a proportion of total clients under the care of State or Territory specialised public mental health services, by selected characteristics, 2013-14 (a), (b), (c), (d)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (d)</i>
<i>Age group</i>										
Less than 15 years	%	52.5	52.4	53.7	48.7	44.6	59.6	58.6	51.9	51.4
15–24 years	%	43.9	40.6	46.2	45.4	50.4	53.8	40.6	51.5	45.0
25–34 years	%	39.1	35.3	43.2	39.9	44.1	43.0	39.5	44.8	40.2
35–44 years	%	35.1	31.0	39.9	34.9	37.0	39.7	35.9	37.5	35.7
45–54 years	%	34.5	30.3	41.2	43.5	35.4	35.7	34.1	43.3	36.3
55–64 years	%	36.3	31.8	44.4	36.2	37.8	36.2	34.8	45.2	37.3
65–74 years	%	40.4	35.5	48.6	43.4	42.3	36.7	44.5	52.2	41.6
75 years or over	%	49.3	46.8	58.2	52.4	50.9	48.1	52.6	70.0	50.7
<i>Gender</i>										
Male	%	39.0	35.3	43.5	42.3	41.5	45.4	38.3	43.2	40.3
Female	%	42.1	38.1	47.2	43.1	43.8	45.0	41.8	49.4	43.1
<i>SEIFA quintiles (e)</i>										
Quintile 1 (most disadvantaged)	%	39.8	34.6	38.9	42.2	41.9	44.1	72.9	42.1	39.5
Quintile 2	%	39.8	35.7	43.3	40.3	39.7	45.0	28.7	32.0	39.9
Quintile 3	%	40.2	38.0	45.5	41.7	41.3	48.8	49.4	29.5	41.4
Quintile 4	%	39.8	37.3	46.9	44.4	42.5	45.9	36.3	41.9	41.7
Quintile 5 (least disadvantaged)	%	41.9	37.4	46.8	46.0	43.6	50.4	40.7	45.0	42.8
<i>Indigenous status</i>										
Aboriginal and Torres Strait Islander	%	32.2	32.2	36.8	38.6	33.5	45.2	38.0	38.5	35.4
Non-Indigenous	%	40.1	36.4	46.1	42.1	40.9	44.6	39.9	51.2	41.4
<i>Remoteness (e)</i>										
Major cities	%	40.1	36.7	43.1	42.0	39.4	58.9	38.5	83.3	40.3
Inner regional	%	40.8	36.9	43.6	44.8	43.9	43.8	51.0	80.6	41.2

Table 12A.34 **New clients as a proportion of total clients under the care of State or Territory specialised public mental health services, by selected characteristics, 2013-14 (a), (b), (c), (d)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (d)</i>
Outer regional	%	39.0	34.7	43.7	40.6	46.2	48.2	70.4	47.0	42.2
Remote	%	43.8	35.1	40.2	45.8	47.5	50.8	na	41.4	43.9
Very remote	%	36.3	50.0	39.0	52.4	36.8	45.2	100.0	43.5	43.4

- (a) Clients in receipt of services include all people who received one or more community service contacts or had one or more days of inpatient or residential care in the data period.
- (b) A new client is defined as a consumer who has not been seen in the five years preceding the first contact with a State or Territory specialised public mental health service in the data period.
- (c) See table 12A.33 and the DQI for specific footnotes regarding each State or Territory.
- (d) Analysis exclude data for clients for who the relevant demographic variable is missing (for example, Indigenous status).
- (e) Remoteness and socioeconomic status have been allocated using the client's usual residence, not the location of the service provider. State/territory is reported for the state/territory of the service provider.

na Not available.

Source: AIHW (unpublished) derived from data provided by State and Territory governments.

TABLE 12A.35

Table 12A.35 **New clients as a proportion of total clients of MBS subsidised mental health services (a), (b), (c)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
2012-13										
New clients	no.	207 845	170 671	133 286	54 394	43 915	13 091	9 367	3 930	636 500
Total clients	no.	573 106	489 338	345 281	141 813	124 312	34 532	23 892	8 050	1 740 323
Proportion of total clients who are new	%	36.3	34.9	38.6	38.4	35.3	37.9	39.2	48.8	36.6
2013-14										
New clients	no.	218 380	180 387	143 630	62 348	45 668	13 829	10 153	4 211	678 606
Total clients	no.	621 649	535 423	385 785	160 493	133 634	37 678	26 393	8 659	1 909 713
Proportion of total clients who are new	%	35.1	33.7	37.2	38.8	34.2	36.7	38.5	48.6	35.5
2014-15										
New clients	no.	228 067	187 335	151 461	68 405	47 944	14 838	10 710	4 654	713 414
Total clients	no.	669 237	576 409	421 761	178 387	144 243	40 761	28 707	9 502	2 069 005
Proportion of total clients who are new	%	34.1	32.5	35.9	38.3	33.2	36.4	37.3	49.0	34.5

(a) Data are calculated based on date of processing of specified MBS mental health items.

(b) State/Territory is allocated based on the postcode recorded for the person at the first service event within each reference period year.

(c) A new client is defined as a patient who has not previously used a MBS mental health item in the five years preceding the first use of a MBS mental health item in the reference period.

Source: Australian Government Department of Health (unpublished).

TABLE 12A.36

Table 12A.36 **New clients as a proportion of total clients under the care of MBS subsidised mental health services, by selected characteristics, 2014-15 (a), (b), (c)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Age group</i>										
Less than 15 years	%	51.7	49.0	52.4	57.3	50.7	54.6	56.5	66.1	51.6
15–24 years	%	38.2	36.0	40.2	42.2	37.4	40.4	38.2	54.7	38.5
25–34 years	%	33.8	31.9	34.5	36.4	32.3	32.9	34.8	47.7	33.6
35–44 years	%	29.7	28.1	31.1	33.7	28.9	30.8	34.1	44.2	30.0
45–54 years	%	28.3	27.5	30.4	33.5	28.1	30.9	32.9	46.3	29.1
55–64 years	%	29.0	28.2	31.2	34.2	28.0	33.7	36.0	43.9	29.8
65–74 years	%	33.5	32.0	35.4	38.1	30.8	37.2	39.1	46.5	33.7
75 years or over	%	40.8	39.2	44.0	46.8	39.5	52.6	43.8	60.7	41.4
<i>Gender</i>										
Male	%	37.3	36.1	39.8	42.2	36.8	40.2	41.0	54.3	38.1
Female	%	31.9	30.1	33.3	35.8	30.8	34.0	35.1	45.4	32.1
<i>Indigenous status</i>										
Aboriginal and Torres Strait Islander	%	34.1	31.0	39.5	44.9	37.3	35.2	38.1	55.2	36.7
Non-Indigenous	%	34.1	32.5	35.8	38.2	33.1	36.5	37.3	48.0	34.4
<i>Remoteness</i>										
Major cities	%	33.6	31.9	34.3	37.5	32.3	..	37.3	..	33.6
Inner regional	%	34.7	33.8	37.4	39.3	34.7	35.1	36.6	..	35.3
Outer regional	%	38.8	38.9	41.3	41.6	37.7	39.3	..	48.8	40.3
Remote	%	45.0	41.7	48.4	52.0	43.4	42.5	..	48.2	47.8
Very remote	%	46.0	..	51.2	54.0	43.2	34.7	..	50.8	50.3

(a) Clients in receipt of services include all people who received one or more community service contacts or had one or more days of inpatient or residential care in the data period.

Table 12A.36 New clients as a proportion of total clients under the care of MBS subsidised mental health services, by selected characteristics, 2014-15 (a), (b), (c)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
--	-------------	------------	------------	------------	-----------	-----------	------------	------------	-----------	-------------

(b) A new client is defined as a consumer who has not been seen in the five years preceding the first contact with a State or Territory specialised public mental health service in the data period.

(c) Remoteness data have been allocated using the client's usual residence, not the location of the service provider. State/territory is reported for the state/territory of the service provider.

.. Not applicable.

Source: Australian Government Department of Health (unpublished).

TABLE 12A.37

Table 12A.37 Proportion of people receiving clinical mental health services by service type and Indigenous status

		<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Age standardised proportion (%) (a)											no.
<i>2007-08</i>											
Public (b), (c)											
Aboriginal and Torres Strait Islander		3.6	2.6	3.5	3.1	4.3	1.3	4.8	2.9	3.3	19 187
Non-Indigenous		1.2	1.1	1.8	1.6	1.5	2.0	1.6	1.9	1.4	276 005
Private (d)											
Aboriginal and Torres Strait Islander		na	na	na	na	na	na	na	..	na	na
Non-Indigenous		na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)											
Aboriginal and Torres Strait Islander		np	np	np	np	np	np	np	np	np	np
Non-Indigenous		np	np	np	np	np	np	np	np	np	np
<i>2008-09</i>											
Public (b), (c)											
Aboriginal and Torres Strait Islander		3.9	2.6	3.4	3.4	4.8	1.1	4.8	3.0	3.4	20 616
Non-Indigenous		1.2	1.1	1.6	1.7	1.6	1.3	1.7	1.9	1.4	277 321
Private (d)											
Aboriginal and Torres Strait Islander		na	na	na	na	na	na	na	..	na	na
Non-Indigenous		na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)											
Aboriginal and Torres Strait Islander		5.9	7.6	3.9	2.4	4.7	5.6	6.7	1.0	4.4	24 603

TABLE 12A.37

Table 12A.37 Proportion of people receiving clinical mental health services by service type and Indigenous status

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Non-Indigenous	6.0	6.5	5.5	4.9	5.7	5.0	4.6	2.7	5.8	1 200 337
<i>2009-10</i>										
Public (b), (c)										
Aboriginal and Torres Strait Islander	4.0	2.6	3.5	3.8	4.8	3.0	4.8	3.7	3.7	22 930
Non-Indigenous	1.2	1.1	1.6	1.7	1.6	1.4	1.7	2.0	1.4	282 620
Private (d)										
Aboriginal and Torres Strait Islander	na	na	na	na	na	na	na	..	na	na
Non-Indigenous	na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)										
Aboriginal and Torres Strait Islander	6.6	8.3	4.2	2.7	5.1	6.2	7.2	1.3	4.8	28 303
Non-Indigenous	6.5	7.1	6.1	5.3	6.4	5.6	5.0	3.2	6.4	1 337 882
<i>2010-11</i>										
Public (b), (c)										
Aboriginal and Torres Strait Islander	3.9	2.5	3.9	4.3	4.9	1.6	5.3	3.7	3.8	24 250
Non-Indigenous	1.2	1.1	1.7	1.8	1.6	1.6	1.8	2.0	1.4	291 381
Private (d)										
Aboriginal and Torres Strait Islander	na	na	na	na	na	na	na	..	na	na
Non-Indigenous	na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)										
Aboriginal and Torres Strait Islander	8.2	9.2	5.2	3.6	6.5	7.6	9.7	1.5	6.0	36 044

TABLE 12A.37

Table 12A.37 Proportion of people receiving clinical mental health services by service type and Indigenous status

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Non-Indigenous	7.1	7.8	6.8	5.7	7.0	6.3	5.5	3.4	7.0	1 486 676
2011-12										
Public (b), (c)										
Aboriginal and Torres Strait Islander	4.5	na	4.2	4.9	5.7	1.0	6.4	3.9	4.3	26 133
Non-Indigenous	1.2	na	1.7	1.8	1.7	0.8	1.9	2.3	1.5	240 556
Private (d)										
Aboriginal and Torres Strait Islander	na	na	na	na	na	na	na	..	na	na
Non-Indigenous	na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)										
Aboriginal and Torres Strait Islander	9.7	11.4	6.4	3.7	7.5	7.6	10.7	1.4	7.0	43 634
Non-Indigenous	7.3	8.1	7.1	5.7	7.3	6.4	5.6	3.7	7.2	1 559 298
2012-13 (f)										
Public (b), (c)										
Aboriginal and Torres Strait Islander	4.9	na	4.5	5.3	5.9	1.4	6.3	4.1	4.7	29 424
Non-Indigenous	1.5	na	1.8	1.9	1.8	1.2	2.2	2.4	1.7	269 241
Private (d)										
Aboriginal and Torres Strait Islander	na	na	na	na	na	na	na	..	na	na
Non-Indigenous	na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)										
Aboriginal and Torres Strait Islander	10.7	12.0	7.1	4.0	8.2	8.8	11.4	1.4	7.7	49 787

TABLE 12A.37

Table 12A.37 Proportion of people receiving clinical mental health services by service type and Indigenous status

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Non-Indigenous	7.9	8.7	7.7	5.9	7.6	7.0	6.2	4.1	7.7	1 690 537
2013-14 (f)										
Public (b), (c)										
Aboriginal and Torres Strait Islander	5.4	2.9	4.5	5.6	6.0	2.0	8.1	4.2	4.8	32 870
Non-Indigenous	1.5	1.1	1.8	1.9	1.9	1.6	2.4	2.3	1.6	345 219
Private (d)										
Aboriginal and Torres Strait Islander	na	na	na	na	na	na	na	..	na	na
Non-Indigenous	na	na	na	na	na	na	na	..	na	na
MBS and DVA (e)										
Aboriginal and Torres Strait Islander	12.0	13.5	8.1	5.1	9.2	9.9	11.8	1.6	8.7	57 620
Non-Indigenous	8.4	9.4	8.4	6.4	8.2	7.7	6.7	4.3	8.3	1 852 094

- (a) Rates are age-standardised to the Australian population as at 30 June 2001.
- (b) Excludes people for whom Indigenous status was missing or not reported. The Indigenous status rates should be interpreted with caution due to the varying and, in some instances, unknown quality of Indigenous identification across jurisdictions.
- (c) Caution should be taken when making inter-jurisdictional comparisons for public data. South Australia submitted data that were not based on unique patient identifier or data matching approaches. This was also the case for data submitted by Tasmania prior to 2012-13. Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. Industrial action during 2011-12 and 2012-13 in Tasmania has limited the available data quality and quantity of the community mental health care data; which represents a large proportion of the overall figures. Australian totals for 2011-12 and 2012-13 only include available data and should therefore be interpreted with caution. Australian totals for 2011-12 and 2012-13 should not be compared to previous years.
- (d) Indigenous information is not collected for private psychiatric hospitals.

Table 12A.37 **Proportion of people receiving clinical mental health services by service type and Indigenous status**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
(e)	DVA data not available by Indigenous status. MBS data are not published for 2007-08. Medicare data presented by Indigenous status have been adjusted for under-identification in the Department of Human Services (DHS) Voluntary Indigenous Identifier (VII) database. Indigenous rates are therefore modelled and should be interpreted with caution. These statistics are not derived from the total Australian Indigenous population, but from those Aboriginal and Torres Strait Islander people who have voluntarily identified as Indigenous to DHS. The statistics have been adjusted to reflect demographic characteristics of the overall Indigenous population, but this adjustment may not address all the differences in the service use patterns of the enrolled population relative to the total Indigenous population. The level of VII enrolment (61 per cent nationally as at August 2012) varies across age-sex-remoteness-State/Territory sub-groups and over time which means that the extent of adjustment required varies across jurisdictions and over time. Indigenous rates should also be interpreted with caution due to small population numbers in some jurisdictions.								
(f)	A change in scope for 'public data' implemented for the 2012-13 collection period means that comparison of rates for years up to and including 2011-12 with years 2012-13 and beyond should not be made. See data quality information for additional information.								
	na Not available. .. Not applicable. np Not published.								

Source: AIHW (unpublished), derived from data provided by State and Territory governments; Private Mental Health Alliance; Centralised Data Management Service data; Australian Government Department of Health and DVA; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period).

TABLE 12A.38

Table 12A.38 Proportion of people receiving clinical mental health services by service type and remoteness area (a)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
	Age standardised proportion (%) (b)									no.
<i>2007-08</i>										
Public (c), (d)										
Major cities	1.2	0.9	1.5	1.3	1.6	..	1.8	..	1.2	173 288
Inner regional	2.6	1.7	2.5	3.9	1.7	np	np	..	2.2	85 003
Outer regional	3.5	2.2	2.2	2.2	2.6	np	..	2.0	2.3	43 447
Remote	4.4	4.3	1.9	0.9	2.0	np	..	2.2	1.9	5 744
Very remote	13.0	..	3.9	4.8	2.1	np	..	2.2	3.6	6 297
Private (c), (e)										
Major cities	0.1	0.1	0.1	0.1	np	..	np	..	0.1	19 261
Inner regional	0.1	–	0.1	0.1	np	np	np	..	0.1	2 973
Outer regional	–	–	–	–	np	np	–	579
Remote	–	–	–	–	np	np	–	69
Very remote	–	..	–	–	np	np	–	30
MBS and DVA (c)										
Major cities	5.3	5.8	5.1	4.6	5.2	..	4.0	..	5.3	764 089
Inner regional	5.1	5.3	4.6	3.7	4.5	4.8	4.6	..	4.9	192 134
Outer regional	3.7	3.7	3.1	3.6	3.2	3.4	..	2.4	3.3	62 986
Remote	2.5	4.7	1.9	1.4	2.5	2.1	..	0.9	1.8	5 668
Very remote	2.6	..	1.2	0.7	2.7	5.5	..	1.2	1.3	2 070
<i>2008-09</i>										
Public (c), (d)										
Major cities	1.2	0.9	1.4	1.3	1.9	..	1.9	..	1.2	180 087
Inner regional	2.7	1.5	2.4	4.0	2.0	np	np	..	2.2	85 135
Outer regional	4.0	2.1	2.2	2.3	2.6	np	..	2.0	2.4	44 963
Remote	5.8	1.5	1.6	0.9	2.5	np	..	2.5	2.0	6 193

TABLE 12A.38

Table 12A.38 Proportion of people receiving clinical mental health services by service type and remoteness area (a)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Very remote	16.2	..	3.1	5.1	2.3	np	..	2.2	3.7	6 544
Private (c), (e)										
Major cities	0.1	0.1	0.2	0.1	np	..	np	..	0.1	20 251
Inner regional	0.1	–	0.1	0.1	np	np	np	..	0.1	3 205
Outer regional	–	–	–	–	np	np	–	645
Remote	0.1	–	–	–	np	np	–	98
Very remote	–	..	–	–	np	np	–	30
MBS and DVA (c)										
Major cities	6.2	6.7	6.1	5.3	6.3	..	4.8	..	6.2	916 074
Inner regional	6.2	6.6	5.7	4.7	5.5	5.6	5.7	..	6.0	239 453
Outer regional	4.7	4.5	4.0	4.4	4.1	4.2	..	3.0	4.2	80 394
Remote	3.0	6.1	2.5	1.9	3.4	2.7	..	1.3	2.4	7 460
Very remote	4.3	..	1.6	0.8	2.4	6.3	..	1.6	1.5	2 557
2009-10										
Public (c), (d)										
Major cities	1.4	0.9	1.6	1.3	1.8	..	2.0	..	1.3	198 917
Inner regional	2.2	1.6	1.8	4.3	2.1	1.4	np	..	2.0	81 749
Outer regional	2.6	2.1	1.8	2.3	2.5	1.2	..	2.0	2.1	39 579
Remote	3.8	1.0	1.5	1.0	2.6	–	..	2.8	1.9	5 798
Very remote	5.5	..	2.4	5.8	2.1	0.7	..	2.6	3.5	6 416
Private (c), (e)										
Major cities	0.1	0.1	0.2	0.2	np	..	np	..	0.1	21 149
Inner regional	0.1	0.1	0.1	0.1	np	np	np	..	0.1	3 416
Outer regional	–	–	–	–	np	np	–	674
Remote	0.1	0.1	–	–	np	np	–	105
Very remote	–	..	–	–	np	np	–	31

TABLE 12A.38

Table 12A.38 Proportion of people receiving clinical mental health services by service type and remoteness area (a)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
MBS and DVA (c)										
Major cities	6.6	7.3	6.7	5.7	6.9	..	5.2	..	6.7	1 011 181
Inner regional	6.8	7.4	6.3	5.2	6.5	6.3	6.4	..	6.7	270 641
Outer regional	5.2	5.4	4.7	4.9	4.6	4.8	..	3.4	4.8	93 109
Remote	3.2	6.3	2.8	2.3	4.4	2.8	..	1.6	2.7	8 759
Very remote	4.9	..	1.7	1.0	2.3	4.9	..	2.0	1.7	2 963
<i>2010-11</i>										
Public (c), (d)										
Major cities	1.4	0.9	1.7	1.8	1.8	..	2.1	..	1.4	214 072
Inner regional	2.2	1.6	1.8	1.6	2.1	1.9	np	..	1.9	76 427
Outer regional	2.5	2.0	1.9	2.5	2.4	1.6	..	2.0	2.1	40 932
Remote	3.5	1.2	1.9	3.0	2.6	0.6	..	2.7	2.6	8 115
Very remote	5.1	..	2.9	2.0	2.5	0.7	..	3.1	2.5	4 820
Private (c), (e)										
Major cities	0.1	0.2	0.2	0.2	np	..	np	..	0.1	22 910
Inner regional	0.1	0.1	0.1	0.1	np	np	np	..	0.1	3 950
Outer regional	–	–	–	–	np	np	–	858
Remote	0.1	0.1	–	0.1	np	np	–	115
Very remote	–	..	–	–	np	np	–	45
MBS and DVA (c)										
Major cities	7.3	7.9	7.4	6.1	7.6	..	5.6	..	7.3	1 124 293
Inner regional	7.6	8.1	6.9	5.9	7.1	6.9	6.4	..	7.4	301 981
Outer regional	5.7	6.3	5.3	5.5	5.1	5.5	..	3.6	5.4	104 578
Remote	3.2	5.8	3.6	2.6	4.0	3.4	..	1.8	3.0	9 668
Very remote	4.4	..	1.9	1.2	2.3	7.4	..	2.0	1.8	3 314

2011-12

TABLE 12A.38

Table 12A.38 Proportion of people receiving clinical mental health services by service type and remoteness area (a)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Public (c), (d)										
Major cities	1.4	na	1.6	1.8	1.9	..	2.1	..	1.6	181 124
Inner regional	2.2	na	2.1	2.0	2.1	1.7	np	..	2.1	59 145
Outer regional	2.6	na	2.2	2.8	2.8	1.4	..	2.3	2.3	39 567
Remote	3.3	na	2.1	2.8	2.7	1.5	..	2.8	2.6	7 915
Very remote	5.2	..	3.0	2.9	2.4	0.9	..	3.3	3.0	6 164
Private (c), (e)										
Major cities	0.1	0.2	0.2	0.2	np	..	np	..	0.2	25 188
Inner regional	0.1	0.1	0.1	0.1	np	np	np	..	0.1	4 112
Outer regional	–	–	0.1	0.1	np	np	0.1	1 104
Remote	0.1	0.2	–	0.1	np	np	–	122
Very remote	–	..	0.1	–	np	np	–	75
MBS and DVA (c)										
Major cities	7.5	8.3	7.9	5.9	7.9	..	5.8	..	7.6	1 191 781
Inner regional	7.9	8.2	7.0	5.8	7.4	6.8	6.4	..	7.6	297 015
Outer regional	6.1	6.3	5.5	4.7	5.4	6.0	..	3.0	5.4	106 181
Remote	3.7	5.7	3.4	2.5	4.1	4.3	..	1.8	3.0	9 465
Very remote	3.1	..	1.9	1.3	2.2	6.5	..	0.6	1.5	2 892
2012-13 (f)										
Public (c), (d)										
Major cities	1.5	na	1.6	1.8	2.0	..	2.2	..	1.7	193 900
Inner regional	2.4	na	2.0	2.0	2.3	1.4	np	..	2.1	61 745
Outer regional	2.8	na	2.2	2.8	3.0	1.2	..	2.4	2.4	41 122
Remote	3.5	na	2.5	3.4	2.8	1.6	..	3.5	3.1	9 415
Very remote	5.4	..	2.6	2.5	4.2	0.7	..	2.9	2.8	5 814
Private (c), (e)										

TABLE 12A.38

Table 12A.38 Proportion of people receiving clinical mental health services by service type and remoteness area (a)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Major cities	0.2	0.2	0.2	0.2	np	..	np	..	0.2	26 968
Inner regional	0.1	0.1	0.1	0.1	np	np	np	..	0.1	4 610
Outer regional	–	0.1	0.1	0.1	np	np	0.1	1 133
Remote	0.1	0.1	–	0.1	np	np	–	137
Very remote	–	..	–	–	np	np	–	49
MBS and DVA (c)										
Major cities	8.0	8.8	8.5	6.1	8.2	..	6.3	..	8.1	1 289 439
Inner regional	8.7	9.0	7.5	6.2	8.0	7.4	5.1	..	8.2	325 303
Outer regional	6.6	6.7	6.0	4.7	5.8	6.6	..	3.4	5.9	116 157
Remote	4.0	6.8	3.5	2.5	4.4	4.8	..	1.9	3.2	10 102
Very remote	2.9	..	1.8	1.3	2.7	5.8	..	0.6	1.5	3 003
2013-14 (f)										
Public (c), (d)										
Major cities	1.6	1.0	1.6	1.9	2.0	..	2.0	..	1.5	243 790
Inner regional	2.4	1.6	2.1	2.2	2.3	1.7	np	..	2.1	81 695
Outer regional	2.9	2.1	2.3	3.1	3.1	1.5	..	2.3	2.5	47 936
Remote	3.4	1.9	2.4	3.6	2.9	1.4	..	3.5	3.1	9 652
Very remote	6.9	..	2.9	2.7	4.8	1.1	..	3.1	3.1	6 415
Private (c), (e)										
Major cities	0.2	0.2	0.2	0.2	np	..	np	..	0.2	27 209
Inner regional	0.1	0.1	0.1	0.1	np	np	np	..	0.1	4 906
Outer regional	0.1	0.1	0.1	0.1	np	np	0.1	1 220
Remote	0.1	0.1	–	0.1	np	np	–	144
Very remote	–	..	–	–	np	np	–	61
MBS and DVA (c)										
Major cities	8.4	9.4	9.2	6.6	8.7	..	6.8	..	8.6	1 404 185

Table 12A.38 **Proportion of people receiving clinical mental health services by service type and remoteness area (a)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Inner regional	9.6	9.9	8.4	6.9	8.6	8.1	np	..	9.1	362 877
Outer regional	7.5	7.5	6.9	5.4	6.3	7.3	..	3.5	6.6	131 005
Remote	4.8	7.4	3.9	3.0	4.9	5.2	..	2.0	3.6	11 456
Very remote	3.6	..	2.2	1.6	3.3	5.3	..	0.6	1.7	3 512

- (a) Not all remoteness areas are represented in each State or Territory. Where a state/territory does not have a particular remoteness category a rate cannot be calculated. Excludes contacts for which demographic information was missing and/or not reported.
- (b) Rates are age-standardised to the Australian population as at 30 June 2001.
- (c) For 2007-08 and 2008-09, disaggregation by remoteness area is based on a person's usual residence, the location of the service provider or a combination of both. For these years, the public data should be interpreted with caution as the methodology used to allocate remoteness area varied across jurisdictions. For 2009-10 data onwards, disaggregation by remoteness area is based on a person's usual residence, not the location of the service provider. State/territory is the state/territory of the service provider.
- (d) Caution should be taken when making inter-jurisdictional comparisons for public data. South Australian data are not based on unique patient identifier or data matching approaches. This was also the case for data submitted by Tasmania prior to 2012-13. Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. Industrial action during 2011-12 and 2012-13 in Tasmania has limited the available data quality and quantity of the community mental health care data; which represents a large proportion of the overall figures. Australian totals for 2011-12 and 2012-13 only include available data and should therefore be interpreted with caution. Australian totals for 2011-12 and 2012-13 should not be compared to previous years.
- (e) Private psychiatric hospital figures are not published for SA, Tasmania, and the ACT due to confidentiality reasons, but are included in the Australia figures.
- (f) Public 2012-13 data are considered a break in series due to a change of scope between 2011-12 and 2012-13. Historical remoteness area data was not re-supplied due to this change in scope. Therefore, changes in public data from 2012-13 onwards with years prior to 2012-13 should not be made.

na Not available. **..** Not applicable. **-** Nil or rounded to zero. **np** Not published.

Source: AIHW (unpublished) derived from data provided by State and Territory governments; State and territory (unpublished) specialised mental health services data; Private Mental Health Alliance (unpublished) Centralised Data Management Service data; Department of Health (unpublished) and DVA (unpublished), MBS Statistics; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period).

TABLE 12A.39

Table 12A.39 Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
	Age standardised proportion (c)									no.
<i>2007-08</i>										
Public (d), (e)										
Quintile 1 (most disadvantaged)	1.8	1.5	1.9	2.0	2.9	2.0	np	1.5	1.9	76 635
Quintile 2	1.9	1.4	2.6	1.4	1.2	2.9	4.3	6.1	1.8	74 505
Quintile 3	1.5	1.2	2.0	2.1	1.0	1.3	3.7	3.8	1.6	67 420
Quintile 4	1.4	0.9	1.7	2.0	1.3	0.9	2.3	0.6	1.4	55 904
Quintile 5 (least disadvantaged)	1.2	0.7	1.2	1.4	2.0	..	1.5	2.5	1.2	48 530
Private (d), (f)										
Quintile 1 (most disadvantaged)	–	0.1	–	0.1	np	np	np	np	0.1	2 556
Quintile 2	–	–	0.1	–	np	np	np	np	0.1	2 351
Quintile 3	0.1	–	0.1	0.1	np	np	np	np	0.1	3 572
Quintile 4	0.1	0.1	0.2	0.1	np	np	np	np	0.1	5 383
Quintile 5 (least disadvantaged)	0.2	0.2	0.2	0.2	np	..	np	np	0.2	9 074
MBS and DVA										
Quintile 1 (most disadvantaged)	4.4	4.9	4.3	2.3	4.5	3.8	3.7	0.7	4.3	176 364
Quintile 2	5.3	5.2	4.1	3.9	4.8	3.9	4.2	2.0	4.9	200 248
Quintile 3	5.2	5.4	4.6	3.9	4.5	4.2	3.9	1.6	4.8	202 268
Quintile 4	5.3	5.5	4.9	3.9	5.0	6.1	4.0	1.7	5.0	206 586
Quintile 5 (least disadvantaged)	5.4	6.3	4.9	4.8	5.4	..	3.9	1.4	5.4	231 002
<i>2008-09</i>										
Public (d), (e)										
Quintile 1 (most disadvantaged)	1.9	1.5	1.7	2.2	2.7	np	np	1.6	1.8	72 356
Quintile 2	2.0	1.4	2.7	1.5	1.3	np	4.6	6.2	1.9	77 089
Quintile 3	1.5	1.2	2.3	2.1	1.3	np	3.8	4.0	1.7	71 113
Quintile 4	1.4	0.8	1.3	2.0	0.8	np	2.4	0.6	1.2	51 399

TABLE 12A.39

Table 12A.39 Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Quintile 5 (least disadvantaged)	1.2	0.7	1.0	1.4	3.5	..	1.6	2.4	1.2	50 798
Private (d), (f)										
Quintile 1 (most disadvantaged)	–	0.1	–	0.1	np	np	np	np	–	2 036
Quintile 2	–	–	0.1	0.1	np	np	np	np	0.1	2 578
Quintile 3	0.1	0.1	0.1	0.1	np	np	np	np	0.1	3 888
Quintile 4	0.1	0.2	0.2	0.1	np	np	np	np	0.1	6 212
Quintile 5 (least disadvantaged)	0.2	0.2	0.2	0.2	np	..	np	np	0.2	9 553
MBS and DVA										
Quintile 1 (most disadvantaged)	5.3	5.8	5.4	2.7	5.6	4.6	4.6	0.9	5.2	218 084
Quintile 2	6.3	6.2	5.1	4.7	5.9	4.7	4.8	2.5	5.9	244 695
Quintile 3	6.1	6.5	5.7	4.8	5.7	4.9	4.8	2.2	5.8	247 895
Quintile 4	6.1	6.5	5.8	4.5	5.7	6.7	4.9	2.0	5.9	250 106
Quintile 5 (least disadvantaged)	6.3	7.2	5.6	5.5	6.3	..	4.6	1.8	6.2	270 901
2009-10										
Public (d), (e)										
Quintile 1 (most disadvantaged)	1.9	1.5	2.6	2.2	2.7	1.0	np	2.6	2.0	85 633
Quintile 2	1.9	1.4	1.8	1.5	2.1	4.2	4.8	2.4	1.8	75 384
Quintile 3	1.5	1.2	1.7	2.2	1.7	1.3	3.8	3.3	1.6	69 386
Quintile 4	1.4	0.8	1.4	2.1	1.2	1.0	2.5	1.6	1.3	56 689
Quintile 5 (least disadvantaged)	1.1	0.7	1.0	1.4	1.0	..	1.7	1.7	1.0	45 247
Private (d), (f)										
Quintile 1 (most disadvantaged)	–	0.1	0.0	0.1	np	np	np	np	–	1 939
Quintile 2	0.1	0.1	0.1	0.1	np	np	np	np	0.1	2 864
Quintile 3	0.1	0.1	0.1	0.1	np	np	np	np	0.1	4 121
Quintile 4	0.1	0.1	0.2	0.2	np	np	np	np	0.1	5 993
Quintile 5 (least disadvantaged)	0.2	0.2	0.2	0.3	np	..	np	np	0.2	10 565

TABLE 12A.39

Table 12A.39 Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
MBS and DVA										
Quintile 1 (most disadvantaged)	5.9	6.5	6.0	3.1	6.4	5.2	5.2	1.1	5.8	246 684
Quintile 2	6.8	6.9	5.7	5.1	6.6	5.1	5.3	3.0	6.5	274 627
Quintile 3	6.6	7.2	6.4	5.2	6.2	5.6	5.2	2.5	6.4	277 661
Quintile 4	6.5	7.1	6.4	4.9	6.2	7.5	5.3	2.3	6.4	278 258
Quintile 5 (least disadvantaged)	6.7	7.6	6.1	5.8	6.9	..	5.0	2.1	6.6	293 715
2010-11										
Public (d), (e)										
Quintile 1 (most disadvantaged)	1.9	1.5	2.9	3.5	2.7	2.0	np	2.9	2.2	93 565
Quintile 2	1.9	1.4	1.9	2.2	2.1	1.4	4.4	2.5	1.9	79 324
Quintile 3	1.6	1.2	1.7	1.9	1.7	1.2	3.7	3.0	1.6	69 526
Quintile 4	1.4	0.8	1.3	1.6	1.3	1.7	2.6	1.7	1.3	55 664
Quintile 5 (least disadvantaged)	1.1	0.7	1.0	1.4	1.0	..	1.7	1.8	1.0	45 973
Private (d), (f)										
Quintile 1 (most disadvantaged)	–	0.1	–	0.1	np	np	np	np	–	2 179.0
Quintile 2	0.1	0.1	0.1	0.1	np	np	np	np	0.1	3 217.0
Quintile 3	0.1	0.1	0.1	0.1	np	np	np	np	0.1	4 752.0
Quintile 4	0.1	0.1	0.2	0.2	np	np	np	np	0.1	6 743.0
Quintile 5 (least disadvantaged)	0.2	0.3	0.2	0.3	np	..	np	np	0.2	10 987.0
MBS and DVA										
Quintile 1 (most disadvantaged)	6.5	7.2	6.6	3.7	7.0	5.9	5.8	1.2	6.5	277 164
Quintile 2	7.6	7.6	6.5	5.5	7.3	5.6	5.9	3.4	7.2	309 010
Quintile 3	7.1	7.9	7.2	5.5	6.7	6.3	5.4	2.8	7.0	307 839
Quintile 4	7.2	7.7	7.2	5.3	6.9	8.0	5.7	2.4	7.0	312 702
Quintile 5 (least disadvantaged)	7.2	8.1	6.6	6.2	7.6	..	5.5	2.2	7.1	319 001

TABLE 12A.39

Table 12A.39 Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
<i>2011-12</i>										
Public (d), (e), (g)										
Quintile 1 (most disadvantaged)	1.9	na	2.8	4.1	3.1	1.7	np	2.9	2.3	81 894
Quintile 2	1.9	na	2.0	2.4	2.1	1.3	3.5	2.9	2.0	64 732
Quintile 3	1.5	na	1.8	2.1	1.7	1.4	3.3	2.9	1.7	58 780
Quintile 4	1.5	na	1.4	1.7	1.5	1.3	2.9	2.6	1.6	46 849
Quintile 5 (least disadvantaged)	1.1	na	1.1	1.5	1.1	..	1.8	2.2	1.3	41 555
Private (d)										
Quintile 1 (most disadvantaged)	na	na	na	na	na	na	na	..	0.1	2 394
Quintile 2	na	na	na	na	na	na	na	..	0.1	3 524
Quintile 3	na	na	na	na	na	na	na	..	0.1	5 461
Quintile 4	na	na	na	na	na	na	na	..	0.2	7 354
Quintile 5 (least disadvantaged)	na	na	na	na	na	..	na	..	0.3	11 868
MBS and DVA										
Quintile 1 (most disadvantaged)	na	na	na	na	na	na	na	na	7.1	306 636
Quintile 2	na	na	na	na	na	na	na	na	7.2	311 718
Quintile 3	na	na	na	na	na	na	na	na	7.3	322 463
Quintile 4	na	na	na	na	na	na	na	na	7.4	328 411
Quintile 5 (least disadvantaged)	na	na	na	na	na	..	na	na	7.2	320 535
<i>2012-13</i>										
Public (d), (e), (g), (h)										
Quintile 1 (most disadvantaged)	2.2	na	3.3	3.2	3.2	1.7	np	3.2	2.6	90 816
Quintile 2	2.1	na	2.1	2.8	2.4	1.2	7.1	2.4	2.2	74 825
Quintile 3	1.7	na	1.5	2.1	1.9	1.0	4.4	3.9	1.7	56 760
Quintile 4	1.6	na	1.3	1.7	1.4	1.0	2.7	2.6	1.5	47 696
Quintile 5 (least disadvantaged)	1.2	na	1.0	1.3	1.0	0.9	1.8	2.1	1.2	41 679

TABLE 12A.39

Table 12A.39 Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Private (d)										
Quintile 1 (most disadvantaged)	na	na	na	na	na	na	na	..	0.1	2 411
Quintile 2	na	na	na	na	na	na	na	..	0.1	3 998
Quintile 3	na	na	na	na	na	na	na	..	0.1	5 506
Quintile 4	na	na	na	na	na	na	na	..	0.2	7 835
Quintile 5 (least disadvantaged)	na	na	na	na	na	na	na	..	0.3	13 147
MBS and DVA										
Quintile 1 (most disadvantaged)	na	na	na	na	na	na	na	na	7.7	336 345
Quintile 2	na	na	na	na	na	na	na	na	7.7	340 118
Quintile 3	na	na	na	na	na	na	na	na	7.9	351 587
Quintile 4	na	na	na	na	na	na	na	na	7.8	355 720
Quintile 5 (least disadvantaged)	na	na	na	na	na	na	na	na	7.6	343 452
2013-14										
Public (d), (e), (g), (h)										
Quintile 1 (most disadvantaged)	2.2	1.6	3.4	3.3	3.3	2.1	5.8	2.9	2.5	109 926
Quintile 2	2.1	1.5	2.1	2.9	2.5	1.6	5.0	1.7	2.1	91 951
Quintile 3	1.8	1.2	1.5	2.3	1.9	1.2	3.6	2.2	1.6	73 186
Quintile 4	1.6	0.9	1.4	1.8	1.4	1.3	2.8	2.0	1.4	62 214
Quintile 5 (least disadvantaged)	1.2	0.7	1.0	1.4	1.0	1.1	1.5	1.5	1.1	50 479
Private (d)										
Quintile 1 (most disadvantaged)	na	na	na	na	na	na	na	..	0.1	2 637
Quintile 2	na	na	na	na	na	na	na	..	0.1	4 295
Quintile 3	na	na	na	na	na	na	na	..	0.1	6 007
Quintile 4	na	na	na	na	na	na	na	..	0.2	7 855
Quintile 5 (least disadvantaged)	na	na	na	na	na	na	na	..	0.3	12 746
MBS and DVA										

Table 12A.39 Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
Quintile 1 (most disadvantaged)	na	na	na	na	na	na	na	na	na	8.4 375 999
Quintile 2	na	na	na	na	na	na	na	na	na	8.4 380 544
Quintile 3	na	na	na	na	na	na	na	na	na	8.5 390 726
Quintile 4	na	na	na	na	na	na	na	na	na	8.4 389 627
Quintile 5 (least disadvantaged)	na	na	na	na	na	na	na	na	na	8.1 375 362

- (a) Socio-Economic Indexes for Areas (SEIFA) quintiles are based on the ABS Index of Relative Socio-economic Disadvantage, with quintile 1 being the most disadvantaged and quintile 5 being the least disadvantaged. SEIFA quintiles represent approximately 20 per cent of the national population, but do not necessarily represent 20 per cent of the population in each State or Territory. Excludes people for whom demographic information was missing and/or not reported.
- (b) Disaggregation by SEIFA is based on a person's usual residence, not the location of the service provider.
- (c) Rates are age-standardised to the Australian population as at 30 June 2001.
- (d) For 2007-08 and 2008-09, disaggregation by SEIFA is based on a person's usual residence, the location of the service provider or a combination of both. For these years, the public data should be interpreted with caution as the methodology used to allocate SEIFA varied across jurisdictions. From 2009-10 onwards, disaggregation by SEIFA is based on a person's usual residence, not the location of the service provider, except for public data for the NT, for which the majority of the data was based on the location of the service. Due to system-related issues impacting data quality, Tasmania was unable to provide data by SEIFA for 2008-09.
- (e) SA submitted data that were not based on unique patient identifier or data matching approaches. This was also the case for data submitted by Tasmania before 2012-13. Therefore caution should be taken when making inter-jurisdictional comparisons.
- (f) Private psychiatric hospital figures are not published for SA, Tasmania, and the ACT due to confidentiality reasons but are included in the Australia figures.
- (g) Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. Industrial action during 2011-12 and 2012-13 in Tasmania has limited the available data quality and quantity of the community mental health care data; which represents a large proportion of the overall figures. Australian totals for 2011-12 and 2012-13 only include available data and should therefore be interpreted with caution.
- (h) A change in scope for public data implemented for the 2012-13 collection period means that comparison of rates for years up to and including 2011-12 with years 2012-13 and beyond should not be made. See the quality statement for additional information.

na Not available. .. Not applicable. – Nil or rounded to zero. np Not published.

Table 12A.39 **Proportion of people receiving clinical mental health services by service type and SEIFA (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Source:</i>	AIHW (unpublished) derived from data provided by State and Territory governments. State and Territory (unpublished) community mental health care data; Private Mental Health Alliance (unpublished) Centralised Data Management Service data; Department of Health (unpublished) and DVA (unpublished), MBS Statistics; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period).								

Table 12A.40 **People receiving clinical State and Territory governments' specialised public mental health services, by age group and gender, 2013-14 (a)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>	
	%									<i>no.</i>
Age groups										
Less than 15 years	1.0	0.5	1.3	1.2	1.8	1.2	0.8	1.4	1.0	43 886
15–24 years	2.9	1.5	3.2	3.1	3.3	2.2	4.4	4.8	2.7	83 682
25–34 years	2.3	1.3	2.5	2.6	2.9	2.0	3.2	4.0	2.2	74 062
35–44 years	2.3	1.5	2.4	2.6	2.9	2.0	3.2	3.7	2.2	71 606
45–54 years	1.9	1.2	1.8	2.0	2.3	1.5	2.5	2.6	1.7	53 688
55–64 years	1.3	0.9	1.2	1.5	1.5	1.1	1.7	1.8	1.2	31 804
65+ years	1.3	1.1	1.2	2.0	1.3	1.8	2.2	1.3	1.3	44 539
Gender (b)										
Male	1.9	1.1	2.0	2.0	2.3	1.6	2.4	2.9	1.8	200 345
Female	1.8	1.1	1.9	2.3	2.3	1.7	2.8	2.8	1.8	202 572
Total (b)	1.8	1.1	2.0	2.1	2.3	1.7	2.6	2.8	1.8	403 311

(a) Caution should be taken when making inter-jurisdictional comparisons for public data. SA data are not based on unique patient identifier or data matching approaches.

(b) Rates are age-standardised to the Australian population as at 30 June 2001.

Source: AIHW (unpublished) derived from data provided by State and Territory governments; State and Territory governments (unpublished) specialised mental health services data; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period).

Table 12A.41 Proportion of people receiving clinical mental health services, by service type and SEIFA IRSD deciles (age-standardised rate) (a), (b), (c)

	<i>Public (d)</i>	<i>Private</i>	<i>MBS and DVA</i>
<i>2007-08</i>			
Decile 1	1.9	0.1	4.1
Decile 2	1.9	–	4.5
Decile 3	1.9	0.1	4.8
Decile 4	1.8	0.1	5.0
Decile 5	1.6	0.1	4.8
Decile 6	1.6	0.1	4.9
Decile 7	1.3	0.1	4.9
Decile 8	1.5	0.1	5.1
Decile 9	1.2	0.2	5.5
Decile 10	1.1	0.2	5.3
<i>2008-09</i>			
Decile 1	1.7	–	5.0
Decile 2	1.8	–	5.5
Decile 3	1.8	0.1	5.8
Decile 4	1.9	0.1	5.9
Decile 5	1.7	0.1	5.8
Decile 6	1.6	0.1	5.9
Decile 7	1.2	0.1	5.7
Decile 8	1.2	0.1	6.0
Decile 9	1.2	0.2	6.4
Decile 10	1.2	0.2	6.0
<i>2009-10</i>			
Decile 1	2.1	–	5.6
Decile 2	2.0	–	6.1
Decile 3	1.8	0.1	6.4
Decile 4	1.7	0.1	6.5
Decile 5	1.6	0.1	6.3
Decile 6	1.6	0.1	6.4
Decile 7	1.4	0.1	6.2
Decile 8	1.2	0.1	6.5
Decile 9	1.1	0.2	6.8
Decile 10	1.0	0.2	6.5
<i>2010-11</i>			
Decile 1	2.2	–	6.3
Decile 2	2.2	0.1	6.7
Decile 3	1.8	0.1	7.2
Decile 4	1.9	0.1	7.2
Decile 5	1.7	0.1	6.9

TABLE 12A.41

Table 12A.41 **Proportion of people receiving clinical mental health services, by service type and SEIFA IRSD deciles (age-standardised rate) (a), (b), (c)**

	<i>Public (d)</i>	<i>Private</i>	<i>MBS and DVA</i>
Decile 6	1.4	0.1	7.0
Decile 7	1.3	0.1	6.9
Decile 8	1.2	0.2	7.1
Decile 9	1.1	0.2	7.3
Decile 10	1.0	0.3	6.9
<i>2011-12</i>			
Decile 1	1.9	–	6.9
Decile 2	1.9	0.1	7.2
Decile 3	1.5	0.1	7.2
Decile 4	1.5	0.1	6.9
Decile 5	1.3	0.1	7.1
Decile 6	1.3	0.1	7.2
Decile 7	1.1	0.1	7.1
Decile 8	1.0	0.2	7.3
Decile 9	1.0	0.2	7.0
Decile 10	0.8	0.3	7.0
<i>2012-13 (e)</i>			
Decile 1	2.2	–	7.6
Decile 2	2.0	0.1	7.9
Decile 3	1.8	0.1	7.9
Decile 4	1.6	0.1	7.5
Decile 5	1.3	0.1	7.8
Decile 6	1.3	0.1	7.9
Decile 7	1.2	0.2	7.8
Decile 8	0.9	0.2	7.9
Decile 9	1.0	0.3	7.6
Decile 10	0.9	0.3	7.6
<i>2013-14 (e)</i>			
Decile 1	2.5	–	8.1
Decile 2	2.4	0.1	8.7
Decile 3	2.2	0.1	8.7
Decile 4	2.0	0.1	8.2
Decile 5	1.6	0.1	8.4
Decile 6	1.6	0.1	8.6
Decile 7	1.5	0.1	8.3
Decile 8	1.3	0.2	8.5
Decile 9	1.2	0.2	8.1
Decile 10	1.0	0.3	8.1

Table 12A.41 Proportion of people receiving clinical mental health services, by service type and SEIFA IRSD deciles (age-standardised rate) (a), (b), (c)

	<i>Public (d)</i>	<i>Private</i>	<i>MBS and DVA</i>
(a)	SEIFA deciles are based on the ABS Index of Relative Socio-economic Disadvantage (IRSD), with decile 1 being the most disadvantaged and decile 10 being the least disadvantaged. SEIFA deciles represent approximately 10 per cent of the national population, but do not necessarily represent 10 per cent of the population in each State or Territory. Excludes people for whom information was missing and/or not reported.		
(b)	Disaggregation by SEIFA is based on a person's usual residence, not the location of the service provider, except for public data for the NT, for which around 89 per cent of the data was based on the location of the service.		
(c)	Rates are age-standardised to the Australian population as at 30 June 2001.		
(d)	For 2007-08 and 2008-09, disaggregation by SEIFA is based on a person's usual residence, the location of the service provider or a combination of both. For these years, the public data should be interpreted with caution as the methodology used to allocate SEIFA varied across jurisdictions. From 2009-10 onwards, disaggregation by SEIFA is based on a person's usual residence, not the location of the service provider. Due to system-related issues impacting data quality, Tasmania was unable to provide data by SEIFA for 2008-09. Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. Industrial action during 2011-12 and 2012-13 in Tasmania has limited the available data quality and quantity of the community mental health care data; which represents a large proportion of the overall figures. Australian totals for 2011-12 and 2012-13 only include available data and should therefore be interpreted with caution.		
(e)	A change in scope for public data implemented for the 2012-13 collection period means that comparison of rates for years up to and including 2011-12 with years 2012-13 and beyond should not be made. See the quality statement for additional information. – Nil or rounded to zero.		
<i>Source:</i>	AIHW (unpublished) derived from data provided by State and Territory governments; State and Territory governments (unpublished) specialised mental health services data; Private Mental Health Alliance (unpublished) Centralised Data Management Service data; Department of Health (unpublished) and DVA (unpublished), MBS Statistics; ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period).		

TABLE 12A.42

Table 12A.42 Proportion of people receiving clinical mental health services by service type (a)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>2007-08</i>										
Public (b)										
Number	no.	109 250	59 352	76 227	37 862	28 400	9 499	6 832	4 559	331 981
Rate	%	1.6	1.1	1.9	1.8	1.9	2.0	2.0	2.1	1.6
Private (c)										
Number	no.	7 256	6 170	4 791	2 183	np	np	np	..	23 044
Rate	%	0.1	0.1	0.1	0.1	np	np	np	..	0.1
MBS and DVA										
Number: Total MBS and DVA (d)	no.	349 679	287 210	189 005	87 638	75 116	20 527	14 163	3 981	1027 330
Rate: Total MBS and DVA (d)	%	5.1	5.5	4.6	4.1	4.8	4.3	4.0	1.8	4.9
Rate: Psychiatrist (e)	%	1.4	1.5	1.3	1.1	1.6	1.0	1.1	0.4	1.4
Rate: Clinical psychologist (f)	%	0.6	0.6	0.4	1.0	0.7	0.9	0.6	0.1	0.6
Rate: GP (g)	%	3.7	4.0	3.2	3.0	3.2	3.2	2.8	1.4	3.5
Rate: Other allied health (h)	%	1.4	1.8	1.4	0.6	0.9	1.1	1.2	0.4	1.3
<i>2008-09</i>										
Public (b)										
Number	no.	112 751	60 034	74 168	39 886	30 777	9 362	7 371	4 930	339 279
Rate	%	1.6	1.1	1.8	1.8	2.0	1.9	2.1	2.2	1.6
Private (c)										
Number	no.	7 575	6 308	5 270	2 629	np	np	np	..	24 348
Rate	%	0.1	0.1	0.1	0.1	np	np	np	..	0.1
MBS and DVA										
Number: Total MBS and DVA (d)	no.	419 027	346 064	235 222	107 077	91 841	24 501	17 119	5 104	1 247 142
Rate: Total MBS and DVA (d)	%	6.0	6.6	5.6	4.9	5.8	5.1	4.8	2.3	5.9
Rate: Psychiatrist (e)	%	1.4	1.5	1.3	1.1	1.6	1.0	1.1	0.4	1.4
Rate: Clinical psychologist (f)	%	0.8	0.8	0.6	1.2	1.1	1.2	0.7	0.2	0.8

TABLE 12A.42

Table 12A.42 Proportion of people receiving clinical mental health services by service type (a)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Rate: GP (g)	%	4.6	4.9	4.2	3.7	4.2	3.9	3.4	1.9	4.4
Rate: Other allied health (h)	%	1.7	2.3	1.8	0.8	1.1	1.3	1.5	0.5	1.7
<i>2009-10</i>										
Public (b)										
Number	no.	116 276	61 130	73 903	41 928	31 208	6 209	7 670	5 450	343 774
Rate	%	1.7	1.1	1.7	1.9	2.0	1.3	2.1	2.3	1.6
Private (c)										
Number	no.	8 145	6 544	5 392	3 047	np	np	np	..	25 536
Rate	%	0.1	0.1	0.1	0.1	np	np	np	..	0.1
MBS										
Number: Total MBS and DVA (d)		460 708	385 085	265 357	119 533	103 225	27 741	18 871	6 146	1387 297
Rate: Total MBS and DVA (d)		6.6	7.2	6.1	5.3	6.5	5.7	5.2	2.7	6.4
Rate: Psychiatrist (e)		1.4	1.5	1.3	1.1	1.7	1.1	1.1	0.4	1.4
Rate: Clinical psychologist (f)		1.0	1.0	0.7	1.4	1.3	1.3	0.9	0.3	1.0
Rate: GP (g)		5.0	5.4	4.7	4.0	4.7	4.3	3.7	2.2	4.8
Rate: Other allied health (h)		2.0	2.6	2.1	1.0	1.2	1.5	1.7	0.7	2.0
<i>2010-11</i>										
Public (b)										
Number	no.	119 380	61 686	78 129	44 980	32 063	7 845	8 101	5 730	357 914
Rate	%	1.7	1.1	1.8	2.0	2.0	1.6	2.2	2.4	1.6
Private (c)										
Number	no.	8 354	7 692	5 673	3 250	np	np	np	..	27 924
Rate	%	0.1	0.1	0.1	0.1	np	np	np	..	0.1
MBS										
Number: Total MBS and DVA (d)		511 672	426 982	300 311	131 892	115 088	31 175	20 838	6 775	1 544 744
Rate: Total MBS and DVA (d)		7.2	7.8	6.8	5.7	7.1	6.4	5.6	2.9	7.0

TABLE 12A.42

Table 12A.42 Proportion of people receiving clinical mental health services by service type (a)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Rate: Psychiatrist (e)		1.4	1.5	1.3	1.1	1.6	1.1	1.2	0.4	1.4
Rate: Clinical psychologist (f)		1.1	1.1	0.9	1.4	1.7	1.4	1.2	0.3	1.1
Rate: GP (g)		5.6	6.1	5.4	4.4	5.4	5.0	4.2	2.4	5.5
Rate: Other allied health (h)		2.3	2.8	2.3	1.2	1.4	1.9	1.7	0.7	2.2
<i>2011-12</i>										
Public (b)										
Number	no.	123 341	na	82 179	47 296	34 090	6 390	8 427	6 437	308 160
Rate	%	1.7	na	1.9	2.0	2.2	1.3	2.3	2.7	1.9
Private (c)										
Number	no.	9 537	8 301	6 578	3 616	np	np	np	..	30 640
Rate	%	0.1	0.1	0.1	0.2	np	np	np	..	0.1
MBS										
Number: Total MBS and DVA (d)		536 353	453 347	320 397	134 105	119 613	32 031	21 926	7 307	1 625 098
Rate: Total MBS and DVA (d)		7.5	8.2	7.2	5.7	7.4	6.5	5.8	3.1	7.3
Rate: Psychiatrist (e)		1.4	1.5	1.4	1.1	1.6	1.1	1.1	0.4	1.4
Rate: Clinical psychologist (f)		1.2	1.3	1.0	1.5	1.9	1.5	1.5	0.4	1.3
Rate: GP (g)		5.8	6.4	5.6	4.3	5.6	5.1	4.4	2.6	5.7
Rate: Other allied health (h)		2.3	2.9	2.4	1.2	1.5	2.0	1.6	0.7	2.3
<i>2012-13</i>										
Public (b)										
Number	no.	129 183	na	86 469	50 267	35 992	6 678	9 058	7 051	324 698
Rate	%	1.8	na	1.9	2.1	2.3	1.3	2.4	2.9	1.9
Private (c)										
Number	no.	10 539	8 642	7 241	3 785	np	np	np	..	32 944
Rate	%	0.1	0.1	0.2	0.2	np	np	np	..	0.1
MBS										

TABLE 12A.42

Table 12A.42 **Proportion of people receiving clinical mental health services by service type (a)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Number: Total MBS and DVA (d)		580 047	492 618	353 147	143 637	126 345	34 848	24 275	8 097	1 763 028
Rate: Total MBS and DVA (d)		8.0	8.8	7.8	5.9	7.8	7.1	6.3	3.4	7.8
Rate: Psychiatrist (e)		1.5	1.6	1.5	1.1	1.7	1.2	1.0	0.3	1.5
Rate: Clinical psychologist (f)		1.4	1.6	1.2	1.5	2.0	1.9	1.7	0.4	1.5
Rate: GP (g)		6.3	6.9	6.1	4.5	5.9	5.6	4.9	2.9	6.1
Rate: Other allied health (h)		2.5	3.1	2.5	1.2	1.7	2.0	1.8	0.7	2.4
<i>2013-14</i>										
Public (b)										
Number	no.	133 513	64 978	89 119	53 166	37 168	8 440	9 825	7 102	403 311
Rate	%	1.8	1.1	2.0	2.1	2.3	1.7	2.6	2.8	1.8
Private (c)										
Number	no.	10 991	8 988	7 550	3 495	np	np	np	..	33 574
Rate	%	0.1	0.2	0.2	0.1	np	np	np	..	0.1
MBS										
Number: Total MBS and DVA (d)		628 834	538 678	394 456	162 357	135 747	37 995	26 809	8 720	1 933 630
Rate: Total MBS and DVA (d)		8.6	9.4	8.6	6.4	8.3	7.8	6.9	3.5	8.4
Rate: Psychiatrist (e)		1.6	1.7	1.7	1.1	1.7	1.3	0.9	0.3	1.5
Rate: Clinical psychologist (f)		1.6	1.8	1.5	1.6	2.3	2.3	1.8	0.4	1.7
Rate: GP (g)		6.9	7.5	6.8	5.0	6.4	6.1	5.5	3.1	6.7
Rate: Other allied health (h)		2.6	3.2	2.8	1.3	1.8	2.0	2.0	0.8	2.5

(a) Rates are age-standardised to the Australian population as at 30 June 2001.

Table 12A.42 **Proportion of people receiving clinical mental health services by service type (a)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
(b)	Caution should be taken when making inter-jurisdictional comparisons for public data. South Australian data are not based on unique patient identifier or data matching approaches. This was also the case for data submitted by Tasmania prior to 2012-13. Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. Tasmania data for 2007-08 and 2008-09 are limited to Community mental health care services only and are therefore not comparable with later years. Industrial action during 2011-12 and 2012-13 in Tasmania has limited the available data quality and quantity of the community mental health care data; which represents a large proportion of the overall figures. Australian public totals and rates for 2011-12 and 2012-13 should be interpreted with caution as they include only available data and are therefore not comparable to previous years.									
(c)	Private psychiatric hospital figures are not published for SA, Tasmania, and the ACT due to confidentiality reasons but are included in the Australia totals.									
(d)	MBS and DVA services are those provided under any of the Medicare/DVA-funded service types described at (e) to (h). People seen by more than one provider type are counted only once in the total.									
(e)	Consultant psychiatrist services are MBS items 134, 136, 138, 140, 142, 289, 291, 293, 296, 297, 299, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 319, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 342, 344, 346, 348, 350, 352, 353, 355, 356, 357, 358, 359, 361, 364, 366, 367, 369, 370, 855, 857, 858, 861, 864, 866, 14224 (as relevant across years).									
(f)	Clinical psychologist services are MBS items: 80000, 80005, 80010, 80015, 80020 and DVA items US01, US02, US03, US04, US05, US06, US07, US08, US50, US51, US99 (as relevant across years).									
(g)	GP services are MBS items 170, 171, 172, 2574, 2575, 2577, 2578, 2700, 2701, 2702, 2704, 2705, 2707, 2708, 2710, 2712, 2713, 2715, 2717, 2719, 2721, 2723, 2725, 2727, 20104 (as relevant across years).									
(h)	Other allied health services are MBS items 10956, 10968, 80100, 80105, 80110, 80115, 80120, 80125, 80130, 80135, 80140, 80145, 80150, 80155, 80160, 80165, 80170, 81325, 81355, 82000, 82015 and DVA items CL20, CL25, CL30, US11, US12, US13, US14, US15, US16, US17, US18, US21, US22, US23, US24, US25, US26, US27, US31, US32, US33, US34, US35, US36, US37, US52, US53, US96, US97, US98 (as relevant across years).									

na Not available. ... **Not applicable.** **np** Not published.

Source: AIHW (unpublished) derived from data provided by State and Territory governments, the Private Mental Health Alliance Centralised Data Management Service data; MBS Statistics Australian Government Department of Health (unpublished) and DVA (unpublished); ABS (unpublished) Estimated Residential Population, 30 June (prior to relevant period).

TABLE 12A.43

Table 12A.43 Services used for mental health problems, Australia, 2007 (per cent) (a), (b)

	With lifetime mental disorder		No lifetime mental disorder (e)	Total
	Symptoms in previous 12 months (c)	No symptoms in previous 12 months (d)		
GP	24.7 ± 2.4	6.2 ± 1.5	2.8 ± 0.9	8.1 ± 0.7
Psychiatrist	7.9 ± 2.7	1.4 ± 0.7	0.6 ± 0.3	2.3 ± 0.6
Psychologist	13.2 ± 2.1	1.8 ± 0.6	0.8 ± 0.3	3.5 ± 0.5
Other mental health professional	7.7 ± 1.6	1.5 ± 0.5	np	2.2 ± 0.4
Other health professional	6.6 ± 1.6	2.1 ± 1.0	1.0 ± 0.4	2.4 ± 0.5
Hospitalisation	2.6 ± 1.1	np	np	0.7 ± 0.3
Total who used health services	34.9 ± 3.1	9.2 ± 1.8	4.7 ± 1.1	11.9 ± 0.9
Total who did not use services for mental health	65.1 ± 3.1	90.8 ± 1.8	95.2 ± 1.1	88.1 ± 0.9

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) Estimates with RSEs greater than 25 per cent are considered unreliable. These estimates are not published.

(c) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(d) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(e) People who did not meet criteria for diagnosis of a lifetime mental disorder.

np Not published.

Source: ABS (unpublished) 2007 Survey of Mental Health and Wellbeing, Cat. no. 4326.0.

TABLE 12A.44

Table 12A.44 **Services used for mental health, by mental disorder status, 2007 (per cent) (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Total who used services for mental health in previous 12 months (c)									
Any 12-month mental disorder (d)	32.5 ± 6.4	37.0 ± 7.1	34.4 ± 7.0	35.8 ± 10.5	35.2 ± 9.1	np	np	np	34.9 ± 3.1
Lifetime mental disorder, with no 12-month symptoms (e)	7.6 ± 2.0	11.5 ± 4.0	9.0 ± 3.3	7.4 ± 3.6	np	np	np	np	9.2 ± 1.8
No lifetime mental disorder (f)	np	4.6 ± 1.8	5.6 ± 1.8	4.8 ± 2.3	np	np	np	np	4.7 ± 1.1
Total	10.9 ± 1.8	13.1 ± 2.2	12.1 ± 2.0	12.0 ± 2.5	11.0 ± 2.7	np	np	np	11.9 ± 0.9
Total who did not use services for mental health in previous 12 months									
Any 12-month mental disorder (d)	67.5 ± 6.4	63.0 ± 7.1	65.6 ± 7.0	64.2 ± 10.5	64.8 ± 9.1	65.5 ± 23.2	np	np	65.1 ± 3.1
Lifetime mental disorder, with no 12-month symptoms (e)	92.4 ± 2.0	88.5 ± 4.0	91.0 ± 3.3	92.6 ± 3.6	90.3 ± 5.7	87.8 ± 13.3	np	np	90.8 ± 1.8
No lifetime mental disorder (f)	95.4 ± 2.3	95.4 ± 1.8	94.4 ± 1.8	95.2 ± 2.3	96.1 ± 2.8	95.2 ± 7.1	np	np	95.2 ± 1.1
Total	89.1 ± 1.8	86.9 ± 2.2	87.9 ± 2.0	88.0 ± 2.5	88.6 ± 2.8	88.7 ± 6.9	81.6 ± 12.2	95.3 ± 6.2	88.1 ± 0.9

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) Estimates with RSEs greater than 25 per cent are considered unreliable. These estimates are not published.

(c) Includes hospitalisations.

(d) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(e) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(f) People who did not meet criteria for diagnosis of a lifetime mental disorder.

np Not published.

Source: ABS (unpublished) *2007 Survey of Mental Health and Wellbeing*, Cat. no. 4326.0.

TABLE 12A.45

Table 12A.45 **Young people who had contact with MBS subsidised primary mental health care services, by age group (a), (b), (c), (d), (e), (f), (g)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (h)</i>
<i>2010-11</i>										
<i>Number of children or young adults who have had contact</i>										
Pre-school (0-<5 years)	no.	2 868	2 527	1 326	645	760	114	76	25	8 341
Primary school (5-<12 years)	no.	21 250	18 890	12 749	5 106	5 037	1 180	803	225	65 242
Secondary school (12-<18 years)	no.	29 381	24 940	17 697	7 392	6 685	2 059	1 384	359	89 900
Youth/young adult (18-<25 years)	no.	49 576	42 417	30 564	14 091	11 699	3 693	2 538	758	155 338
All children and young people aged <25 years	no.	103 075	88 774	62 335	27 235	24 181	7 045	4 800	1 366	318 819
<i>Proportion of population who had contact with MBS-subsidised primary mental health services</i>										
Pre-school (0-<5 years)	%	0.6	0.7	0.4	0.4	0.8	0.4	0.3	0.1	0.6
Primary school (5-<12 years)	%	3.4	4.1	3.1	2.5	3.8	2.7	2.7	0.9	3.4
Secondary school (12-<18 years)	%	5.4	6.1	5.0	4.1	5.5	5.1	5.2	1.8	5.3
Youth/young adult (18-<25 years)	%	7.2	7.6	6.9	5.9	7.4	8.1	5.7	2.9	7.1
All children and young people aged <25 years	%	4.4	5.0	4.1	3.5	4.7	4.3	3.8	1.5	4.4
<i>2011-12</i>										
<i>Number of children or young adults who have had contact</i>										
Pre-school (0-<5 years)	no.	3 249	2 783	1 485	724	765	129	97	33	9 266
Primary school (5-<12 years)	no.	23 830	22 469	14 868	5 618	5 510	1 383	866	283	74 830
Secondary school (12-<18 years)	no.	32 882	28 198	19 921	8 282	7 399	2 311	1 515	439	100 950
Youth/young adult (18-<25 years)	no.	53 901	46 621	33 628	14 626	12 716	3 782	2 790	870	168 937
All children and young people aged <25 years	no.	113 861	100 072	69 902	29 250	26 391	7 605	5 268	1 624	353 982
<i>Proportion of population who had contact with MBS-subsidised primary mental health services</i>										
Pre-school (0-<5 years)	%	0.7	0.8	0.5	0.5	0.8	0.4	0.4	0.2	0.6
Primary school (5-<12 years)	%	3.8	4.8	3.6	2.7	4.1	3.1	2.8	1.2	3.8
Secondary school (12-<18 years)	%	6.1	6.9	5.6	4.5	6.1	5.7	5.8	2.3	5.9
Youth/young adult (18-<25 years)	%	7.8	8.4	7.5	6.0	8.1	8.3	6.3	3.4	7.7

TABLE 12A.45

Table 12A.45 Young people who had contact with MBS subsidised primary mental health care services, by age group (a), (b), (c), (d), (e), (f), (g)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (h)</i>
All children and young people aged <25 years	%	4.9	5.6	4.6	3.7	5.2	4.7	4.2	1.8	4.8
<i>2012-13</i>										
<i>Number of children or young adults who have had contact</i>										
Pre-school (0-<5 years)	no.	3 778	3 279	1 914	763	841	119	83	28	10 805
Primary school (5-<12 years)	no.	27 396	26 535	17 774	6 551	6 195	1 586	983	302	87 325
Secondary school (12-<18 years)	no.	38 242	33 217	24 143	10 293	8 528	2 703	1 985	427	119 542
Youth/young adult (18-<25 years)	no.	60 739	52 016	38 351	16 055	13 774	4 285	3 220	992	189 438
All children and young people aged <25 years	no.	130 155	115 047	82 181	33 662	29 337	8 693	6 272	1 749	407 110
<i>Proportion of population who had contact with MBS-subsidised primary mental health services</i>										
Pre-school (0-<5 years)	%	0.8	0.9	0.6	0.5	0.8	0.4	0.3	0.1	0.7
Primary school (5-<12 years)	%	4.3	5.6	4.2	3.0	4.6	3.6	3.2	1.2	4.4
Secondary school (12-<18 years)	%	7.0	8.1	6.7	5.5	7.0	6.8	7.6	2.2	7.0
Youth/young adult (18-<25 years)	%	8.8	9.3	8.5	6.4	8.7	9.5	7.4	3.8	8.5
All children and young people aged <25 years	%	5.5	6.4	5.3	4.1	5.7	5.4	5.0	2.0	5.5
<i>2013-14</i>										
<i>Number of children or young adults who have had contact</i>										
Pre-school (0-<5 years)	no.	4 191	3 417	2 292	840	971	159	98	51	12 019
Primary school (5-<12 years)	no.	30 859	29 925	21 884	7 784	7 342	1 948	1 163	357	101 264
Secondary school (12-<18 years)	no.	42 464	35 174	27 742	11 273	9 259	3 024	2 246	469	131 657
Youth/young adult (18-<25 years)	no.	66 526	56 911	43 070	18 648	15 051	4 736	3 588	1 054	209 589
All children and young people aged <25 years	no.	144 040	125 428	94 988	38 545	32 624	9 866	7 095	1 930	454 528
<i>Proportion of population who had contact with MBS-subsidised primary mental health services</i>										
Pre-school (0-<5 years)	%	0.9	0.9	0.7	0.5	1.0	0.5	0.4	0.3	0.8
Primary school (5-<12 years)	%	4.8	6.1	5.1	3.5	5.4	4.4	3.6	1.4	5.0
Secondary school (12-<18 years)	%	7.8	8.6	7.6	6.0	7.7	7.7	8.6	2.4	7.7

TABLE 12A.45

Table 12A.45 **Young people who had contact with MBS subsidised primary mental health care services, by age group (a), (b), (c), (d), (e), (f), (g)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (h)</i>
Youth/young adult (18-<25 years)	%	9.5	10.1	9.4	7.4	9.6	10.5	8.4	4.0	9.3
All children and young people aged <25 years	%	6.0	6.8	6.0	4.6	6.3	6.2	5.6	2.1	6.1
<i>2014-15</i>										
<i>Number of children or young adults who have had contact</i>										
Pre-school (0-<5 years)	no.	4 609	3 652	2 599	928	1 066	168	128	56	13 206
Primary school (5-<12 years)	no.	34 639	33 696	25 440	9 029	8 184	2 222	1 375	406	114 992
Secondary school (12-<18 years)	no.	46 246	37 731	30 973	12 581	10 111	3 381	2 359	582	143 965
Youth/young adult (18-<25 years)	no.	73 402	62 085	48 032	21 225	16 616	5 108	4 038	1 191	231 697
All children and young people aged <25 years	no.	158 896	137 165	107 045	43 763	35 978	10 878	7 900	2 235	503 859
<i>Proportion of population who had contact with MBS subsidised primary mental health services</i>										
Pre-school (0-<5 years)	%	1.0	1.0	0.8	0.5	1.1	0.5	0.5	0.3	0.9
Primary school (5-<12 years)	%	5.3	6.8	5.8	3.9	5.9	5.0	4.2	1.6	5.6
Secondary school (12-<18 years)	%	8.5	9.2	8.5	6.7	8.5	8.7	9.1	2.9	8.4
Youth/young adult (18-<25 years)	%	10.3	10.9	10.4	8.5	10.6	11.3	9.8	4.7	10.2
All children and young people aged <25 years	%	6.6	7.4	6.8	5.2	7.0	6.8	6.2	2.5	6.7

- (a) Totals do not equal the sum of all mental health providers as data excludes psychiatrists. MBS items included for this indicator are as follows:
- Clinical psychologist services: MBS items 80000, 80005, 80010, 80015, 80020
 - GP services: MBS items 170, 171, 172, 2574, 2575, 2577, 2578, 2700, 2701, 2702, 2704, 2705, 2707, 2708, 2710, 2712, 2713, 2715, 2717, 2719, 2721, 2723, 2725, 2727
 - Other allied health services: MBS items 10956, 10968, 80100, 80105, 80110, 80115, 80120, 80125, 80130, 80135, 80140, 80145, 80150, 80155, 80160, 80165, 80170, 81325, 81355, 82000, 82015.
- (b) Data are based on the date the claim was processed.
- (c) Age of the patient is based on age at last service during the reporting period. Note that in previous years, data supplied for this indicator calculated each patient's age at 30 June of each reference year. The derived rates may differ to those published in previous reports.
- (d) A person is counted if any of the specified mental health item has been used in the reference period.
- (e) A patient is allocated to a state/territory based on their location as at the last service in the reference period.

Table 12A.45 **Young people who had contact with MBS subsidised primary mental health care services, by age group (a), (b), (c), (d), (e), (f), (g)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i> (h)
(f)	The allocation to the state or territory uses a concordance (ABS ASGS 2011 Postcode to Remoteness Area/State) and splits a person where the postcode covers more than one state/territory, therefore the totals may not equal the sum of the individual cells due to rounding.									
(g)	The population data used in this table are the June estimate before the relevant financial year. For 2012-13 data, the estimate is June 2012. The derived rates may differ to those published in previous reports.									
(h)	The sum of the states and territories may not add to the Australian totals as the Australian totals include young people who could not be allocated to a State or Territory.									

Source: Australian Government Department of Health (unpublished); ABS (unpublished) *Australian Demographic Statistics*, Cat. no. 3101.0.

TABLE 12A.46

Table 12A.46 Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by selected characteristics (per cent) (a), (b), (c), (d), (e), (f), (g)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (h)</i>
<i>2010-11</i>									
Gender									
Male	3.8	4.3	3.5	2.8	4.0	3.4	2.9	1.2	3.7
Female	5.1	5.7	4.8	4.3	5.5	5.3	4.8	1.9	5.1
Remoteness areas (f)									
Major cities	4.3	4.8	4.4	3.7	5.0	..	3.8	..	4.4
Inner regional	5.1	5.7	4.5	3.5	4.8	4.7	5.0
Outer regional	3.7	4.7	3.2	3.1	3.5	3.6	..	1.8	3.4
Remote	1.9	3.2	2.0	1.5	2.6	2.2	..	0.9	1.8
Very Remote	1.5	..	1.0	0.7	0.9	2.9	..	0.2	0.7
SEIFA quintiles (f)									
Quintile 1 (most disadvantaged)	4.5	4.7	3.6	4.8	3.8	2.9	4.3	0.4	4.1
Quintile 2	3.7	5.2	4.9	3.2	6.3	9.6	2.4	3.7	4.4
Quintile 3	5.0	5.0	5.3	2.4	4.5	4.6	3.9	1.2	4.5
Quintile 4	4.1	5.3	3.7	3.2	5.6	6.1	4.9	1.5	4.3
Quintile 5 (least disadvantaged)	5.1	4.6	3.5	5.2	3.6	..	3.5	2.0	4.5
Indigenous status									
Aboriginal and Torres Strait Islander	5.3	7.1	3.3	2.0	3.9	5.8	6.3	0.6	3.9
Non-Indigenous	4.4	5.0	4.2	3.6	4.8	4.2	3.8	2.2	4.4
<i>2011-12</i>									
Gender									
Male	4.2	4.8	3.9	3.0	4.4	3.8	3.2	1.5	4.1
Female	5.6	6.4	5.3	4.4	6.0	5.7	5.2	2.3	5.6
Remoteness areas (f)									
Major cities	4.7	5.5	5.0	3.8	5.5	..	4.2	..	4.9
Inner regional	5.6	6.2	4.8	4.1	5.5	4.9	-	..	5.4
Outer regional	4.1	5.0	3.6	3.0	3.7	4.2	..	2.0	3.7
Remote	2.4	3.6	1.9	1.4	2.6	2.6	..	1.2	1.8
Very Remote	1.6	..	0.9	0.8	0.8	3.6	..	0.3	0.7
SEIFA quintiles (f)									
Quintile 1 (most disadvantaged)	4.5	5.6	4.1	5.6	4.7	3.4	2.0	0.5	4.5
Quintile 2	4.3	5.2	5.2	5.5	5.2	7.4	5.7	2.8	4.9
Quintile 3	4.9	6.3	5.4	2.4	6.7	5.2	13.6	2.0	4.9
Quintile 4	5.7	5.3	4.2	3.0	5.8	6.1	6.8	2.1	4.8
Quintile 5 (least disadvantaged)	5.6	5.6	4.2	4.9	3.7	..	3.3	1.3	5.0

TABLE 12A.46

Table 12A.46 Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by selected characteristics (per cent) (a), (b), (c), (d), (e), (f), (g)

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aust (h)
Indigenous status									
Aboriginal and Torres Strait Islander	6.1	7.9	3.8	2.2	4.5	6.2	6.5	0.8	4.5
Non-Indigenous	4.8	5.6	4.7	3.8	5.2	4.6	4.1	2.6	4.9
2012-13									
Gender									
Male	4.7	5.4	4.4	3.2	4.8	4.3	3.8	1.5	4.6
Female	6.4	7.3	6.2	5.0	6.6	6.6	6.2	2.4	6.4
Remoteness areas (f)									
Major cities	5.4	6.2	5.8	4.3	6.0	..	4.9	..	5.6
Inner regional	6.5	7.2	5.4	4.8	6.3	5.6	-	..	6.2
Outer regional	4.7	5.4	4.2	3.3	4.1	5.0	..	2.2	4.3
Remote	2.7	4.0	1.9	1.7	3.1	3.3	..	1.3	2.0
Very Remote	1.5	..	0.9	0.9	1.3	3.6	..	0.3	0.8
SEIFA quintiles (f)									
Quintile 1 (most disadvantaged)	5.1	5.8	5.3	3.4	5.6	5.1	5.8	0.5	5.1
Quintile 2	5.6	6.4	5.5	3.9	5.7	5.4	5.1	2.1	5.5
Quintile 3	5.9	6.7	5.5	4.1	5.6	5.4	5.1	2.0	5.7
Quintile 4	5.6	6.5	5.3	4.1	5.9	5.9	5.2	2.1	5.6
Quintile 5 (least disadvantaged)	5.6	6.3	5.1	4.3	5.7	6.2	4.8	2.0	5.4
Indigenous status									
Aboriginal and Torres Strait Islander	7.1	9.1	4.6	2.4	5.3	6.6	8.1	0.8	5.2
Non-Indigenous	5.5	6.4	5.4	4.3	5.7	5.3	4.9	2.8	5.5
2013-14									
Gender									
Male	5.2	5.9	5.2	3.7	5.4	4.9	4.3	1.7	5.1
Female	7.0	7.8	7.0	5.6	7.3	7.5	6.9	2.6	7.0
Remoteness areas (f)									
Major cities	5.8	6.6	6.4	4.7	6.6	..	5.5	..	6.0
Inner regional	7.3	7.9	6.3	5.5	7.1	6.4	-	..	7.1
Outer regional	5.5	6.0	5.0	4.0	4.6	5.6	..	2.5	4.9
Remote	3.7	5.3	2.5	2.0	3.6	3.7	..	1.1	2.4
Very Remote	2.3	..	1.1	1.3	1.6	4.4	..	0.3	1.0
SEIFA quintiles (f)									
Quintile 1 (most disadvantaged)	5.6	6.2	6.0	3.7	6.2	5.6	6.3	0.5	5.6
Quintile 2	6.2	7.0	6.3	4.3	6.4	6.3	5.8	2.3	6.2

Table 12A.46 Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by selected characteristics (per cent) (a), (b), (c), (d), (e), (f), (g)

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aust (h)
Quintile 3	6.4	7.1	6.2	4.9	6.2	6.2	5.9	1.9	6.3
Quintile 4	6.0	7.0	6.0	4.5	6.6	6.8	5.9	2.4	6.2
Quintile 5 (least disadvantaged)	6.1	6.7	5.7	4.7	6.3	7.2	5.3	2.2	5.9
Indigenous status									
Aboriginal and Torres Strait Islander	7.7	10.4	5.7	2.9	5.7	6.7	8.6	0.9	5.9
Non-Indigenous	6.0	6.8	6.1	4.8	6.4	6.1	5.5	3.0	6.1
2014-15 (i)									
Gender									
Male	5.7	6.4	5.9	4.3	6.1	5.6	4.8	2.1	5.7
Female	7.6	8.4	7.7	6.2	7.9	8.1	7.6	3.0	7.6
Remoteness areas (f)									
Major cities	6.3	7.1	7.1	5.4	7.3	..	6.2	..	6.6
Inner regional	8.1	8.9	7.2	6.1	8.0	7.1	2.1	..	7.9
Outer regional	6.3	7.0	5.6	4.7	4.9	6.3	..	3.6	5.7
Remote	3.6	5.7	2.9	2.3	3.9	4.8	..	1.3	2.7
Very Remote	2.5	..	1.3	1.6	1.6	3.6	..	1.1	1.4
Indigenous status									
Aboriginal and Torres Strait Islander	8.8	11.2	6.3	3.4	7.2	7.3	9.9	1.2	6.7
Non-Indigenous	6.5	7.4	6.8	5.3	7.0	6.8	6.1	3.4	6.7

(a) Totals do not equal the sum of all mental health providers as data excludes psychiatrists. MBS items included for this indicator are as follows:

– Clinical psychologist services: MBS items 80000, 80005, 80010, 80015, 80020

– GP services: MBS items 170, 171, 172, 2574, 2575, 2577, 2578, 2700, 2701, 2702, 2704, 2705, 2707, 2708, 2710, 2712, 2713, 2715, 2717, 2719, 2721, 2723, 2725, 2727

– Other allied health services: MBS items 10956, 10968, 80100, 80105, 80110, 80115, 80120, 80125, 80130, 80135, 80140, 80145, 80150, 80155, 80160, 80165, 80170, 81325, 81355, 82000, 82015.

(b) Data are based on the date the claim was processed.

(c) Age of the patient is based on age at last service during the reporting period. Note that in previous years, data supplied for this indicator calculated each patient's age at 30 June of each reference year. The derived rates may differ to those published in previous reports.

(d) A person is counted if any of the specified mental health item has been used in the reference period.

(e) A patient is allocated to a state/territory based on their location as at the last service in the reference period.

(f) The allocation to the state or territory uses a concordance (ABS ASGS 2011 Postcode to Remoteness Area/State) and splits a person where the postcode covers more than one state/territory, therefore the totals may not equal the sum of the individual cells due to rounding.

(g) The derived rates may differ to those published in previous reports as they may be derived using updated populations.

Table 12A.46 Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by selected characteristics (per cent) (a), (b), (c), (d), (e), (f), (g)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i> (h)
--	------------	------------	------------	-----------	-----------	------------	------------	-----------	-----------------

(h) The sum of the states and territories may not add to the Australian totals as the Australian totals include young people who could not be allocated to a State or Territory.

(i) SEIFA quintile proportions are not available as the populations required to derive them are not available.

Source: Australian Government Department of Health (unpublished); ABS (unpublished) *Australian Demographic Statistics*, Cat. no. 3101.0.

Table 12A.47 Proportion of young people (aged < 25 years) who had contact with MBS subsidised primary mental health care services, by service type (per cent) (a), (b), (c), (d), (e), (f)

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aust (g)
2010-11									
GP services	3.8	4.2	3.6	3.0	4.1	3.6	3.3	1.4	3.8
Clinical psychologist services	0.9	0.9	0.7	1.1	1.6	1.1	1.0	0.2	0.9
Other allied health services	1.8	2.2	1.7	0.9	1.1	1.6	1.5	0.5	1.7
2011-12									
GP services	4.2	4.7	4.0	3.1	4.4	3.9	3.6	1.6	4.1
Clinical psychologist services	1.0	1.1	0.9	1.2	1.9	1.2	1.3	0.2	1.1
Other allied health services	2.0	2.5	1.9	1.0	1.3	1.8	1.4	0.7	1.9
2012-13									
GP services	4.8	5.4	4.6	3.5	4.9	4.5	4.2	1.7	4.7
Clinical psychologist services	1.2	1.3	1.0	1.3	2.0	1.6	1.5	0.2	1.3
Other allied health services	2.2	2.8	2.2	1.1	1.5	1.9	1.7	0.6	2.1
2013-14									
GP services	5.2	5.8	5.3	4.0	5.4	5.2	4.8	1.9	5.2
Clinical psychologist services	1.4	1.5	1.2	1.4	2.3	2.0	1.7	0.2	1.5
Other allied health services	2.3	2.9	2.5	1.3	1.7	2.0	1.9	0.7	2.3
2014-15									
GP services	5.8	6.4	5.9	4.6	6.0	5.8	5.3	2.3	5.8
Clinical psychologist services	1.5	1.6	1.3	1.5	2.6	2.2	2.0	0.2	1.6
Other allied health services	2.5	3.2	2.8	1.5	1.8	2.2	2.0	0.8	2.5

(a) Data excludes psychiatrists. MBS items included for this indicator are as follows:

– Clinical psychologist services: MBS items 80000, 80005, 80010, 80015, 80020

– GP services: MBS items 170, 171, 172, 2574, 2575, 2577, 2578, 2700, 2701, 2702, 2704, 2705, 2707, 2708, 2710, 2712, 2713, 2715, 2717, 2719, 2721, 2723, 2725, 2727

– Other allied health services: MBS items 10956, 10968, 80100, 80105, 80110, 80115, 80120, 80125, 80130, 80135, 80140, 80145, 80150, 80155, 80160, 80165, 80170, 81325, 81355, 82000, 82015.

(b) Data are based on the date the claim was processed.

(c) Age of the patient is based on age at last service during the reporting period. Note that in previous years, data supplied for this indicator calculated each patient's age at 30 June of each reference year.

(d) A person is counted if any of the specified mental health item has been used in the reference period.

(e) A patient is allocated to a state/territory based on their location as at the last service in the reference period.

(f) The population data used in this table are the June estimate before the the relevant financial year. For 2012-13 data, the estimate is June 2012.

(g) The Australian total rates include young people who could not be allocated to a State or Territory.

Source: Australian Government Department of Health (unpublished); ABS (unpublished) *Australian Demographic Statistics*, Cat. no. 3101.0; ABS (2014) *Australian Demographic Statistics*, Cat. no. 3101.0.

Table 12A.48 **Specialised public mental health services reviewed against National Standards for Mental Health Services, 30 June (a)**

		<i>NSW (b)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Expenditure on services assessed at level 1										
2006	\$'000	641 641	574 931	380 642	129 288	153 479	32 236	36 950	22 820	1 971 986
2007	\$'000	556 183	586 248	410 814	95 750	190 360	33 997	46 838	25 537	1 945 727
2008	\$'000	770 511	635 893	526 682	134 530	104 592	42 635	48 458	28 062	2 291 362
2009	\$'000	880 733	681 385	586 763	187 961	100 433	50 559	54 558	30 202	2 572 592
2010	\$'000	851 044	714 515	611 262	178 483	270 545	16 252	54 835	32 326	2 729 262
2011	\$'000	921 406	762 949	699 580	212 630	276 680	45 469	57 536	35 230	3 011 479
2012	\$'000	901 034	525 579	759 987	299 748	124 058	–	62 122	39 291	2 711 818
2013	\$'000	1 069 928	667 682	768 420	432 525	127 752	8 272	64 666	–	3 139 245
2014	\$'000	998 999	615 783	640 373	431 270	–	–	67 471	45 466	2 799 361
Expenditure on services assessed at level 2										
2006	\$'000	–	–	602	12 993	2 013	11 126	–	–	26 734
2007	\$'000	18 413	–	236	168 105	1 409	3 363	–	–	191 526
2008	\$'000	33 962	190	1 770	170 831	1 594	–	–	–	208 347
2009	\$'000	44 946	70	1 234	171 349	1 175	6 171	–	–	224 946
2010	\$'000	217 392	4 117	1 671	174 807	–	–	–	–	397 987
2011	\$'000	236 712	86	–	–	–	49 232	–	–	286 030
2012	\$'000	64 055	272	1 330	53 701	157 099	–	–	–	276 456
2013	\$'000	2 767	103	–	10 096	81 609	–	–	41 329	135 904
2014	\$'000	–	–	–	5 786	220 192	–	–	–	225 978
Expenditure on services assessed at level 3										
2006	\$'000	94 363	18 628	14 377	147 659	42 422	14 212	–	–	331 661
2007	\$'000	220 311	13 383	51 891	45 173	31 781	8 970	–	–	371 509
2008	\$'000	63 334	148	16 771	38 271	135 413	18 753	–	–	272 689
2009	\$'000	71 549	21 630	1 772	16 283	164 555	21 880	–	–	297 669
2010	\$'000	486	23 010	52 296	38 423	2 116	74 572	–	–	190 903
2011	\$'000	490	16 128	3 692	124 290	10 518	–	–	–	155 119
2012	\$'000	177 030	15 709	–	84 463	–	88 003	–	–	365 206
2013	\$'000	168 117	61 161	1 492	54 206	78 580	5 165	–	–	368 720
2014	\$'000	245 526	52 865	141 737	71 053	128 496	90 603	–	–	730 280
Expenditure on services assessed at level 4										
2006	\$'000	46 246	1 073	4 326	–	1 418	2 328	–	–	55 391
2007	\$'000	61 105	1 107	3 694	–	2 180	24 165	–	–	92 252
2008	\$'000	37 887	4 911	462	2 220	3 507	16 235	–	–	65 223
2009	\$'000	3 107	4 143	655	6 304	2 220	2 653	–	–	19 082
2010	\$'000	12 602	8 940	815	7 927	6 611	–	–	–	36 895
2011	\$'000	12 122	15 616	1 971	98 024	1 124	–	–	–	128 858
2012	\$'000	101 432	287 982	926	38 667	16 194	–	–	–	445 201
2013	\$'000	69 759	122 910	940	7 041	15 526	76 378	–	572	293 127
2014	\$'000	145 364	238 818	1 065	19 934	–	–	–	–	405 180

Table 12A.48 **Specialised public mental health services reviewed against National Standards for Mental Health Services, 30 June (a)**

		<i>NSW (b)</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Expenditure on specialised public mental health services										
2006	\$'000	782 250	594 633	399 947	289 939	199 332	59 901	36 950	22 820	2 385 771
2007	\$'000	856 012	600 739	466 636	309 027	225 730	70 494	46 838	25 537	2 601 014
2008	\$'000	905 693	641 143	545 686	345 852	245 106	77 623	48 458	28 062	2 837 621
2009	\$'000	1 000 336	707 227	590 424	381 897	268 383	81 263	54 558	30 202	3 114 289
2010	\$'000	1 081 524	750 582	666 043	399 640	279 273	90 824	54 835	32 326	3 355 046
2011	\$'000	1 170 730	794 780	705 243	434 944	288 323	94 701	57 536	35 230	3 581 486
2012	\$'000	1 243 551	829 543	762 243	476 579	297 351	88 003	62 122	39 291	3 798 683
2013	\$'000	1 310 571	851 856	770 852	503 868	303 467	89 815	64 666	41 901	3 936 995
2014	\$'000	1 389 889	907 465	783 174	528 043	348 688	90 603	67 471	45 466	4 160 799
Per cent of expenditure on services assessed at level 1										
2006	%	82.0	96.7	95.2	44.6	77.0	53.8	100.0	100.0	82.7
2007	%	65.0	97.6	88.0	31.0	84.3	48.2	100.0	100.0	74.8
2008	%	85.1	99.2	96.5	38.9	42.7	54.9	100.0	100.0	80.7
2009	%	88.0	96.3	99.4	49.2	37.4	62.2	100.0	100.0	82.6
2010	%	78.7	95.2	91.8	44.7	96.9	17.9	100.0	100.0	81.3
2011	%	78.7	96.0	99.2	48.9	96.0	48.0	100.0	100.0	84.1
2012	%	72.5	63.4	99.7	62.9	41.7	–	100.0	100.0	71.4
2013	%	81.6	78.4	99.7	85.8	42.1	9.2	100.0	–	79.7
2014	%	71.9	67.9	81.8	81.7	–	–	100.0	100.0	67.3
Per cent of expenditure on services assessed at level 2										
2006	%	–	–	0.2	4.5	1.0	18.6	–	–	1.1
2007	%	2.2	–	0.1	54.4	0.6	4.8	–	–	7.4
2008	%	3.7	–	0.3	49.4	0.7	–	–	–	7.3
2009	%	4.5	–	0.2	44.9	0.4	7.6	–	–	7.2
2010	%	20.1	0.5	0.3	43.7	–	–	–	–	11.9
2011	%	20.2	–	–	–	–	52.0	–	–	8.0
2012	%	5.2	–	0.2	11.3	52.8	–	–	–	7.3
2013	%	0.2	–	–	2.0	26.9	–	–	98.6	3.5
2014	%	–	–	–	1.1	63.1	–	–	–	5.4
Per cent of expenditure on services assessed at level 3										
2006	%	12.1	3.1	3.6	50.9	21.3	23.7	–	–	13.9
2007	%	25.7	2.2	11.1	14.6	14.1	12.7	–	–	14.3
2008	%	7.0	–	3.1	11.1	55.2	24.2	–	–	9.6
2009	%	7.2	3.1	0.3	4.3	61.3	26.9	–	–	9.6
2010	%	–	3.1	7.9	9.6	0.8	82.1	–	–	5.7
2011	%	–	2.0	0.5	28.6	3.6	–	–	–	4.3
2012	%	14.2	1.9	–	17.7	–	100.0	–	–	9.6
2013	%	12.8	7.2	0.2	10.8	25.9	5.8	–	–	9.4
2014	%	17.7	5.8	18.1	13.5	36.9	100.0	–	–	17.6
Per cent of expenditure on services assessed at level 4										
2006	%	5.9	0.2	1.1	–	0.7	3.9	–	–	2.3

Table 12A.48 **Specialised public mental health services reviewed against National Standards for Mental Health Services, 30 June (a)**

		<i>NSW</i> (b)	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
2007	%	7.1	0.2	0.8	–	1.0	34.3	–	–	3.5
2008	%	4.2	0.8	0.1	0.6	1.4	20.9	–	–	2.3
2009	%	0.3	0.6	0.1	1.7	0.8	3.3	–	–	0.6
2010	%	1.2	1.2	0.1	2.0	2.4	–	–	–	1.1
2011	%	1.0	2.0	0.3	22.5	0.4	–	–	–	3.6
2012	%	8.2	34.7	0.1	8.1	5.4	–	–	–	11.7
2013	%	5.3	14.4	0.1	1.4	5.1	85.0	–	1.4	7.4
2014	%	10.5	26.3	0.1	3.8	–	–	–	–	9.7

- (a) There is variation across jurisdictions in the method used to assign an assessment level (1, 2, 3 or 4) to service units. In some jurisdictions, if an organisation with multiple service units is assessed at a particular level all the expenditure on the organisation's units is 'counted' at that assessment level. In other jurisdictions, assessment levels are assigned at the service unit and this may or may not be consistent with the other units within the organisation. The approach can also vary across organisations within a single jurisdiction.
- (b) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.

– Nil or rounded to zero.

Source: AIHW (unpublished) MHE NMDS.

Table 12A.49 Recurrent expenditure on community-based services as a proportion of total spending on mental health services (per cent) (a), (b), (c)

	<i>NSW</i> (d)	<i>Vic</i>	<i>Qld</i> (e)	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
2005-06	43.5	63.5	43.3	51.3	45.1	62.4	79.2	60.4	51.0
2006-07	43.7	63.3	48.1	52.0	45.6	59.3	73.3	65.7	51.5
2007-08	46.1	62.6	49.1	53.1	47.9	57.8	71.5	64.0	52.5
2008-09	44.5	62.6	51.8	53.6	49.9	57.7	74.2	62.6	52.7
2009-10	44.6	62.8	54.0	54.4	52.4	54.8	74.7	65.4	53.4
2010-11	44.3	63.2	55.5	53.4	56.5	56.5	73.2	64.3	53.9
2011-12	43.7	64.2	56.3	53.4	58.8	56.8	74.4	63.9	54.2
2012-13	41.8	64.1	55.3	52.9	61.5	58.8	73.3	62.1	53.5
2013-14	41.3	64.7	55.2	53.1	59.6	59.7	73.0	63.7	53.3

(a) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure estimates.

(b) Due to the ongoing validation of the NMDS, data could differ from previous reports.

(c) Recurrent expenditure exclude indirect and aged care residential expenditure.

(d) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.

(e) Queensland does not fund community-based residential services, but funds extended treatment (campus-based and non-campus-based) services that provide longer term inpatient treatment and rehabilitation services with clinical staffing for 24 hours a day, 7 days a week.

Source: AIHW (unpublished) MHE NMDS.

TABLE 12A.50

Table 12A.50 Specialised public mental health services episodes with completed consumer outcomes measures collected (a), (b)

<i>Unit</i>	<i>NSW</i>	<i>Vic (c)</i>	<i>Qld</i>	<i>WA</i>	<i>SA Tas (c)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (c)</i>	
2007-08									
Group A: People discharged from hospital (d)									
no.	5 989	3 740	4 419	2 564	2 657	324	40	92	19 825
%	29.7	28.0	42.0	43.0	50.4	19.7	4.6	16.1	34.0
Group B: People discharged from community-based ambulatory care (e)									
no.	2 126	3 938	6 065	1 196	1 457	366	np	51	15 199
%	12.0	33.9	39.5	21.5	30.4	22.3	np	6.2	25.7
Group C: People in ongoing community-based ambulatory care (f)									
no.	5 073	5 307	5 917	2 760	3 097	705	159	305	23 323
%	16.5	27.4	31.5	26.1	39.7	19.3	5.6	23.3	24.8
2008-09									
Group A: People discharged from hospital (d)									
no.	5 605	6 350	2 205	2 944	2 360	321	46	104	19 935
%	27.8	46.5	20.6	47.9	46.3	20.2	4.9	18.2	33.8
Group B: People discharged from community-based ambulatory care (e)									
no.	1 985	6 804	3 577	1 162	1 420	305	np	25	15 278
%	10.3	62.3	19.3	18.8	27.2	21.2	np	3.3	23.7
Group C: People in ongoing community-based ambulatory care (f)									
no.	5 108	6 472	5 759	3 558	3 340	712	175	383	25 507
%	16.1	34.0	34.0	30.9	37.7	21.3	5.6	25.0	27.1
2009-10									
Group A: People discharged from hospital (d)									
no.	6 146	7 845	1 736	2 945	2 490	316	67	146	21 691
%	30.2	55.7	16.2	44.4	46.9	–	7.6	26.3	36.1
Group B: People discharged from community-based ambulatory care (e)									
no.	2 024	8 618	2 706	1 329	1 510	291	np	48	16 526
%	9.9	77.3	17.7	20.6	28.9	24.0	np	6.0	27.0
Group C: People in ongoing community-based ambulatory care (f)									
no.	5 943	7 895	6 544	4 064	3 201	685	335	396	29 063
%	17.5	44.1	32.0	35.0	36.3	30.1	10.0	23.8	29.4
2010-11									
Group A: People discharged from hospital (d)									
no.	5 937	8 249	2 515	3 236	2 288	443	87	200	22 955
%	30.1	57.1	22.6	45.3	39.3	30.8	8.9	34.6	37.5
Group B: People discharged from community-based ambulatory care (e)									
no.	2 309	10 243	3 537	1 351	1 473	583	np	50	19 546
%	11.0	80.4	21.7	18.4	25.2	39.2	np	6.6	28.6
Group C: People in ongoing community-based ambulatory care (f)									
no.	6 020	8 165	7 146	4 453	3 150	703	466	354	30 457
%	18.1	45.7	35.1	36.3	36.3	31.8	13.8	20.3	30.8

Table 12A.50 Specialised public mental health services episodes with completed consumer outcomes measures collected (a), (b)

<i>Unit</i>	<i>NSW</i>	<i>Vic (c)</i>	<i>Qld</i>	<i>WA</i>	<i>SA Tas (c)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (c)</i>	
2011-12 (c)									
Group A: People discharged from hospital (d)									
no.	6 095	na	3 377	3 614	2 307	774	33	223	16 423
%	29.4	na	28.8	49.5	42.5	50.4	3.3	36.2	34.0
Group B: People discharged from community-based ambulatory care (e)									
no.	2 501	na	3 227	1 332	1 438	294	np	48	8 840
%	12.7	na	18.1	16.7	22.6	17.7	np	5.1	15.9
Group C: People in ongoing community-based ambulatory care (f)									
no.	7 498	na	7 133	3 651	3 200	541	276	402	22 701
%	21.1	na	34.1	29.1	34.8	23.3	7.8	19.7	26.5
2012-13 (c)									
Group A: People discharged from hospital (d)									
no.	7 088	na	4 664	3 623	2 149	1 010	66	225	18 825
%	32.3	na	36.5	45.7	42.3	54.8	6.0	33.7	36.7
Group B: People discharged from community-based ambulatory care (e)									
no.	2 403	na	4 521	1 404	1 396	392	26	64	10 206
%	12.0	na	23.6	15.7	20.3	40.8	4.7	5.5	17.4
Group C: People in ongoing community-based ambulatory care (f)									
no.	6 460	na	8 254	3 855	3 409	618	302	412	23 310
%	18.9	na	38.9	29.6	36.1	39.0	8.1	18.0	27.4
2013-14									
Group A: People discharged from hospital (d)									
no.	6 289	7 311	5 524	3 415	1 838	669	258	262	25 566
%	27.9	46.1	42.8	41.6	38.1	36.9	25.3	39.1	37.7
Group B: People discharged from community-based ambulatory care (e)									
no.	2 871	9 748	5 288	1 454	1 495	440	np	34	21 330
%	13.9	86.8	26.7	15.1	19.7	25.6	np	2.2	29.4
Group C: People in ongoing community-based ambulatory care (f)									
no.	6 407	6 836	8 684	3 687	3 634	843	294	396	30 781
%	19.8	37.2	23.6	27.3	39.5	25.0	6.9	11.1	27.2

(a) These data were prepared by the Australian Mental Health Outcomes and Classification Network, using data submitted by State and Territory governments to the Australian Government Department of Health. To be counted as an episode for which consumer outcome measures are collected, data need to be completed correctly (a specified minimum number of items completed) and have a 'matching pair' — that is, a beginning and end rating are needed to enable an outcome score to be determined.

(b) Estimates of the number of episodes with complete outcome data for state and territory mental health services for all years are based on an analytic approach that compares the number of episodes with 'matched pairs' outcomes data to data submitted for the various mental health National Minimum Data Sets.

Table 12A.50 **Specialised public mental health services episodes with completed consumer outcomes measures collected (a), (b)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic (c)</i>	<i>Qld</i>	<i>WA</i>	<i>SA Tas (c)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (c)</i>
(c)	Data are not available for Victoria for 2011-12 and 2012-13. All totals for 2011-12 and 2012-13 exclude Victoria. Industrial action in Tasmania has limited the available data quality and quantity of the 2011-12 and 2012-13 data.								
(d)	Group A covers people who received a discrete episode of inpatient care within a state/territory designated psychiatric inpatient unit during the reference year. The defining characteristic of the group is that the episode of inpatient care commenced, and was completed, within the year. Outcome scores were calculated as the difference between the total score recorded at admission and discharge. The analysis excludes episodes where length of stay was three days or less because it is not meaningful to compare admission and discharge ratings for short duration episodes.								
(e)	Group B covers people who received relatively short term community care from a state/territory mental health service during the reference year. The defining characteristic of the group is that the episode of community care commenced, and was completed, within the year. Outcome scores were calculated as the difference between the total score recorded at admission to, and discharge from, community care. A subgroup of people whose episode of community care completed because they were admitted to hospital is not included in this analysis.								
(f)	Group C covers people receiving relatively long term community care from a state/territory mental health service. It includes people who were receiving care for the whole of the reference year, and those who commenced community care sometime after 1 July who continued under care for the rest of the year. The defining characteristic of the group is that all remained in ongoing care when the year ended (30 June). Outcome scores were calculated as the difference between the total score recorded on the first occasion rated and the last occasion rated in the year.								
	na Not available. – Nil or rounded to zero. np Not published.								

Source: AIHW (unpublished) from data provided by the Australian Mental Health Outcomes and Classification Network.

Table 12A.51 Rate and duration of seclusion events in public specialised mental health acute inpatient units (a)

	NSW (b)	Vic (c)	Qld (d)	WA (e)	SA (f), (g)	Tas (h)	ACT (i), (j), (k)	NT (d), (l), (m)	Aust
<i>Seclusion events per 1000 patient bed days</i>									
2008-09	11.0	18.8	18.2	15.3	na	15.4	13.3	na	15.5
2009-10	11.5	19.4	15.0	11.6	7.6	11.5	1.7	23.0	13.5
2010-11	9.4	15.1	17.2	8.3	7.7	14.7	0.7	19.9	11.8
2011-12	9.2	13.3	13.3	4.7	10.1	11.9	1.3	25.9	10.4
2012-13	8.5	10.9	12.7	6.0	9.1	19.7	0.9	16.4	9.6
2013-14	7.4	9.2	11.1	5.2	4.6	15.2	1.1	21.6	8.1
2014-15	7.7	7.5	11.4	4.3	5.0	10.1	2.7	31.0	7.8
<i>Average duration of seclusion events (no. of hours) (n), (o)</i>									
2013-14	6.0	9.5	3.8	2.4	na	4.1	2.1	6.4	6.0
2014-15	5.8	8.0	3.4	2.7	na	2.5	2.2	7.9	5.4

- (a) Variation in jurisdictional legislation may result in differences in the definition of a seclusion event. Data reported by jurisdictions may therefore vary and comparisons should be made with caution.
- (b) NSW does not have a centralised database for the collection of seclusion data. Services report seclusion rates regularly to the NSW Ministry of Health. Services are required to maintain local seclusion registers, which may be audited by NSW Official Visitors. Seclusion rates are a Key Performance Indicator (KPI) in regular performance reporting to NSW Local Health Districts. NSW seclusion rates include bed days for some forensic services managed by correctional facilities.
- (c) Victoria has fewer beds per capita than other jurisdictions, and as such, it may be useful to view the rate of seclusion events in a broader population context (rates per capita).
- (d) Queensland and the NT do not report any acute forensic services, however forensic patients can and do access acute care through general units.
- (e) The WA data include seclusion events (numerator) and patient days (denominator) for both acute and non-acute specialised public mental health units. WA also does not have a centralised data base for the collection of seclusion data. Services provide seclusion data from their own data bases.
- (f) Recent data reporting improvements will impact on SA data. Bed days used to calculate SA's seclusion rates are estimated based on 100 per cent occupied bed numbers, which are fluctuating in relation to new infrastructure projects. The number of bed days is an estimate which affects the rate of seclusion reported for SA and fluctuations in bed numbers can relate to new infrastructure projects. During 2010-11, a substantial number of seclusion events in one particular hospital were for a small number of patients with over half of these patient-requested events. This may have impacted on the overall seclusion rate reported for the state for 2010-11.
- (g) For 2008-09, SA was unable to supply seclusion data.
- (h) The increase in the state-wide Tasmanian seclusion rate for 2012-13 and 2013-14 data is due to a small number of clients having an above average number of seclusion events.
- (i) For the ACT, when interpreting these data, the relative small size of the ACT should be noted, with a total of between 63 and 70 acute inpatient beds reported between 2008-09 and 2013-14.

Table 12A.51 Rate and duration of seclusion events in public specialised mental health acute inpatient units (a)

	<i>NSW</i> (b)	<i>Vic</i> (c)	<i>Qld</i> (d)	<i>WA</i> (e)	<i>SA</i> (f), (g)	<i>Tas</i> (h)	<i>ACT</i> (i), (j), (k)	<i>NT</i> (d), (l), (m)	<i>Aust</i>
(j)									ACT activities initiated as part of the Beacon Site project included the implementation of a clinical review committee inclusive of clinical staff, consumers and carer representation to review episodes of seclusion for systemic issues on a case-by-case basis. This has led to a number of reforms over several years that had a direct impact on the use of seclusion and its reduction to the low levels now reported.
(k)									In the ACT, work is progressive and ongoing as part of a larger process of providing a place of improved safety and security, both for people experiencing an acute episode of mental ill health leading to an inpatient admission, visitors and for the staff who work in this challenging environment.
(l)									Due to the low ratio of beds per person in the NT compared with other jurisdictions, the apparent rate of seclusion is inflated when reporting seclusion per patient day compared with reporting on a population basis. Due to the low number of beds in the NT, high rates of seclusion for a few individuals has a disproportional effect on the rate of seclusion reported. NT seclusion data is therefore not directly comparable with other jurisdictions.
(m)									The NT was unable to supply seclusion data for 2008-09.
(n)									SA report seclusion duration in 4 hour blocks and therefore the mean duration cannot be calculated. The national average seclusion duration figure excludes SA.
(o)									Due to data comparability issues for events occurring in forensic services, all forensic service events are excluded from the average duration analysis.

na Not available.

Source: AIHW (2015) *Mental Health Services in Australia Online*, mhsa.aihw.gov.au/home/ (accessed 16 December 2015).

Table 12A.52 Rate and duration of seclusion events in public specialised mental health acute inpatient units (per 1000 patient bed days), by target population (a), (b)

	<i>Events per 1000 patient bed days</i>							<i>Average duration of events (no. of hours) (c)</i>
	<i>2008-09</i>	<i>2009-10</i>	<i>2010-11</i>	<i>2011-12</i>	<i>2012-13</i>	<i>2013-14</i>	<i>2014-15</i>	<i>2014-15</i>
General	17.1	15.4	13.1	11.6	10.3	9.6	9.1	5.6
Child and adolescent	17.0	11.4	16.6	18.1	14.5	9.6	12.0	1.6
Older people	3.7	2.9	1.2	0.7	0.7	0.5	0.4	4.4
Mixed	15.1	13.3	12.3	10.3	10.0
Forensic (b)	10.8	6.9	5.2	6.8	8.9	5.3	4.9	78.1
Total	15.5	13.5	11.8	10.4	9.6	8.1	7.8	5.4

- (a) See table 12A.51 for general caveats regarding seclusion data.
- (b) Excludes some public sector acute forensic mental health hospital services operated in correctional facilities.
- (c) SA report seclusion duration in 4 hour blocks and therefore the mean duration cannot be calculated. The national average seclusion duration figure excludes SA.
- .. Not applicable.

Source: AIHW (2015) *Mental Health Services in Australia Online*, mhsa.aihw.gov.au/home/ (accessed 16 December 2015).

TABLE 12A.53

Table 12A.53 **Consumer and carer participation (a), (b), (c), (d)**

	NSW (e), (f)	Vic	Qld	WA (g)	SA	Tas	ACT (h)	NT (h)	Aust
<i>Number of consumer and carer consultants</i>									
Number of paid consumer workers (FTE)									
2005-06	27.3	19.6	9.8	0.5	2.8	–	1.3	–	61.3
2006-07	24.8	19.0	10.3	0.8	2.1	–	–	–	57.0
2007-08	27.9	20.0	9.7	1.2	4.7	–	–	–	63.5
2008-09	23.5	17.1	13.6	3.6	6.3	0.5	–	–	64.6
2009-10	21.5	17.7	14.1	5.1	5.7	0.5	–	–	64.6
2010-11	20.5	17.9	17.8	3.3	8.4	0.5	–	–	68.5
2011-12	23.9	19.1	19.5	2.0	8.2	1.5	–	–	74.2
2012-13	17.5	19.4	14.3	4.2	12.5	–	–	0.2	68.0
2013-14	26.5	16.2	8.6	4.7	12.9	0.5	–	0.2	69.5
Number of paid carer workers (FTE)									
2005-06	2.7	11.7	0.4	–	–	–	–	–	14.8
2006-07	8.6	13.6	0.9	–	–	–	–	–	23.1
2007-08	7.0	15.5	1.5	0.8	1.8	–	–	–	26.6
2008-09	10.3	14.3	2.7	0.5	2.4	0.5	–	–	30.6
2009-10	13.7	15.8	4.8	1.0	1.5	0.5	–	–	37.3
2010-11	13.7	17.9	5.3	1.0	5.0	0.5	–	–	43.4
2011-12	15.9	18.5	6.4	0.2	4.2	0.6	–	–	45.8
2012-13	10.2	18.6	3.0	0.2	3.6	1.0	–	–	36.6
2013-14	7.7	18.0	5.2	–	4.8	1.0	–	–	36.8
Number of paid direct care (including, consumer and carer worker positions) (FTE)									
2005-06	6 494.5	5 270.0	3 633.8	2 332.3	1 691.3	607.7	331.3	151.9	20 512.8
2006-07	6 732.0	5 338.0	3 875.8	2 427.1	1 800.9	656.2	321.8	158.5	21 310.3
2007-08	6 777.3	5 440.8	4 233.4	2 537.7	1 963.3	639.7	314.7	167.5	22 074.4

TABLE 12A.53

Table 12A.53 **Consumer and carer participation (a), (b), (c), (d)**

	<i>NSW</i> (e), (f)	<i>Vic</i>	<i>Qld</i>	<i>WA</i> (g)	<i>SA</i>	<i>Tas</i>	<i>ACT</i> (h)	<i>NT</i> (h)	<i>Aust</i>
2008-09	7 025.6	5 634.4	4 405.7	2 670.5	1 977.3	652.6	313.8	193.3	22 873.2
2009-10	7 357.2	5 703.9	4 361.7	2 724.8	2 025.3	682.5	334.5	196.3	23 386.1
2010-11	7 637.3	5 912.7	4 694.2	2 856.0	2 121.6	687.3	338.4	205.3	24 452.7
2011-12	7 898.4	6 049.5	4 991.9	3 017.4	2 045.6	646.8	345.1	216.1	25 210.6
2012-13	8 033.2	6 000.9	5 086.1	3 146.3	2 033.4	628.9	375.5	249.4	25 553.7
2013-14	8 174.4	6 262.7	4 839.6	3 189.6	2 102.5	650.1	390.8	271.2	25 880.9
Paid consumer workers (FTE) per 1000 paid direct care (h)									
2005-06	4.2	3.7	2.7	0.2	1.7	–	3.9	–	3.0
2006-07	3.7	3.6	2.7	0.3	1.2	–	–	–	2.7
2007-08	4.1	3.7	2.3	0.5	2.4	–	–	–	2.9
2008-09	3.3	3.0	3.1	1.4	3.2	0.8	–	–	2.8
2009-10	2.9	3.1	3.2	1.9	2.8	0.7	–	–	2.8
2010-11	2.7	3.0	3.8	1.2	4.0	0.7	–	–	2.8
2011-12	3.0	3.2	3.9	0.7	4.0	2.3	–	–	2.9
2012-13	2.2	3.2	2.8	1.3	6.1	–	–	0.7	2.7
2013-14	3.2	2.6	1.8	1.5	6.1	0.8	–	0.6	2.7
Paid carer workers (FTE) per 1000 paid direct care (FTE) (h)									
2005-06	0.4	2.2	0.1	–	–	–	–	–	0.7
2006-07	1.3	2.5	0.2	–	–	–	–	–	1.1
2007-08	1.0	2.9	0.4	0.3	0.9	–	–	–	1.2
2008-09	1.5	2.5	0.6	0.2	1.2	0.8	–	–	1.3
2009-10	1.9	2.8	1.1	0.4	0.8	0.7	–	–	1.6
2010-11	1.8	3.0	1.1	0.4	2.4	0.7	–	–	1.8
2011-12	2.0	3.1	1.3	0.1	2.1	0.9	–	–	1.8
2012-13	1.3	3.1	0.6	0.1	1.8	1.6	–	–	1.4

Table 12A.53 **Consumer and carer participation (a), (b), (c), (d)**

	NSW (e), (f)	Vic	Qld	WA (g)	SA	Tas	ACT (h)	NT (h)	Aust
2013-14	0.9	2.9	1.1	–	2.3	1.5	–	–	1.4

- (a) Non-government organisations are included only where they provide staffed residential services.
- (b) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of relevant items.
- (c) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (d) Data up to 2009-10 were restricted to consumer/carer consultants. In 2010-11, the definitions were altered to include a broader range of roles in the contemporary mental health environment, transitioning to mental health consumer and carer workers. These improved definitions should promote greater consistency between jurisdictions. Comparisons between data up to 2009-10 with data from 2010-11 should not be made.
- (e) NSW advised that the government has no authority to require consumer participation in services delivered through the primary care program.
- (f) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year.
- (g) WA has advised that this information does not represent the full range of consumer and carer participation. Genuine engagement with consumers and carers is one of the key principles of the Mental Health Commission's Strategic Policy document *Mental Health 2020*. The Commission has allocated funding to establish and support Consumers of Mental Health WA Inc., a peak body that provides systemic advocacy and is run for and by consumers. Other examples include provision of funding to develop the capacity of non-government organisations to employ people with a lived experience of mental illness and awarding scholarships to people with a lived experience to complete approved university and polytechnic studies in mental health. Several key consumer and carer advisory groups are supported and provided with financial assistance and collectively, these groups provide advice and representations on consumer and carer issues. The Commission funds Carers Association of WA for the provision of systemic advocacy services and the Mental Health Carers ARAFMI (WA) for a range of services including individual advocacy.
- (h) Consumer and carer workers are not employed in the ACT (except in 2005-06). The NT do not employ carer staff and employed consumer staff in 2012-13 only.

– Nil or rounded to zero.

Source: AIHW (unpublished) MHE NMDS.

TABLE 12A.54

Table 12A.54 Rates of community follow-up for people within the first seven days of discharge from hospital

	<i>Unit</i>	<i>NSW</i>	<i>Vic (a)</i>	<i>Qld</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (c)</i>	<i>ACT (d)</i>	<i>NT</i>	<i>Aust (e)</i>
<i>2005-06</i>										
Overnight separations from acute psychiatric inpatient services	no.	24 891	14 957	14 326	6 222	5 352	na	1 136	1 004	67 888
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	10 695	8 938	6 488	2 703	1 611	na	769	188	31 392
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	43.0	59.8	45.3	43.4	30.1	na	67.7	18.7	46.2
<i>2006-07</i>										
Overnight separations from acute psychiatric inpatient services	no.	26 656	15 602	13 534	5 994	5 430	na	1 100	888	69 204
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	11 539	9 303	6 833	2 756	1 532	na	759	342	33 064
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	43.3	59.6	50.5	46.0	28.2	na	69.0	38.5	47.8
<i>2007-08</i>										
Overnight separations from acute psychiatric inpatient services	no.	27 103	16 400	13 600	5 863	5 590	2 116	1 148	854	72 674
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	10 856	9 803	7 094	2 778	1 941	433	827	348	34 080

TABLE 12A.54

Table 12A.54 Rates of community follow-up for people within the first seven days of discharge from hospital

	<i>Unit</i>	<i>NSW</i>	<i>Vic (a)</i>	<i>Qld</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (c)</i>	<i>ACT (d)</i>	<i>NT</i>	<i>Aust (e)</i>
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	40.1	59.8	52.2	47.4	34.7	20.5	72.0	40.7	46.9
<i>2008-09</i>										
Overnight separations from acute psychiatric inpatient services	no.	27 035	16 429	14 147	6 272	5 435	2 121	1 233	780	73 452
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	11 078	10 132	6 228	3 070	2 222	461	901	323	34 415
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	41.0	61.7	44.0	48.9	40.9	21.7	73.1	41.4	46.9
<i>2009-10</i>										
Overnight separations from acute psychiatric inpatient services	no.	26 403	16 552	14 061	6 439	5 509	1 758	1 184	742	72 648
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	11 864	10 591	6 417	3 227	2 301	456	873	289	36 018
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	44.9	64.0	45.6	50.1	41.8	25.9	73.7	38.9	49.6
<i>2010-11</i>										
Overnight separations from acute psychiatric inpatient services	no.	26 932	17 156	14 634	7 524	5 825	1 730	1 185	771	75 757

TABLE 12A.54

Table 12A.54 Rates of community follow-up for people within the first seven days of discharge from hospital

	<i>Unit</i>	<i>NSW</i>	<i>Vic (a)</i>	<i>Qld</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (c)</i>	<i>ACT (d)</i>	<i>NT</i>	<i>Aust (e)</i>
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	12 811	11 730	7 696	3 683	2 662	505	932	308	40 327
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	47.6	68.4	52.6	49.0	45.7	29.2	78.6	39.9	53.2
<i>2011-12</i>										
Overnight separations from acute psychiatric inpatient services	no.	27 407	na	15 187	7 800	5 987	1 655	1 306	781	60 123
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	14 348	na	9 838	3 992	3 064	531	1 015	313	33 101
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	52.4	na	64.8	51.2	51.2	32.1	77.7	40.1	55.1
<i>2012-13</i>										
Overnight separations from acute psychiatric inpatient services	no.	28 297	na	15 916	8 705	5 436	1 667	1 307	889	62 217
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	16 828	na	11 598	4 669	2 935	347	966	414	37 757
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	59.5	na	72.9	53.6	54.0	20.8	73.9	46.6	60.7
<i>2013-14</i>										

TABLE 12A.54

Table 12A.54 **Rates of community follow-up for people within the first seven days of discharge from hospital**

	<i>Unit</i>	<i>NSW</i>	<i>Vic (a)</i>	<i>Qld</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (c)</i>	<i>ACT (d)</i>	<i>NT</i>	<i>Aust (e)</i>
Overnight separations from acute psychiatric inpatient services	no.	29 200	18 214	16 401	9 144	4 909	1 855	1 238	952	81 913
Overnight acute separations with community mental health contact recorded in the seven days following separation	no.	18 603	13 159	12 081	5 215	2 836	1 103	898	456	54 351
Overnight separations from acute psychiatric inpatient services with community mental health contact recorded in the seven days following separation	%	63.7	72.2	73.7	57.0	57.8	59.5	72.5	47.9	66.4

- (a) For public sector community mental health services, Victorian data are unavailable for 2011-12 and 2012-13 due to service level collection gaps resulting from protected industrial action during this period.
- (b) SA submitted data that was not based on unique patient identifier but is based on a limited data matching approach. Therefore caution needs to be taken when making interjurisdictional comparisons.
- (c) Industrial action in Tasmania limited the available data quality and quantity of 2011-12 and 2012-13 community data. Tasmanian data are not available for 2005-06 and 2006-07. Data submitted up to 2012-13 were not based on unique patient identifiers or data matching approaches.
- (d) From 2012-13, the ACT has refined its calculation methodology and as such, comparisons to previous years' results should be viewed with caution.
- (e) Due to data supply issues, Australian totals should be interpreted with caution.

na Not available.

Source: AIHW (unpublished), from data provided by State and Territory governments.

Table 12A.55 Rates of community follow-up within first seven days of discharge from a psychiatric admission, by State and Territory, by Indigenous status and remoteness (a), (b)

	<i>Unit</i>	<i>NSW</i>	<i>Vic (c)</i>	<i>Qld</i>	<i>WA</i>	<i>SA (d)</i>	<i>Tas (e)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (f)</i>
<i>2011-12</i>										
Indigenous status										
Aboriginal and Torres Strait Islander	%	45.2	na	61.3	40.3	45.4	na	87.9	32.5	48.3
Non-Indigenous	%	53.0	na	65.4	52.3	52.6	na	78.2	47.9	56.6
Remoteness										
Major cities	%	52.5	na	71.2	52.9	53.5	na	79.5	50.0	55.2
Inner regional	%	54.6	na	63.5	50.7	41.3	na	51.9	25.0	59.3
Outer regional	%	52.8	na	67.7	43.9	41.4	na	100.0	48.9	56.7
Remote	%	39.5	na	59.3	47.5	31.0	na	..	43.1	45.4
Very remote	%	36.4	na	61.9	28.4	34.5	na	..	25.8	33.4
<i>2012-13</i>										
Indigenous status										
Aboriginal and Torres Strait Islander	%	53.9	na	72.2	47.3	39.4	15.1	68.3	40.2	55.3
Non-Indigenous	%	60.0	na	73.2	54.3	55.9	21.3	74.5	53.1	61.4
Remoteness										
Major cities	%	59.4	na	71.2	55.1	56.8	–	75.6	16.7	61.6
Inner regional	%	62.7	na	78.2	52.2	40.2	18.6	32.6	33.3	61.0
Outer regional	%	59.8	na	76.3	47.4	38.6	24.6	19.3	53.8	60.6
Remote	%	38.7	na	64.8	55.7	47.4	31.5	..	54.8	52.3
Very remote	%	60.9	na	72.2	40.4	39.2	22.2	..	33.1	41.7
<i>2013-14</i>										
Indigenous status										
Aboriginal and Torres Strait Islander	%	59.5	69.3	72.9	55.8	39.0	55.6	70.8	39.3	60.5
Non-Indigenous	%	64.2	72.4	74.0	57.2	59.7	59.8	72.7	55.1	66.9
Remoteness										
Major cities	%	65.1	72.2	71.4	57.4	59.8	14.4	76.2	21.1	66.8
Inner regional	%	61.9	73.3	81.1	57.3	50.0	59.6	10.6	37.5	67.6
Outer regional	%	63.2	76.2	78.4	58.3	53.8	63.3	–	49.6	68.4
Remote	%	48.2	69.2	69.6	60.2	40.0	44.7	..	66.7	59.6
Very remote	%	57.7	62.5	74.7	47.0	34.7	53.8	..	29.5	44.0

(a) The Indigenous status rates should be interpreted with caution due to the varying and, in some instances, unknown quality of Indigenous identification across jurisdictions. Excludes people for whom demographic information was missing or not reported.

(b) Disaggregation by remoteness area is based on a person's usual residence, not the location of the service provider. State/territory is the state/territory of the service provider. Excludes people for whom demographic information was missing or not reported.

Table 12A.55 Rates of community follow-up within first seven days of discharge from a psychiatric admission, by State and Territory, by Indigenous status and remoteness (a), (b)

	<i>Unit</i>	<i>NSW</i>	<i>Vic (c)</i>	<i>Qld</i>	<i>WA</i>	<i>SA (d)</i>	<i>Tas (e)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (f)</i>
--	-------------	------------	----------------	------------	-----------	---------------	----------------	------------	-----------	-----------------

(c) For public sector community mental health services, Victorian data for 2011-12 and 2012-13 are unavailable due to service level collection gaps resulting from protected industrial action during this period.

(d) SA submitted data that was not based on unique patient identifier but is based on a limited data matching approach. Therefore caution needs to be taken when making interjurisdictional comparisons.

(e) Industrial action in Tasmania limited the available data quality and quantity of 2011-12 and 2012-13 community data.

(f) Due to data supply issues, Australian totals should be interpreted with caution.

na Not available. **..** Not applicable. **–** Nil or rounded to zero.

Source: AIHW (unpublished), from data provided by State and Territory governments.

Table 12A.56 Rates of community follow-up within first seven days of discharge from a psychiatric admission, by age group, gender and SEIFA quintiles, 2013-14

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA (a)</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Age group</i>										
Less than 15 years	%	54.6	61.2	59.6	50.8	15.8	66.7	–	50.0	56.4
15–24 years	%	63.0	69.5	70.8	58.7	44.8	56.3	79.3	38.2	64.5
25–34 years	%	64.5	74.1	71.8	55.4	59.2	56.8	76.1	47.4	66.4
35–44 years	%	64.3	75.0	74.9	54.5	58.4	54.1	76.2	53.5	67.2
45–54 years	%	64.7	75.4	77.7	60.0	63.6	66.1	67.1	53.3	68.8
55–64 years	%	64.9	73.3	77.3	62.4	62.9	63.2	70.3	52.4	68.6
65+ years	%	58.4	65.5	79.5	55.8	64.0	75.3	52.4	50.0	63.7
<i>Gender</i>										
Male	%	61.9	72.2	73.2	53.5	57.4	56.2	72.5	48.3	65.1
Female	%	65.6	72.3	74.1	60.3	58.1	62.8	72.6	47.3	67.6
<i>SEIFA quintiles</i>										
Quintile 1 (most disadvantaged)	%	64.9	73.4	76.3	60.0	55.2	61.8	20.0	32.9	67.5
Quintile 2	%	62.4	72.2	77.1	59.3	56.9	61.1	66.7	47.2	66.1
Quintile 3	%	62.9	73.9	74.6	57.3	53.5	58.5	40.0	63.4	67.1
Quintile 4	%	65.9	71.2	71.0	58.2	66.4	55.3	79.1	42.8	67.5
Quintile 5 (least disadvantaged)	%	65.9	71.8	70.1	52.3	61.2	52.4	75.0	43.9	65.9

(a) SA submitted data that was not based on unique patient identifier but is based on a limited data matching approach. Therefore caution needs to be taken when making interjurisdictional comparisons.

– Nil or rounded to zero.

Source: AIHW (unpublished), from data provided by State and Territory governments.

TABLE 12A.57

Table 12A.57 Readmissions to hospital within 28 days of discharge (a), (b), (c), (d)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (b)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>2005-06</i>										
Overnight separations from psychiatric acute inpatient services	no.	25 087	14 957	14 211	6 644	5 352	2 617	1 136	1 004	71 008
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 057	2 098	2 696	933	629	334	152	140	11 039
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	16.2	14.0	19.0	14.0	11.8	12.8	13.4	13.9	15.5
<i>2006-07</i>										
Overnight separations from psychiatric acute inpatient services	no.	26 767	15 602	13 432	6 476	5 430	1 901	1 100	888	71 596
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 526	2 309	2 110	822	491	126	123	123	10 630
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	16.9	14.8	15.7	12.7	9.0	6.6	11.2	13.9	14.8
<i>2007-08</i>										
Overnight separations from psychiatric acute inpatient services	no.	27 202	16 400	13 296	6 447	5 590	2 046	1 148	848	72 977

TABLE 12A.57

Table 12A.57 Readmissions to hospital within 28 days of discharge (a), (b), (c), (d)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (b)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 716	2 484	2 059	857	616	167	114	111	11 124
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	17.3	15.1	15.5	13.3	11.0	8.2	9.9	13.1	15.2
<i>2008-09</i>										
Overnight separations from psychiatric acute inpatient services	no.	27 101	16 429	13 827	6 881	5 431	1 823	1 233	780	73 505
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 344	2 317	2 124	956	507	113	68	86	10 515
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	16.0	14.1	15.4	13.9	9.3	6.2	5.5	11.0	14.3
<i>2009-10</i>										
Overnight separations from psychiatric acute inpatient services	no.	26 447	16 552	13 928	7 321	5 503	1 758	1 184	742	73 435
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 094	2 300	2 106	1 009	455	196	51	75	10 286

TABLE 12A.57

Table 12A.57 **Readmissions to hospital within 28 days of discharge (a), (b), (c), (d)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (b)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	15.5	13.9	15.1	13.8	8.3	11.1	4.3	10.1	14.0
<i>2010-11</i>										
Overnight separations from psychiatric acute inpatient services	no.	27 083	17 156	14 457	8 403	5 825	1 730	1 185	771	76 610
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 274	2 427	2 207	1 187	523	242	63	105	11 028
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	15.8	14.1	15.3	14.1	9.0	14.0	5.3	13.6	14.4
<i>2011-12</i>										
Overnight separations from psychiatric acute inpatient services	no.	27 463	17 910	15 192	8 719	5 987	1 655	1 306	781	79 013
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 298	2 554	2 294	1 218	551	191	165	88	11 359
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	15.7	14.3	15.1	14.0	9.2	11.5	12.6	11.3	14.4

TABLE 12A.57

Table 12A.57 Readmissions to hospital within 28 days of discharge (a), (b), (c), (d)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (b)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>2012-13</i>										
Overnight separations from psychiatric acute inpatient services	no.	28 157	18 912	15 916	9 638	5 437	1 667	1 307	889	81 923
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 141	2 771	2 275	1 317	420	212	188	95	11 419
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	14.7	14.7	14.3	13.7	7.7	12.7	14.4	10.7	13.9
<i>2013-14</i>										
Overnight separations from psychiatric acute inpatient services	no.	29 204	19 281	16 401	10 095	5 101	1 856	1 238	965	84 141
Overnight acute separations that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	no.	4 170	2 842	2 196	1 444	354	251	133	105	11 495
Proportion of overnight separations from psychiatric acute inpatient services that were followed by a readmission to a psychiatric acute inpatient service within 28 days of discharge	%	14.3	14.7	13.4	14.3	6.9	13.5	10.7	10.9	13.7

(a) Data are based on all 'in scope' separations from State and Territory psychiatric inpatient units, defined as those for which it is meaningful to examine readmission rates. The following separations were excluded: same day separations; overnight separations that occur through discharge/transfer to another hospital; statistical discharge — type change; left against medical advice/discharge at own risk and death.

Table 12A.57 **Readmissions to hospital within 28 days of discharge (a), (b), (c), (d)**

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA (b)</i>	<i>Tas (b)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
(b)	For the purposes of this indicator, a readmission for any of the separations identified as 'in-scope' is defined as an admission to any other public psychiatric acute unit within the jurisdiction that occurs within 28 days of the date of the original separation. For this to occur a system of unique client identifiers needs to be in place that allows individuals to be 'tracked' across units. Such systems have been available in all states/territories for the full period (2005-06 to 2013-14), with the exception of Tasmania (which introduced such a system in 2012-13) and SA (which has not yet introduced such a system). Undercounting of readmissions may have occurred in SA and Tasmania in the years that the system of unique identifiers is not in place. For SA therefore only readmissions to same hospital are identified, in all years' data, rather than readmissions to any hospital.									
(c)	No distinction is made between planned and unplanned readmissions because data collection systems in most Australian mental health services do not include a reliable and consistent method to distinguish a planned from an unplanned admission to hospital.									
(d)	For data before 2012-13, states and territories differed in the overnight separations that they count as 'in scope'. NSW and Queensland excluded separations where length of stay is one night only and the procedure code for ECT is recorded and the ACT excluded all overnight separations with the procedure code for ECT, whereas the others (Victoria, WA, SA, Tasmania and the NT) include all overnight separations for the procedure code for ECT. For 2012-13, the exclusion of overnight stays of one night with an ECT procedure code became a business rule for the calculation of data for this indicator. The change was considered likely to be minimal, therefore, historical data updates were not considered mandatory. The change is also unlikely to alter the interpretability of long term data trends.									
(e)	For Qld, inpatient identifiers are unique at a hospital level. A routine linkage program is utilised to create a unique identifier for reporting purposes.									

Source: AIHW (unpublished), from data provided by State and Territory governments.

Table 12A.58 Readmissions to hospital within 28 days of discharge, by selected characteristics, 2013-14 (a), (b), (c), (d)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Age group</i>										
Less than 15 years	%	12.3	19.7	15.2	11.8	3.5	–	–	–	15.0
15–24 years	%	13.9	20.1	14.0	15.7	9.4	15.1	13.5	6.8	15.3
25–34 years	%	15.9	15.1	14.2	16.0	5.2	15.3	12.7	12.4	14.7
35–44 years	%	14.5	13.3	13.4	14.2	7.2	14.0	10.5	12.2	13.5
45–54 years	%	14.0	13.3	13.0	15.4	6.8	12.2	6.1	13.6	13.2
55–64 years	%	12.9	13.7	12.8	11.2	7.2	11.0	11.7	7.1	12.3
65–74 years	%	13.1	9.1	9.1	9.6	5.9	7.7	10.8	10.0	10.0
75 years or over	%	9.1	4.3	7.2	6.1	6.7	7.1	1.7	–	6.5
<i>Gender</i>										
Male	%	13.7	13.8	12.6	12.7	6.2	13.1	8.1	12.3	12.8
Female	%	14.9	15.7	14.2	15.8	7.8	14.0	13.2	8.7	14.5
<i>SEIFA quintiles (d)</i>										
Quintile 1 (most disadvantaged)	%	13.4	14.7	13.4	13.4	6.2	13.9	–	11.7	13.2
Quintile 2	%	14.0	13.2	12.9	15.3	7.3	13.2	25.6	5.7	13.2
Quintile 3	%	13.6	16.6	12.9	14.0	5.0	15.9	6.7	8.5	14.0
Quintile 4	%	14.1	14.3	12.8	15.5	7.6	11.9	13.2	7.9	13.5
Quintile 5 (least disadvantaged)	%	14.5	13.7	14.3	12.1	11.2	4.8	9.2	5.9	13.6
<i>Indigenous status</i>										
Aboriginal and Torres Strait Islander	%	17.7	21.9	14.3	18.6	9.3	13.3	6.3	12.7	16.4
Non-Indigenous	%	14.0	14.7	13.4	13.9	7.2	13.7	11.1	9.4	13.6
<i>Remoteness (d)</i>										
Major cities	%	13.9	15.4	13.7	14.3	7.5	7.2	11.7	–	13.8
Inner regional	%	14.3	13.0	12.5	12.3	3.8	13.5	0.2	–	13.1
Outer regional	%	11.9	12.5	12.6	16.2	5.5	14.8	–	11.5	12.5
Remote	%	15.2	15.4	2.0	12.4	3.3	6.0	na	9.2	9.7
Very remote	%	3.8	12.5	9.6	7.8	7.8	15.4	na	12.6	10.2

(a) Data are based on all 'in scope' separations from State and Territory psychiatric inpatient units, defined as those for which it is meaningful to examine readmission rates. The following separations were excluded: same day separations; overnight separations that occur through discharge/transfer to another hospital; statistical discharge — type change; left against medical advice/discharge at own risk and death.

(b) For the purposes of this indicator, a readmission for any of the separations identified as 'in-scope' is defined as an admission to any other public psychiatric acute unit within the jurisdiction that occurs within 28 days of the date of the original separation. For this to occur a system of unique client identifiers needs to be in place that allows individuals to be 'tracked' across units. Such systems have been available in all states/territories for the full period (2005-06 to 2013-14), with the exception of Tasmania (which introduced such a system in 2012-13) and SA (which has not yet introduced such a system). Undercounting of readmissions may have occurred in SA and Tasmania in the years that the system of unique identifiers is not in place.

Table 12A.58 Readmissions to hospital within 28 days of discharge, by selected characteristics, 2013-14 (a), (b), (c), (d)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (e)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
(c)	No distinction is made between planned and unplanned readmissions because data collection systems in most Australian mental health services do not include a reliable and consistent method to distinguish a planned from an unplanned admission to hospital.									
(d)	Remoteness and socioeconomic status have been allocated using the client's usual residence, not the location of the service provider. State/territory is reported for the state/territory of the service provider.									
(e)	For Qld, inpatient identifiers are unique at a hospital level. A routine linkage program is utilised to create a unique identifier for reporting purposes.									

na Not available. – Nil or rounded to zero.

Source: AIHW (unpublished), from data provided by State and Territory governments.

TABLE 12A.59

Table 12A.59 **Average recurrent real costs per inpatient bed day, public hospitals, by target population (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g), (h)	WA	SA (i)	Tas (j)	ACT (j), (k)	NT (j)	Aust
General mental health services									
2005-06	795.46	686.96	691.04	939.53	790.52	735.62	732.56	1 161.64	769.29
2006-07	804.22	704.75	706.36	952.63	900.27	913.27	959.86	1 059.33	795.52
2007-08	792.98	757.59	785.56	1 004.65	935.58	941.72	1 014.19	1 246.55	827.77
2008-09	831.84	796.14	778.58	1 048.53	1 028.21	890.05	906.46	1 278.25	857.11
2009-10	836.05	810.26	823.00	1 044.19	1 012.36	1 092.54	901.57	1 300.69	872.42
2010-11	884.37	823.63	822.12	1 153.54	967.98	1 066.37	945.49	1 345.73	902.51
2011-12	937.85	837.02	856.04	1 166.49	964.89	918.43	963.75	1 611.97	934.27
2012-13	961.85	845.11	844.69	1 239.66	904.50	930.31	894.77	1 401.96	945.47
2013-14	997.57	870.35	884.02	1 245.42	1 125.90	898.57	1 063.98	1 571.39	989.55
Child and adolescent mental health services									
2005-06	1 445.77	1 470.24	1 460.88	1 299.28	1 309.31	1 437.98
2006-07	1 474.90	1 472.81	1 563.52	1 587.66	1 656.71	1 514.81
2007-08	1 483.51	1 485.09	1 648.56	1 191.33	2 222.49	1 509.91
2008-09	1 457.63	1 623.40	1 741.99	1 600.34	1 996.86	1 588.45
2009-10	1 703.51	1 586.26	1 673.39	1 598.67	2 148.99	1 671.40
2010-11	1 997.82	1 583.12	1 672.56	2 075.87	1 926.34	1 816.95
2011-12	1 816.50	1 488.27	1 690.30	2 262.84	1 909.70	1 752.31
2012-13	1 612.89	1 505.39	1 356.01	2 120.11	2 176.82	1 572.39
2013-14	1 794.50	1 405.75	1 485.45	2 251.54	2 162.64	1 687.70
Older people's mental health services									
2005-06	651.11	595.73	522.44	805.85	582.57	623.31
2006-07	672.06	625.74	571.37	792.24	618.20	..	2 662.51	..	653.60
2007-08	661.99	669.38	611.57	793.77	669.71	..	1 024.25	..	675.71
2008-09	714.14	677.16	615.97	853.27	759.06	..	1 065.36	..	718.40

Table 12A.59 **Average recurrent real costs per inpatient bed day, public hospitals, by target population (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g), (h)	WA	SA (i)	Tas (j)	ACT (j), (k)	NT (j)	Aust
2009-10	721.43	685.89	628.81	828.64	769.66	..	655.35	..	718.37
2010-11	788.24	714.94	642.48	849.16	717.40	..	654.82	..	743.67
2011-12	807.75	715.03	651.47	959.03	729.12	..	639.15	..	767.75
2012-13	884.72	736.48	619.19	1 018.39	760.42	..	771.80	..	806.22
2013-14	840.25	740.14	684.25	1 053.51	737.71	..	874.23	..	816.14
Forensic mental health services									
2005-06	610.75	875.66	912.50	1 199.31	1 000.99	546.45	..	1 086.06	825.32
2006-07	545.67	886.16	936.66	1 070.94	1 118.48	1 092.40	..	748.03	819.69
2007-08	555.44	895.81	1 060.49	1 024.76	1 153.92	1 575.81	866.04
2008-09	819.87	794.95	1 039.83	1 220.91	1 084.69	1 607.60	934.65
2009-10	885.48	929.13	1 094.58	1 138.76	1 079.30	2 047.32	990.24
2010-11	978.21	883.37	1 196.64	1 042.86	1 055.32	2 397.93	1 023.34
2011-12	959.00	834.71	1 330.64	1 224.35	1 043.93	1 478.06	1 009.92
2012-13	1 022.05	822.00	1 322.81	1 205.34	903.37	2 443.07	1 041.34
2013-14	1 089.08	871.01	1 162.26	1 322.37	949.34	2 170.86	1 072.14

- (a) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).
- (b) Depreciation is excluded for all years.
- (c) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure items.
- (d) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (e) Includes government expenditure and funded patients days in services managed and operated by private and non-government entities.
- (f) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year. For further details see the DQI for this indicator.

Table 12A.59 **Average recurrent real costs per inpatient bed day, public hospitals, by target population (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f)	Vic	Qld (g), (h)	WA	SA (i)	Tas (j)	ACT (j), (k)	NT (j)	Aust
(g)	Queensland Government has advised that it provides older people's mental health inpatient services using a number of different service models including campus and noncampus based options. All service types are reported as older people's mental health services, which may have the effect of lowering the average patient day costs compared to jurisdictions who report 'older people's care units' separately.								
(h)	Data for a small number of <i>Youth</i> services have been rolled into the General services category at the request of Queensland Government.								
(i)	For SA, any increases in admitted patient expenditure in 2013-14 partly relate to genuine increases in mental health services. However, a significant proportion of the increases relate to improved identification and allocation of direct care and general overhead expenditure to mental health services.								
(j)	Child and adolescent mental health services were not available, or could not be separately identified, in Tasmania, the ACT and the NT. Older People's Mental Health Services programs were not available, or could not be separately identified, in Tasmania and the ACT for 2005-06, and the NT. Tasmanian figures include child and adolescent mental health services within the general mental health services category. Forensic mental health services were not provided separately in the ACT and in the NT from 2007-08.								
(k)	ACT average costs for older person's mental health services are based on a new 20 bed unit opened in March 2007. During 2006-07, only 6–10 beds operated due to issues related to staffing resources. This has artificially inflated the average cost of older people's mental health services relative to other jurisdictions and other years.								
	.. Not applicable.								

Source: AIHW (unpublished) MHE NMDS.

TABLE 12A.60

Table 12A.60 **Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f), (g)	Vic (h)	Qld (i), (j), (k)	WA (l)	SA (h), (m)	Tas (h)	ACT (h), (n)	NT (h)	Aust
General mental health services									
<i>Acute</i>									
2005-06	901.03	711.69	777.13	943.08	871.93	727.83	732.56	1 161.64	837.35
2006-07	918.01	724.40	788.11	951.22	984.51	976.08	959.86	1 059.33	867.44
2007-08	897.50	789.39	903.59	996.37	987.72	935.25	1 014.19	1 246.55	902.62
2008-09	911.02	823.50	886.53	1 050.74	1 072.60	932.45	906.46	1 278.25	923.22
2009-10	923.28	837.46	896.74	1 057.30	1 053.01	1 170.13	901.57	1 300.69	936.80
2010-11	964.61	845.48	899.46	1 199.24	988.52	1 181.97	945.49	1 345.73	964.34
2011-12	1 009.77	840.69	912.13	1 191.33	930.46	980.63	963.75	1 611.97	977.31
2012-13	1 051.52	852.51	951.76	1 277.67	932.49	918.03	894.77	1 401.96	1 010.52
2013-14	1 091.72	878.53	993.99	1 287.11	1 140.01	977.63	1 063.98	1 571.39	1 060.56
<i>Non-acute</i>									
2005-06	556.26	523.97	554.90	925.92	595.06	756.64	589.72
2006-07	532.25	571.88	575.48	957.78	641.59	769.54	598.00
2007-08	550.47	558.43	592.88	1 058.84	742.93	964.24	612.43
2008-09	637.00	617.94	606.40	1 035.41	836.12	773.67	663.29
2009-10	639.14	635.32	722.38	969.30	829.60	873.67	700.44
2010-11	702.72	678.14	718.44	1 019.29	878.26	777.19	745.90
2011-12	766.02	809.50	775.07	1 078.00	1 175.53	762.97	815.32
2012-13	743.70	793.45	699.76	1 093.34	727.64	968.41	766.03
2013-14	761.42	811.05	717.74	1 069.22	1 030.31	709.22	783.29
Child and adolescent mental health services									
<i>Acute</i>									
2005-06	1 628.66	1 470.24	1 473.10	1 299.28	1 309.31	1 490.89
2006-07	1 411.58	1 472.81	1 663.24	1 529.45	1 656.71	1 510.94

TABLE 12A.60

Table 12A.60 **Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f), (g)	Vic (h)	Qld (i), (j), (k)	WA (l)	SA (h), (m)	Tas (h)	ACT (h), (n)	NT (h)	Aust
2007-08	1 530.93	1 485.09	1 657.08	1 081.58	2 222.49	1 510.67
2008-09	1 564.89	1 623.40	1 712.09	1 496.84	1 996.86	1 625.18
2009-10	1 800.92	1 586.26	1 543.29	1 383.50	2 148.99	1 650.35
2010-11	1 958.53	1 583.12	1 584.65	1 873.31	1 926.34	1 753.68
2011-12	1 879.00	1 488.27	1 537.53	2 145.62	1 909.70	1 722.92
2012-13	1 831.86	1 505.39	1 298.82	2 120.11	2 176.82	1 649.64
2013-14	2 059.03	1 405.75	1 402.23	2 044.84	2 162.64	1 735.46
<i>Non-acute</i>									
2005-06	1 128.48	..	1 410.51	1 208.61
2006-07	1 630.60	..	1 268.59	1 840.04	1 530.89
2007-08	1 380.16	..	1 621.20	2 009.41	1 506.74
2008-09	1 233.53	..	1 859.75	2 098.14	1 446.55
2009-10	1 444.69	..	2 228.17	3 115.02	1 774.05
2010-11	2 170.07	..	2 084.88	4 681.87	2 270.26
2011-12	1 633.74	..	2 520.86	4 733.92	1 915.12
2012-13	991.29	..	1 645.00	1 165.45
2013-14	1 073.19	..	2 575.22	5 968.32	1 397.37
Older people's mental health services									
<i>Acute</i>									
2005-06	695.91	595.73	755.52	835.78	748.06	698.85
2006-07	727.57	625.74	845.20	833.61	857.12	..	2 662.51	..	742.57
2007-08	743.61	669.38	928.60	819.25	870.82	..	1 024.25	..	766.72
2008-09	776.40	677.16	844.23	881.76	849.43	..	1 065.36	..	779.27
2009-10	771.35	685.89	877.47	922.44	937.01	..	655.35	..	793.75
2010-11	849.57	714.94	873.12	912.75	848.12	..	654.82	..	815.51

TABLE 12A.60

Table 12A.60 **Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f), (g)	Vic (h)	Qld (i), (j), (k)	WA (l)	SA (h), (m)	Tas (h)	ACT (h), (n)	NT (h)	Aust
2011-12	833.89	715.03	898.64	1 042.31	867.06	..	639.15	..	839.78
2012-13	922.16	736.48	889.24	1 092.29	862.67	..	771.80	..	884.48
2013-14	930.39	740.14	855.59	1 134.97	952.98	..	874.23	..	905.60
<i>Non-acute</i>									
2005-06	591.28	..	443.73	659.41	490.93	513.08
2006-07	592.87	..	474.53	653.62	491.58	525.64
2007-08	591.27	..	499.34	706.97	552.85	561.83
2008-09	653.44	..	525.61	756.21	694.71	631.95
2009-10	667.36	..	535.36	529.44	632.32	607.07
2010-11	723.89	..	555.91	468.35	586.59	628.24
2011-12	778.21	..	560.44	461.79	558.80	645.43
2012-13	840.89	..	528.25	527.23	639.46	672.11
2013-14	729.96	..	609.78	525.54	477.84	635.25
Forensic mental health services									
<i>Acute</i>									
2005-06	439.44	970.35	..	1 225.06	1 180.70	546.45	..	1 086.06	784.84
2006-07	491.61	1 058.69	..	1 083.52	1 273.66	1 092.40	..	748.03	828.36
2007-08	468.87	954.28	..	1 024.85	1 217.38	1 575.81	810.51
2008-09	742.29	866.12	..	1 220.90	1 346.42	1 607.60	923.52
2009-10	892.88	987.59	..	1 138.76	1 420.47	2 047.32	1 043.41
2010-11	1 026.59	940.59	..	1 133.87	1 418.40	2 397.93	1 093.73
2011-12	995.83	813.73	..	1 331.24	1 444.33	1 478.06	1 021.97
2012-13	1 065.44	812.08	..	1 327.44	915.84	2 443.07	1 071.40
2013-14	1 147.85	802.91	..	1 291.18	1 369.82	2 170.86	1 101.65
<i>Non-acute</i>									

TABLE 12A.60

Table 12A.60 **Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f), (g)	Vic (h)	Qld (i), (j), (k)	WA (l)	SA (h), (m)	Tas (h)	ACT (h), (n)	NT (h)	Aust
2005-06	688.89	787.91	912.50	1 172.61	954.87	841.54
2006-07	584.69	734.74	936.66	1 057.90	1 076.41	815.48
2007-08	626.69	848.61	1 060.49	1 024.62	1 136.95	895.61
2008-09	896.46	685.16	1 039.83	1 220.91	1 017.15	942.25
2009-10	878.62	832.50	1 094.58	1 138.75	994.47	943.88
2010-11	931.82	788.38	1 196.64	678.44	962.03	957.09
2011-12	929.29	867.35	1 330.64	796.39	942.81	999.57
2012-13	989.10	837.22	1 322.81	746.07	900.07	1 017.32
2013-14	1 043.74	989.30	1 162.26	1 577.39	839.76	1 048.23

- (a) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).
- (b) Depreciation is excluded for all years.
- (c) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure items.
- (d) Hospital inpatient expenditure can include expenditure on government funded public hospital services managed and operated by private and non government entities.
- (e) Mainstreaming has occurred at different rates across jurisdictions. Differences in costs can reflect differences in the rate of this institutional change (that is, the mainstreaming of mental health services).
- (f) Caution is required when interpreting NSW data. Seven residential mental health services in 2006-07 were reclassified as non-acute older person specialised hospital services in 2007-08, reflecting a change in function of those units.
- (g) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year. For further details see the DQI for this indicator.

Table 12A.60 Average recurrent costs per inpatient bed day, public hospitals, by target population and care type (2013-14 dollars) (a), (b), (c), (d), (e)

	<i>NSW (f), (g)</i>	<i>Vic (h)</i>	<i>Qld (i), (j), (k)</i>	<i>WA (l)</i>	<i>SA (h), (m)</i>	<i>Tas (h)</i>	<i>ACT (h), (n)</i>	<i>NT (h)</i>	<i>Aust</i>
(h)	Child and adolescent mental health services were not available, or could not be separately identified, in Tasmania, the ACT and the NT. Tasmanian figures include child and adolescent mental health services within the general mental health services category. Victoria and SA did not have non-acute child and adolescent mental health services units. Older People's Mental Health Services programs were not available, or could not be separately identified, in Tasmania and the NT. Older People's Mental Health Services in non-acute units were not available in Victoria and the ACT. Forensic mental health services were not provided separately in the ACT and in the NT from 2007-08.								
(i)	Queensland Government has advised that it provides older people's mental health inpatient services using a number of different service models including campus and noncampus based options. All service types are reported as older people's mental health services, which may have the effect of lowering the average patient day costs compared to jurisdictions who report 'older people's care units' separately.								
(j)	Caution is required when interpreting Queensland data. Several Forensic services reported in 2008-09 were reclassified as General services in 2009-10 to more accurately reflect the function of these services. Queensland does not report any acute forensic services; however, forensic patients can and do access acute care through general units, which may also impact on the comparability of both cost and length of stay data.								
(k)	Data for a small number of hospital beds reported by Queensland as youth specialised mental health hospital beds are included in the General category at the request of Queensland Government.								
(l)	Caution is required when interpreting WA data. A review of services resulted in the reclassification of beds between the acute and non-acute categories for the 2010-11 collection, to more accurately reflect the function of these services. In addition, during 2010-11, the child and adolescent non-acute inpatient service initiated the closure of beds in order to carry out a complete refurbishment. The service ceased operating in late 2011, but reopened in 2013-14.								
(m)	For SA, any increases in admitted patient expenditure in 2013-14 partly relate to genuine increases in mental health services. However, a significant proportion of the increases relate to improved identification and allocation of direct care and general overhead expenditure to mental health services.								
(n)	ACT average costs for older people's mental health services are based on a new 20 bed unit opened in March 2007. During 2006-07, only 6-10 beds operated due to issues related to staffing resources. This has artificially inflated the average cost of older people's mental health services relative to other jurisdictions and other years.								
	.. Not applicable.								

Source: AIHW (unpublished) MHE NMDS.

Table 12A.61 Average length of stay, public hospitals acute units, by target population (no. of days) (a), (b)

	NSW (c)	Vic	Qld (d)	WA	SA	Tas (e)	ACT (e)	NT (e)	Aust
<i>2010-11</i>									
General mental health services	14.8	14.5	11.4	14.9	13.4	12.0	15.2	12.6	13.8
Child and adolescent mental health services	21.7	10.4	11.2	8.0	4.2	11.8
Older people's mental health services	35.4	32.6	20.7	51.3	45.6	..	36.3	..	35.5
Total	16.0	16.1	11.8	17.3	15.2	12.0	17.5	12.6	15.1
<i>2011-12</i>									
General mental health services	14.6	14.4	11.6	13.8	12.2	12.6	14.5	10.7	13.5
Child and adolescent mental health services	22.1	7.3	11.5	7.2	3.1	10.5
Older people's mental health services	41.2	30.5	11.3	49.8	41.2	..	36.8	..	31.9
Total	16.0	15.4	11.6	16.0	13.8	12.6	16.9	10.7	14.6
<i>2012-13</i>									
General mental health services	13.9	13.8	10.2	15.2	15.0	12.9	15.9	11.8	13.2
Child and adolescent mental health services	21.7	7.5	11.1	6.4	3.6	10.7
Older people's mental health services	42.2	29.7	21.9	50.3	39.1	..	40.3	..	35.7
Total	15.3	14.9	10.7	17.3	16.4	12.9	18.2	11.8	14.4
<i>2013-14</i>									
General mental health services	14.0	13.4	10.0	14.3	13.6	11.3	15.3	10.6	12.9
Child and adolescent mental health services	19.8	7.9	10.3	10.2	4.7	11.1
Older people's mental health services	43.0	31.1	20.2	41.6	38.8	..	30.2	..	35.3
Total	15.4	14.7	10.3	16.5	15.3	11.3	17.0	10.6	14.1

(a) The quality of the separations data used to derive the results in this table is variable across jurisdictions. Until recently, these separations data were not subject to in depth scrutiny. It is expected that the quality of these data will improve over time.

Table 12A.61 Average length of stay, public hospitals acute units, by target population (no. of days) (a), (b)

	<i>NSW (c)</i>	<i>Vic Qld (d)</i>	<i>WA</i>	<i>SA</i>	<i>Tas (e)</i>	<i>ACT (e)</i>	<i>NT (e)</i>	<i>Aust</i>
(b) There is a mismatch between the inpatient bed days and the separations used to derive this indicator for the relevant reference periods.								
– Patients days for clients who separated in the reference period (for example, 2012-13) that were during the previous period (2011-12) are excluded.								
– Patient days for clients who remain in hospital (that is, are not included in the separations data) are included.								
The 'average length of stay' data reported here may not match data reported elsewhere (such as the Australian Institute of Health and Welfare's [AIHW's] Mental Health Services in Australia publication) due to differences in scope, for example these data include separations and days within the reference period only.								
(c) The quality of the NSW 2010-11 MHE NMDS data has been affected by the reconfiguration of the service system during the year. For further details see the DQI for this indicator.								
(d) Separations for a small number of hospital beds reported by Queensland as youth specialised mental health hospital beds were included in the general category at the request of Queensland Government.								
(e) Child and adolescent mental health services were not available, or could not be separately identified, in Tasmania, the ACT and the NT. Tasmanian figures include child and adolescent mental health services within the general mental health services category. Older People's Mental Health Services programs were not available, or could not be separately identified, in Tasmania and the NT.								
.. Not applicable.								

Source: AIHW (unpublished) MHE NMDS.

TABLE 12A.62

Table 12A.62 **Average recurrent cost per inpatient bed day, by public hospital type (2013-14 dollars) (a), (b), (c), (d), (e), (f)**

	NSW (g), (h)	Vic (i)	Qld (j)	WA (k)	SA (l)	Tas (m)	ACT (m), (n)	NT (m), (n)	Aust
Psychiatric hospitals (acute units)									
2005-06	850.40	970.35	..	944.54	974.93	907.16
2006-07	815.55	1 058.69	..	976.94	1 108.26	925.03
2007-08	763.06	954.28	..	954.19	1 201.42	901.69
2008-09	775.79	866.12	..	1 021.14	1 161.62	927.43
2009-10	1 005.37	987.59	..	1 022.75	1 204.68	1 036.14
2010-11	985.40	940.59	..	1 204.44	1 010.20	1 028.43
2011-12	1 112.31	813.73	..	1 255.50	956.52	1 075.65
2012-13	1 164.02	812.08	..	1 328.67	1 062.99	1 133.79
2013-14	1 160.95	802.91	..	1 315.17	1 296.14	1 148.88
Psychiatric hospitals (non-acute units)									
2005-06	597.41	787.91	733.06	967.45	590.83	664.63
2006-07	562.48	734.74	746.45	971.47	616.20	656.17
2007-08	576.52	848.61	821.69	956.23	691.99	693.81
2008-09	662.99	685.16	807.03	1 023.21	791.83	740.74
2009-10	653.70	832.50	828.31	1 012.65	765.52	740.27
2010-11	721.53	788.38	846.80	966.52	766.59	781.84
2011-12	766.56	867.35	934.47	1 012.27	843.00	847.56
2012-13	766.84	837.22	906.13	1 062.95	733.63	829.58
2013-14	782.94	989.30	879.07	1 123.37	723.60	837.87
Psychiatric hospitals (all units)									
2005-06	684.33	875.66	733.06	954.59	735.59	744.21
2006-07	651.53	886.16	746.45	974.52	784.67	743.17
2007-08	640.18	895.81	821.69	954.78	856.48	763.32
2008-09	693.98	794.95	807.03	1 021.73	925.35	800.88

TABLE 12A.62

Table 12A.62 **Average recurrent cost per inpatient bed day, by public hospital type (2013-14 dollars) (a), (b), (c), (d), (e), (f)**

	NSW (g), (h)	Vic (i)	Qld (j)	WA (k)	SA (l)	Tas (m)	ACT (m), (n)	NT (m), (n)	Aust
2009-10	752.16	929.13	828.31	1 019.83	912.83	834.26
2010-11	798.26	883.37	846.80	1 090.05	853.75	856.44
2011-12	880.49	834.71	934.47	1 152.53	888.91	924.19
2012-13	902.73	822.00	906.13	1 217.33	857.56	932.37
2013-14	917.63	871.01	879.07	1 238.53	906.59	945.08
General acute hospital with a psychiatric unit or ward (acute units)									
2006-07	912.38	731.94	834.79	934.06	905.14	1 002.96	1 036.08	1 056.13	857.00
2007-08	903.82	789.51	942.83	975.52	889.94	1 062.02	1 016.12	1 246.55	897.84
2008-09	933.95	822.42	926.19	1 044.11	989.64	1 067.18	934.29	1 278.25	926.21
2009-10	918.90	837.29	928.99	1 054.62	998.41	1 334.55	849.91	1 300.69	932.11
2010-11	989.51	849.76	934.34	1 133.59	973.82	1 380.98	879.67	1 345.73	969.58
2011-12	1 005.50	843.52	945.63	1 171.07	937.67	1 090.92	888.43	1 611.97	974.05
2012-13	1 054.15	859.11	971.51	1 242.34	895.86	1 152.50	869.51	1 401.96	1 005.89
2013-14	1 111.51	874.72	1 014.41	1 267.47	1 081.86	1 181.06	1 025.44	1 571.39	1 057.94
General acute hospital with a psychiatric unit or ward (non-acute units)									
2006-07	685.91	571.88	548.16	792.30	..	769.54	601.68
2007-08	632.91	558.43	563.70	1 035.43	..	964.24	622.18
2008-09	800.01	617.94	580.82	992.74	..	773.67	678.77
2009-10	875.27	635.32	634.08	770.39	..	873.67	717.38
2010-11	932.04	678.14	632.43	933.86	..	777.19	739.92
2011-12	927.49	809.50	665.70	942.07	..	762.97	802.04
2012-13	899.75	793.45	586.12	796.83	..	968.41	753.30
2013-14	881.22	811.05	666.56	853.44	..	709.22	787.84
General acute hospital with a psychiatric unit or ward (all units)									
2006-07	896.28	716.31	741.38	919.68	905.14	944.25	1 036.08	1 056.13	820.99

TABLE 12A.62

Table 12A.62 **Average recurrent cost per inpatient bed day, by public hospital type (2013-14 dollars) (a), (b), (c), (d), (e), (f)**

	NSW (g), (h)	Vic (i)	Qld (j)	WA (k)	SA (l)	Tas (m)	ACT (m), (n)	NT (m), (n)	Aust
2007-08	865.61	765.28	821.52	981.05	889.94	1 043.70	1 016.12	1 246.55	853.92
2008-09	919.89	801.77	820.28	1 038.60	989.64	1 000.91	934.29	1 278.25	890.37
2009-10	914.08	816.56	840.94	1 021.32	998.41	1 231.48	849.91	1 300.69	900.92
2010-11	983.32	832.49	842.36	1 117.11	973.82	1 229.69	879.67	1 345.73	936.90
2011-12	991.78	840.45	861.23	1 152.59	937.67	1 013.05	888.43	1 611.97	946.14
2012-13	1 025.82	852.82	844.89	1 209.44	895.86	1 113.04	869.51	1 401.96	962.77
2013-14	1 067.66	868.85	915.50	1 235.40	1 081.86	1 059.69	1 025.44	1 571.39	1 014.11

- (a) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).
- (b) Depreciation is excluded for all years.
- (c) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure items.
- (d) Due to the ongoing validation of NMDS, data could differ from previous reports.
- (e) Hospital inpatient expenditure can include expenditure on government funded public hospital services managed and operated by private and non-government entities.
- (f) Mainstreaming has occurred at different rates across jurisdictions. The client profile and service costs can be very different for those of a jurisdiction in which general psychiatric treatment still occurs mostly in psychiatric hospitals. For example, Victorian data for psychiatric hospitals comprise mainly forensic services as nearly all general psychiatric treatment occurs in mainstreamed units in general acute hospitals.
- (g) Caution is required when interpreting NSW data. Seven residential mental health services in 2006-07 were reclassified as non-acute older person specialised hospital services in 2007-08, reflecting a change in function of those units.
- (h) The quality of the NSW 2010-11 MHE NMDS data used for this Report has been affected by the reconfiguration of the service system during the year. For further details see the DQI for this indicator.
- (i) Mainstreaming has occurred at different rates in different jurisdictions. In Victoria's case, the data for psychiatric hospitals comprises mainly forensic services, since nearly all general psychiatric treatment occurs in mainstreamed units in general acute hospitals. This means that the client profile and service costs are very different from those of a jurisdiction where general psychiatric treatment still occurs mostly in psychiatric hospitals.

Table 12A.62 **Average recurrent cost per inpatient bed day, by public hospital type (2013-14 dollars) (a), (b), (c), (d), (e), (f)**

	NSW (g), (h)	Vic (i)	Qld (j)	WA (k)	SA (l)	Tas (m)	ACT (m), (n)	NT (m), (n)	Aust
(j)	Queensland data for public acute hospitals include costs associated with extended treatment services (campus-based and non-campus-based) that report through general acute hospitals. Queensland does not provide acute services in psychiatric hospitals.								
(k)	Caution is required when interpreting WA data. A review of services resulted in the reclassification of beds between the acute and non-acute categories for the 2010-11 collection, to more accurately reflect the function of these services.								
(l)	For SA, any increases in admitted patient expenditure in 2013-14 partly relate to genuine increases in mental health services. However, a significant proportion of the increases relate to improved identification and allocation of direct care and general overhead expenditure to mental health services.								
(m)	Tasmania, the ACT and the NT do not have public psychiatric hospitals.								
(n)	SA, the ACT and the NT do not have non-acute units in public acute hospitals with a psychiatric unit or ward.								
	.. Not applicable.								

Source: AIHW (unpublished) MHE NMDS.

TABLE 12A.63

Table 12A.63 **Average recurrent cost per patient day for community residential services (2013-14 dollars) (a), (b), (c), (d), (e)**

	<i>NSW</i> (f), (g), (h)	<i>Vic</i> (h)	<i>Qld</i> (i)	<i>WA</i> (j), (k), (h)	<i>SA</i> (k), (l)	<i>Tas</i> (m), (n)	<i>ACT</i> (h), (n)	<i>NT</i> (k), (l), (o)	<i>Aust</i>
General adult units									
2005-06									
24-hour staffed units	308.06	494.69	..	377.59	280.99	457.07	549.40	..	444.55
non-24-hour staffed units	100.38	174.01	..	139.94	327.25	360.11	87.35	132.82	157.27
2006-07									
24-hour staffed units	304.48	480.35	..	491.26	278.22	476.46	605.20	..	448.29
non-24-hour staffed units	96.74	158.26	..	157.59	333.87	239.14	121.14	306.95	150.44
2007-08									
24-hour staffed units	281.92	465.96	..	525.76	435.81	546.39	622.21	308.36	451.79
non-24-hour staffed units	185.71	155.11	..	194.75	534.98	241.78	115.17	..	174.06
2008-09									
24-hour staffed units	318.46	498.89	..	446.71	439.05	581.49	756.28	280.23	480.34
non-24-hour staffed units	240.43	154.53	..	177.59	324.11	272.59	109.67	..	178.77
2009-10									
24-hour staffed units	273.60	531.13	..	347.96	424.49	447.92	721.97	362.16	470.64
non-24-hour staffed units	206.93	160.27	..	162.85	287.14	247.37	124.50	..	175.32
2010-11									
24-hour staffed units	314.46	567.42	..	555.94	503.23	505.86	692.03	381.32	529.83
non-24-hour staffed units	193.42	165.83	..	146.06	292.71	237.19	117.19	..	169.47
2011-12									
24-hour staffed units	285.81	513.30	..	390.57	520.95	515.94	688.11	325.22	479.93
non-24-hour staffed units	184.63	166.36	..	156.52	356.32	208.91	140.78	..	171.52
2012-13									
24-hour staffed units	260.19	522.56	..	421.03	472.72	657.80	692.47	394.12	487.65
non-24-hour staffed units	135.17	160.26	..	176.45	235.99	254.48	123.42	..	172.64

TABLE 12A.63

Table 12A.63 **Average recurrent cost per patient day for community residential services (2013-14 dollars) (a), (b), (c), (d), (e)**

	<i>NSW</i> (f), (g), (h)	<i>Vic</i> (h)	<i>Qld</i> (i)	<i>WA</i> (j), (k), (h)	<i>SA</i> (k), (l)	<i>Tas</i> (m), (n)	<i>ACT</i> (h), (n)	<i>NT</i> (k), (l), (o)	<i>Aust</i>
2013-14									
24-hour staffed units	249.60	587.88	..	545.56	511.06	516.96	668.45	442.30	535.58
non-24-hour staffed units	130.74	162.30	..	181.78	239.15	228.64	95.98	..	170.38
Older people's care units									
2005-06									
24-hour staffed units	339.28	357.86	506.17	177.40	..	360.11
non-24-hour staffed units	127.15	127.15
2006-07									
24-hour staffed units	400.93	333.42	533.68	186.23	..	351.08
non-24-hour staffed units	329.86	329.86
2007-08									
24-hour staffed units	222.10	331.38	838.30	194.35	..	345.52
non-24-hour staffed units	177.27	177.27
2008-09									
24-hour staffed units	197.90	360.85	556.14	258.07	..	366.91
non-24-hour staffed units	235.22	235.22
2009-10									
24-hour staffed units	229.61	354.57	796.57	208.94	..	369.12
non-24-hour staffed units	227.73	227.73
2010-11									
24-hour staffed units	243.77	367.38	739.48	223.39	..	379.17
non-24-hour staffed units	299.28	299.28
2011-12									
24-hour staffed units	246.78	364.55	718.59	263.20	..	376.71

TABLE 12A.63

Table 12A.63 **Average recurrent cost per patient day for community residential services (2013-14 dollars) (a), (b), (c), (d), (e)**

	NSW (f), (g), (h)	Vic (h)	Qld (i)	WA (j), (k), (h)	SA (k), (l)	Tas (m), (n)	ACT (h), (n)	NT (k), (l), (o)	Aust
non-24-hour staffed units
2012-13									
24-hour staffed units	243.01	376.33	849.99	264.38	..	391.98
non-24-hour staffed units
2013-14									
24-hour staffed units	271.47	387.81	634.86	196.37	..	397.86
non-24-hour staffed units

(a) Depreciation is excluded for all years.

(b) Unit costs are not casemix adjusted.

(c) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).

(d) See AIHW *Mental Health Services in Australia* on-line publication (<http://mhsa.aihw.gov.au/resources/expenditure/data-source/>) for a full description of the derivation of expenditure items.

(e) Due to the ongoing validation of the NMDS, data could differ from previous reports.

(f) Caution is required when interpreting NSW data. Seven residential mental health services in 2006–07 were reclassified as non-acute older person specialised hospital services in 2007–08, reflecting a change in function of those units.

(g) The quality of the NSW 2010-11 *MHE NMDS* data used for this Report has been affected by the reconfiguration of the service system during the year. For further details see the DQI for this indicator.

(h) A small number of residential beds reported by NSW and the ACT as child and adolescent residential mental health service beds were included in the general category at the request of these jurisdictions. Expenditure for a small number of residential beds reported by Victoria, WA and the ACT as youth specialised mental health residential beds were included in the general category at the request of these jurisdictions.

(i) Queensland does not fund community residential services, however, it funds a number of extended treatment services, both campus and non-campus based, which provide longer term inpatient treatment and rehabilitation services with a full clinical staffing 24 hours a day 7 days a week. Queensland does not report these beds as community residential beds as it considers these beds to be substantially different to beds described as such in other states and territories.

Table 12A.63 **Average recurrent cost per patient day for community residential services (2013-14 dollars) (a), (b), (c), (d), (e)**

	<i>NSW</i> (f), (g), (h)	<i>Vic</i> (h)	<i>Qld</i> (i)	<i>WA</i> (j), (k), (h)	<i>SA</i> (k), (l)	<i>Tas</i> (m), (n)	<i>ACT</i> (h), (n)	<i>NT</i> (k), (l), (o)	<i>Aust</i>
(j)	Caution is required when interpreting WA data. Several residential services reported as 24-hour staffed services in 2009-10 transitioned to a non-24-hour staffed model of care as of 1 July 2010.								
(k)	WA, SA and the NT do not have any community residential services that are aged care units.								
(l)	A small number of residential services reported by SA in 2013-14, and the NT in 2012-13 and 2013-14, as forensic are included in the general category at the request of those jurisdictions.								
(m)	Tasmanian services include both acute and rehabilitation units which have higher unit costs than extended care units.								
(n)	Tasmania and the ACT do not have any community-based residential services that are non-24 hour staffed older people's units. From 2011-12, NSW did not have non-24 hour staffed older people's units.								
(o)	General adult 24-hour residential services were not provided in the NT until 2007-08. From 2007-08, general non-24-hour staffed units are not provided.								
	.. Not applicable.								

Source: AIHW (unpublished) MHE NMDS.

Table 12A.64 Average cost, and treatment days per episode, of ambulatory care (a), (b), (c)

	NSW (d)	Vic (e)	Qld	WA	SA (f)	Tas (g)	ACT	NT	Aust
<i>Average treatment days per episode of ambulatory care</i>									
2005-06	6.7	7.8	4.9	4.5	4.8	4.7	8.2	4.0	6.0
2006-07	6.8	7.7	5.2	4.5	5.0	4.1	8.0	4.0	6.1
2007-08	8.0	7.7	5.4	4.6	5.2	5.9	8.0	3.9	6.5
2008-09	7.2	7.6	4.5	4.8	5.3	6.0	8.0	3.9	6.1
2009-10	7.6	7.6	4.9	4.9	5.3	5.2	8.2	3.5	6.3
2010-11	7.5	7.7	5.2	5.0	5.5	5.5	8.2	3.6	6.4
2011-12	8.0	na	5.8	5.0	5.4	4.5	8.6	3.6	6.4
2012-13	7.8	na	6.4	4.8	5.4	3.9	8.4	4.0	6.5
2013-14	8.2	7.1	6.5	4.9	5.4	6.0	8.4	4.4	6.8
<i>Average cost per treatment day of ambulatory care (2013-14 \$) (h)</i>									
2005-06	262.42	278.74	307.58	463.44	443.65	670.22	281.65	483.17	314.12
2006-07	274.14	282.04	341.69	457.60	384.10	591.05	307.87	555.22	324.24
2007-08	265.03	302.79	366.84	487.15	344.45	456.94	292.77	614.11	328.18
2008-09	277.36	315.25	456.96	473.36	332.52	424.79	303.10	567.20	347.58
2009-10	263.97	318.48	524.20	445.49	338.80	412.17	267.11	618.99	348.87
2010-11	273.01	342.42	482.37	435.62	357.62	352.63	266.02	625.06	353.06
2011-12	259.84	na	449.59	457.04	348.39	491.98	263.31	573.22	346.36
2012-13	232.50	na	373.55	448.12	344.52	682.88	241.56	447.30	313.97
2013-14	218.17	366.92	344.03	426.38	345.34	317.71	217.39	431.66	304.47

- (a) Non-uniquely identifiable consumers' have been excluded from the episodes of ambulatory care and treatment days data.
- (b) Recurrent expenditure data used to derive this measure have been adjusted (that is, reduced) to account for proportion of clients in the *CMHC NMDS* that were defined as 'non-uniquely identifiable consumers'. Therefore, it does not match recurrent expenditure on ambulatory care reported elsewhere.
- (c) Due to the ongoing validation of *NMDS*, data could differ from previous reports.
- (d) The quality of the NSW 2010-11 MHE *NMDS* data used for this Report has been affected by the reconfiguration of the service system during the year. For further details see the *DQI* for this indicator.
- (e) Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. The total only includes those jurisdictions that have provided data.
- (f) For SA, any increases in ambulatory expenditure in 2013-14 partly relate to genuine increases in mental health services. However, a significant proportion of the increases relate to improved identification and allocation of direct care and general overhead expenditure to mental health services.
- (g) Industrial action in Tasmania has limited the available data quality and quantity of the 2011-12 and 2012-13 data.
- (h) Time series financial data are adjusted to 2013-14 dollars using the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services (table 12A.98).

na Not available.

Source: AIHW (unpublished) *CMHC NMDS* and *MHE NMDS*.

Table 12A.65 Risk status recent drinkers (in last 12 months) aged 14 years or over, 2013 (per cent)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Lifetime status</i>									
Abstainers (a)	24.3	23.5	19.9	18.3	20.6	16.9	17.5	16.7	22.0
Low risk (b)	58.9	60.4	60.0	60.0	60.9	64.4	60.5	53.6	59.9
Risky (c)	16.7	16.1	20.2	21.6	18.5	18.6	22.0	29.7	18.2
<i>Single occasion</i>									
Abstainers (a)	24.3	23.5	19.9	18.3	20.6	16.9	17.5	16.7	22.0
Low risk (d)	41.2	40.8	39.5	38.2	39.5	42.4	38.3	31.4	40.2
Risky									
At least yearly (e)	10.7	10.8	12.4	12.1	12.1	11.2	15.0	12.2	11.4
At least monthly (f)	11.3	11.8	12.8	13.6	12.9	14.3	13.1	14.0	12.2
At least weekly (g)	12.4	13.1	15.4	17.8	14.9	15.2	16.1	25.7	14.2
Total risky	34.5	35.7	40.6	43.5	39.9	40.7	44.2	51.9	37.8

(a) Not consumed alcohol in the previous 12 months.

(b) On average, had no more than 2 standard drinks per day.

(c) On average, had more than 2 standard drinks per day.

(d) Never had more than 4 standard drinks on any occasion.

(e) Had more than 4 standard drinks at least once a year, but not as often as monthly.

(f) Had more than 4 standard drinks at least once a month, but not as often as weekly.

(g) Had more than 4 standard drinks at least once a week.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.66

Table 12A.66 **Recent drinkers lifetime and single occasion risk, people aged 14 years or older, by social characteristics, 2013 (per cent)**

	<i>Abstainers (a)</i>	<i>Lifetime risk</i>		<i>Single occasion risk</i>		
		<i>Low risk (b)</i>	<i>Risky (c)</i>	<i>Low risk (d)</i>	<i>At least yearly (e)</i>	<i>At least weekly (f)</i>
<i>Socioeconomic status</i>						
Quintile 1 (lowest)	31.1	53.0	15.9	36.3	19.5	13.0
Quintile 2	24.4	57.6	18.0	39.6	21.5	14.6
Quintile 3	20.9	59.9	19.2	39.9	24.3	15.0
Quintile 4	18.6	62.4	19.0	42.3	24.4	14.7
Quintile 5 (highest)	16.7	65.0	18.4	42.1	27.6	13.6
<i>Geography</i>						
Major cities	23.1	60.2	16.7	40.4	23.5	13.0
Inner regional	18.9	62.0	19.1	41.8	24.4	14.9
Outer regional	20.5	56.9	22.6	38.1	23.6	17.8
Remote/Very remote	17.5	47.6	34.9	30.8	22.8	28.9
<i>Indigenous status</i>						
Aboriginal and/or Torres Strait Islander	27.9	49.4	22.7	22.4	29.8	19.9
Non-Indigenous	21.7	60.2	18.1	40.6	23.5	14.1

(a) Not consumed alcohol in the previous 12 months.

(b) On average, had no more than 2 standard drinks per day.

(c) On average, had more than 2 standard drinks per day.

(d) Never had more than 4 standard drinks on any occasion.

(e) Had more than 4 standard drinks at least once a year but not as often as weekly.

(f) Had more than 4 standard drinks at least once a week.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

Table 12A.67 Recent alcohol and illicit drug use, people aged 14 years or over, by substance, 2013 (per cent) (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Alcohol	75.8	76.7	80.4	81.8	79.5	83.2	82.6	83.6	78.2
Illicit drugs									
Cannabis	9.5	9.1	11.1	11.3	11.0	11.8	10.1	17.1	10.2
Ecstasy	2.4	2.4	2.4	2.6	2.8	*2.9	2.9	3.7	2.5
Meth/amphetamines (c)	1.4	1.9	2.3	3.8	2.2	*3.0	2.2	*2.8	2.1
Cocaine	2.7	2.0	2.0	1.6	*1.2	**1.2	2.8	*2.4	2.1
Hallucinogens	1.0	1.3	1.2	1.9	*1.6	*1.1	*1.7	*1.8	1.3
Inhalants	0.8	0.9	0.8	*0.5	*0.4	*1.7	*1.1	*0.8	0.8
Heroin	*<0.1	*0.1	**<0.1	*0.3	**<0.1	–	**0.3	**<0.1	0.1
Ketamine	*0.3	*0.3	**0.2	–	**0.3	*0.8	**0.2	**0.4	0.3
GHB	*<0.1	**<0.1	**<0.1	**0.1	–	**0.7	–	**<0.1	*<0.1
Synthetic Cannabinoids	1.0	1.0	1.5	*2.5	*0.9	*0.9	*0.8	2.8	1.2
New and Emerging Psychoactive Substances	*0.2	*0.5	*0.5	*0.5	*0.4	**1.1	**0.5	*0.6	0.4
Injected drugs	*0.3	*0.2	*0.3	*0.6	*0.3	*0.9	**0.2	*0.3	0.3
<i>Any illicit (d)</i>	11.4	11.0	12.6	13.7	12.5	13.3	12.4	19.0	12.0

- (a) Recent means used in the previous 12 months. For alcohol 'recent use' includes daily, weekly and less than weekly drinkers.
- (b) Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with relative standard errors greater than 50 per cent should be considered as unreliable for most practical purposes. Estimates that have RSEs greater than 50 per cent are marked with " ** " and those with RSEs of between 25 per cent and 50 per cent are marked with " * ".
- (c) Use for non-medical purposes.
- (d) Illicit use of at least 1 of 12 drugs (excluding pharmaceuticals) in the previous 12 months in 2013.
– Nil or rounded to zero.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

Table 12A.68 Use of cannabis, by age group, 2013 (per cent) (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
14–19 years	14.6	12.5	14.7	19.1	*12.3	*19.2	*22.2	24.9	14.8
20–29 years	20.8	20.4	21.6	18.3	22.3	26.7	19.3	23.9	20.8
30–39 years	10.7	10.6	14.0	15.9	13.8	16.7	8.7	18.3	12.3
40–49 years	8.3	7.2	12.0	10.3	13.8	*10.1	8.6	13.1	9.5
50–59 years	6.6	7.0	8.1	6.9	9.3	9.1	*4.1	12.7	7.3
60 years or over	*1.6	*0.7	*0.7	*1.7	*0.9	*1.5	**1.4	*7.0	1.2
14 years or over	9.5	9.1	11.1	11.3	11.0	11.8	10.1	17.1	10.2

(a) Recent use means used in the previous 12 months.

(b) Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with relative standard errors greater than 50 per cent should be considered as unreliable for most practical purposes. Estimates that have RSEs greater than 50 per cent are marked with " ** " and those with RSEs of between 25 per cent and 50 per cent are marked with " * ".

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

Table 12A.69 Risk status recent drinkers (in last 12 months) aged 14 years or over, 2010 (per cent)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
<i>Lifetime status</i>									
Abstainers (a)	22.2	21.3	17.1	17.3	19.3	14.6	13.7	13.9	19.9
Low risk (b)	58.8	59.9	59.1	59.6	60.9	65.7	66.5	56.2	59.6
Risky (c)	19.0	18.8	23.7	23.0	19.7	19.7	19.8	29.8	20.5
<i>Single occasion</i>									
Abstainers (a)	22.2	21.3	17.1	17.3	19.3	14.6	13.7	13.9	19.9
Low risk (d)	40.5	39.9	37.0	38.6	41.6	44.3	41.1	34.5	39.6
Risky									
At least yearly (e)	10.8	11.6	12.0	12.5	11.2	11.7	15.3	11.4	11.6
At least monthly (f)	11.2	12.5	15.3	13.4	11.4	13.3	16.1	15.1	12.8
At least weekly (g)	15.3	14.6	18.5	18.2	16.5	16.0	13.8	25.1	16.2
Total risky	37.3	38.8	45.9	44.0	39.1	41.0	45.2	51.6	40.6

(a) Not consumed alcohol in the previous 12 months.

(b) On average, had no more than 2 standard drinks per day.

(c) On average, had more than 2 standard drinks per day.

(d) Never had more than 4 standard drinks on any occasion.

(e) Had more than 4 standard drinks at least once a year, but not as often as monthly.

(f) Had more than 4 standard drinks at least once a month, but not as often as weekly.

(g) Had more than 4 standard drinks at least once a week.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.70

Table 12A.70 Recent alcohol and illicit drug use, people aged 14 years or over, by substance, 2010 (per cent) (a), (b)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Alcohol	78.2	79.1	83.2	83.0	81.0	85.6	86.5	86.3	80.5
Illicit drugs									
Cannabis	9.3	9.4	11.0	13.4	11.3	8.6	9.5	16.5	10.3
Ecstasy	2.9	3.1	2.7	3.7	3.3	*1.7	*2.3	3.2	3.0
Meth/amphetamines (c)	1.6	2.3	1.9	3.4	2.5	*1.1	*1.2	*2.1	2.1
Cocaine	2.7	2.3	1.3	2.2	1.7	*0.8	*1.8	**0.5	2.1
Hallucinogens	0.8	1.8	1.4	1.9	1.0	*1.0	*1.5	*2.6	1.4
Inhalants	0.6	0.6	0.6	*0.4	*0.6	*0.8	**0.6	*1.5	0.6
Heroin	*0.2	*0.3	*0.1	*0.3	*0.2	**0.1	**0.3	**0.1	0.2
GHB	*0.2	*0.2	*0.1	**0.1	**0.1	–	**<0.1	–	0.1
<i>Any illicit (d)</i>	11.4	11.0	12.3	15.4	12.7	9.6	11.4	18.8	12.0

(a) Recent means used in the previous 12 months. For alcohol 'recent use' includes daily, weekly and less than weekly drinkers.

(b) Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with relative standard errors greater than 50 per cent should be considered as unreliable for most practical purposes. Estimates that have RSEs greater than 50 per cent are marked with " ** " and those with RSEs of between 25 per cent and 50 per cent are marked with " * ".

(c) Use for non-medical purposes.

(d) Includes ketamine and injected drugs, but excludes pharmaceuticals.

– Nil or rounded to zero.

Source: AIHW (2011) *2010 National Drug Strategy Household Survey Report*, Drug statistics series no. 25, Cat. no. PHE 145, Canberra.

TABLE 12A.71

Table 12A.71 **Lifetime risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent)**

	<i>Abstainers (a)</i>			<i>Low risk (b)</i>			<i>Risky (c)</i>		
	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>
<i>Males</i>									
14–19 years	30.4	36.9	47.3	51.7	43.6	43.1	17.8	19.5	9.6
20–29 years	11.2	14.1	14.2	49.5	49.1	54.5	39.2	36.8	31.2
18–24 years	12.1	14.9	16.5	49.2	46.0	55.9	38.6	39.1	27.6
25–34 years	10.9	13.6	14.5	55.5	53.1	55.1	33.5	33.3	30.4
35–44 years	10.9	12.3	14.1	58.0	55.6	55.7	31.0	32.1	30.2
45–54 years	10.0	12.1	15.6	58.5	55.5	56.5	31.4	32.3	27.8
55–64 years	12.0	14.2	15.3	58.1	57.2	55.2	29.7	28.6	29.6
65–74 years	16.9	17.1	19.1	57.2	57.9	54.9	25.8	25.0	26.0
75+ years	22.8	23.7	25.9	59.2	59.6	58.6	17.9	16.7	15.5
Total	14.2	16.7	19.1	56.0	53.8	54.5	29.7	29.6	26.5
<i>Females</i>									
14–19 years	29.3	36.6	46.1	59.3	52.5	47.5	11.2	10.9	6.4
20–29 years	14.9	15.9	19.4	67.5	66.4	68.4	17.5	17.7	12.2
18–24 years	14.5	14.1	18.0	65.2	63.8	67.5	20.2	22.1	14.6
25–34 years	13.8	18.5	21.3	72.8	69.0	69.0	13.2	12.4	9.7
35–44 years	13.5	16.3	18.3	73.3	71.9	70.5	13.1	11.9	11.2
45–54 years	15.6	18.9	18.1	71.7	68.4	68.6	12.5	12.7	13.2
55–64 years	22.2	22.6	20.4	67.9	67.5	69.0	9.8	10.0	10.5
65–74 years	33.9	31.0	29.6	59.1	62.8	63.1	6.9	6.2	7.3
75+ years	40.2	41.8	44.8	55.3	53.4	52.4	4.4	4.7	2.8
Total	20.5	23.0	24.8	67.5	65.4	65.2	11.8	11.6	10.0
<i>All people</i>									
14–19 years	29.9	36.7	46.7	55.4	48.0	45.2	14.5	15.3	8.1
20–29 years	13.0	15.0	16.8	58.4	57.6	61.3	28.5	27.4	21.9
18–24 years	13.3	14.6	17.2	56.9	54.4	61.5	29.7	31.0	21.3
25–34 years	12.4	16.1	17.9	64.3	61.1	62.1	23.2	22.8	20.0
35–44 years	12.2	14.3	16.2	65.7	63.9	63.2	22.0	21.7	20.6
45–54 years	12.7	15.5	16.9	65.0	62.0	62.7	22.1	22.5	20.4
55–64 years	17.4	18.4	17.8	63.3	62.3	62.1	19.2	19.3	20.1
65–74 years	25.5	24.1	24.3	58.2	60.3	59.0	16.2	15.5	16.6
75+ years	32.4	34.3	36.6	57.0	56.0	55.1	10.5	9.7	8.3
Total	17.4	19.9	22.0	61.8	59.6	59.9	20.6	20.5	18.2

(a) Not consumed alcohol in the previous 12 months.

(b) On average, had no more than 2 standard drinks per day.

(c) On average, had more than 2 standard drinks per day.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.72

Table 12A.72 **Single occasion risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent) (a), (b)**

	Low risk (c)			Single occasion risky											
				At least yearly (d)			At least monthly (e)			At least weekly (f)			Every day/most days (g)		
	2007	2010	2013	2007	2010	2013	2007	2010	2013	2007	2010	2013	2007	2010	2013
<i>Males</i>															
14–19 years	21.2	18.0	17.3	10.8	7.87	7.3	16.4	17.8	16.4	18.4	16.9	11.3	2.6	2.5	**0.5
20–29 years	15.8	18.5	19.7	11.7	11.4	14.7	22.5	22.4	22.2	32.2	27.6	25.2	6.4	5.9	4.1
18–24 years	14.4	15.8	17.2	12.3	9.2	13.1	18.8	22.5	25.4	36.5	32.6	25.7	5.7	5.0	2.1
25–34 years	20.0	21.0	21.7	16.3	14.9	15.3	23.2	22.4	21.5	23.0	20.9	21.8	6.3	7.2	5.3
35–44 years	28.6	28.0	27.7	16.3	16.6	16.6	17.4	16.3	17.3	17.5	17.7	15.0	9.1	9.1	9.3
45–54 years	35.1	34.6	34.9	15.4	13.7	12.9	16.9	14.4	14.6	12.5	14.0	13.6	9.9	11.3	8.4
55–64 years	44.1	42.6	42.3	12.5	11.8	10.6	11.6	11.0	12.1	8.4	10.0	10.0	11.3	10.4	9.7
65–74 years	55.2	56.8	53.5	7.8	7.0	6.6	6.2	6.0	6.9	5.0	5.2	4.9	8.8	7.9	9.0
75+ years	62.0	63.1	61.9	5.7	3.6	3.9	3.4	3.0	2.1	2.1	2.3	2.1	3.8	4.4	4.2
Total	32.8	32.5	32.7	13.1	12.0	12.1	15.7	15.2	15.4	16.2	15.8	14.1	7.9	7.9	6.7
<i>Females</i>															
14–19 years	27.7	23.4	22.1	11.5	8.8	10.2	18.0	19.0	14.6	11.8	11.6	6.4	1.4	*0.6	**0.6
20–29 years	27.3	28.6	31.6	19.1	17.7	17.5	19.9	19.1	18.8	16.6	17.1	11.0	1.9	1.7	1.8
18–24 years	21.8	25.2	26.0	17.7	13.5	15.8	22.6	22.5	24.3	21.7	22.8	14.4	1.4	1.9	*1.6
25–34 years	38.9	35.0	40.6	19.7	18.2	16.6	15.4	16.7	12.5	9.7	9.8	7.1	2.2	1.8	1.8
35–44 years	47.4	46.9	47.9	18.5	16.1	15.1	11.2	11.4	9.5	6.6	7.3	6.6	2.7	2.0	2.7
45–54 years	58.0	55.6	53.8	13.0	10.9	10.5	6.4	7.0	7.6	4.2	4.7	6.9	2.8	2.9	3.1
55–64 years	62.1	62.1	63.5	7.5	7.1	7.3	4.0	4.3	4.1	2.4	1.9	1.7	1.7	2.1	2.9
65–74 years	59.9	63.8	63.0	3.0	1.92	2.7	1.7	1.8	2.1	0.3	*0.5	*1.1	1.1	1.1	1.5
75+ years	56.2	54.3	52.3	0.8	*1.4	*1.3	1.1	*0.8	*1.0	0.5	**0.5	**0.2	1.1	*1.2	*0.5
Total	47.7	46.6	47.6	12.8	11.1	10.8	10.0	10.3	9.1	6.8	7.1	5.7	2.0	1.9	2.1
<i>All people</i>															

TABLE 12A.72

Table 12A.72 **Single occasion risk status recent drinkers (in last 12 months) aged 14 years or over, by age group (per cent) (a), (b)**

	Low risk (c)			Single occasion risky											
				At least yearly (d)			At least monthly (e)			At least weekly (f)			Every day/most days (g)		
	2007	2010	2013	2007	2010	2013	2007	2010	2013	2007	2010	2013	2007	2010	2013
14–19 years	24.4	20.6	19.6	11.1	8.3	8.7	17.2	18.4	15.5	15.1	14.3	8.9	2.0	1.6	*0.5
20–29 years	21.4	23.5	25.5	15.3	14.5	16.0	21.2	20.8	20.5	24.6	22.4	18.2	4.2	3.9	3.0
18–24 years	18.0	20.2	21.4	14.9	11.3	14.4	20.6	22.5	24.9	29.3	27.9	20.3	3.7	3.6	1.8
25–34 years	29.6	28.0	31.2	18.0	16.5	15.9	19.3	19.5	17.0	16.2	15.4	14.4	4.3	4.5	3.5
35–44 years	38.1	37.7	37.9	17.4	16.3	15.8	14.2	13.8	13.4	12.0	12.4	10.7	5.9	5.5	5.9
45–54 years	46.4	45.1	44.5	14.2	12.3	11.7	11.7	10.7	11.0	8.4	9.4	10.2	6.4	7.1	5.7
55–64 years	53.6	52.3	52.9	9.9	9.5	9.0	7.6	7.6	8.2	5.2	6.0	5.9	6.2	6.3	6.3
65–74 years	57.6	60.3	58.3	5.4	4.5	4.7	3.9	3.9	4.5	2.6	2.8	3.0	4.9	4.4	5.2
75+ years	58.8	57.9	56.4	3.0	2.3	2.4	2.1	1.7	1.5	1.2	1.2	1.0	2.3	2.5	2.1
Total	40.3	39.6	40.2	12.9	11.6	11.4	12.8	12.8	12.2	11.4	11.4	9.9	4.9	4.8	4.4

(a) Data on abstainers are in table 12A.69.

(b) Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with relative standard errors greater than 50 per cent should be considered as unreliable for most practical purposes. Estimates that have RSEs greater than 50 per cent are marked with " ** " and those with RSEs of between 25 per cent and 50 per cent are marked with " * ".

(c) Never had more than 4 standard drinks on any occasion.

(d) Had more than 4 standard drinks at least once a year but not as often as monthly.

(e) Had more than 4 standard drinks at least once a month but not as often as weekly.

(f) Had more than 4 standard drinks at least once a week but not as often as most days.

(g) Had more than 4 standard drinks on most days or every day.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.73

Table 12A.73 **Selected illicit drug use, by substance and age group (per cent)**
(a), (b)

	1995	1998	2001	2004	2007	2010	2013
<i>Cannabis</i>							
14–19 years	29.2	35.1	24.6	17.9	12.9	15.7	14.8
20–29 years	33.5	36.9	29.3	26.0	20.8	21.3	20.8
30–39 years	13.4	20.3	16.1	15.9	12.1	13.6	12.3
40–49 years	5.2	11.5	8.7	8.7	8.3	9.4	9.5
50–59 years	1.5	6.3	3.3	3.2	3.8	5.5	7.3
60 years or over	0.3	1.2	0.5	0.3	0.5	0.5	1.2
14 years or over	13.1	17.9	12.9	11.3	9.1	10.3	10.2
<i>Ecstasy</i>							
14–19 years	0.6	3.1	5.0	4.3	5.0	2.8	3.0
20–29 years	4.0	8.4	10.4	12.0	11.2	9.9	8.6
30–39 years	0.5	1.3	2.4	4.0	4.7	3.9	2.6
40 years or over	–	0.4	0.2	0.3	0.6	0.5	0.5
14 years or over	0.9	2.4	2.9	3.4	3.5	3.0	2.5
<i>Meth/amphetamines</i>							
14–19 years	2.7	5.9	6.2	4.4	1.6	1.6	*2.0
20–29 years	8.4	12.0	11.2	10.7	7.3	5.9	5.8
30–39 years	1.3	2.6	3.1	4.1	3.9	3.4	3.1
40 years or over	0.2	0.5	0.4	0.4	0.4	0.5	0.6
14 years or over	2.1	3.7	3.4	3.2	2.3	2.1	2.1
<i>Cocaine</i>							
14–19 years	1.1	0.8	1.5	1.0	1.1	1.3	*1.1
20–29 years	4.0	3.9	4.3	3.0	5.1	6.5	5.9
30–39 years	0.8	1.8	1.5	1.8	2.9	3.7	3.5
40 years or over	–	0.3	0.3	0.2	0.3	0.4	0.7
14 years or over	1.0	1.4	1.3	1.0	1.6	2.1	2.1

(a) Used in the previous 12 months.

(b) Results subject to RSEs of between 25 per cent and 50 per cent are marked with " * " and should be considered with caution.

– Nil or rounded to zero.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.74

Table 12A.74 **Selected illicit drug use by people aged 18 years or over, by level of psychological distress and self-reported health conditions (per cent) (a), (b)**

	<i>Not used drug in last 12 months</i>			<i>Used drug in last 12 months</i>			<i>All people (18+)</i>
	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2013</i>
<i>Any illicit drug</i>							
<i>Level of psychological distress</i>							
Low	71.7	71.8	71.6	52.7	57.3	56.7	69.3
Moderate	19.9	19.6	19.7	29.2	25.8	25.9	20.6
High	6.6	6.5	6.5	13.4	12.8	11.1	7.2
Very high	1.7	2.1	2.1	4.7	4.1	6.4	2.8
<i>Self-reported health condition (c)</i>							
Diabetes	5.9	5.7	6.7	2.8	3.3	3.6	6.3
Heart diseases (d)	19.6	20.4	21.7	8.8	10.1	11.6	20.4
Asthma	8.2	8.3	9.2	10.5	10.3	10.9	9.5
Cancer	2.9	3.0	3.1	1.1	1.4	2.0	2.9
Mental illness (e)	10.3	10.8	12.6	16.1	18.7	20.7	13.9
<i>Cannabis</i>							
<i>Level of psychological distress</i>							
Low	70.1	71.1	70.7	52.8	56.7	57.1	69.3
Moderate	20.8	19.8	20.1	28.0	27.0	25.9	20.6
High	7.2	6.8	6.8	14.6	12.7	11.0	7.2
Very high	1.9	2.3	2.4	4.6	3.6	5.9	2.8
<i>Self-reported health condition (c)</i>							
Diabetes	5.8	5.8	6.8	1.4	2.0	1.4	6.3
Heart diseases (d)	19.0	20.5	21.8	5.8	5.9	7.2	20.4
Asthma	8.4	8.5	9.5	10.5	10.0	9.8	9.5
Cancer	2.8	3.0	3.2	0.8	0.9	1.1	2.9
Mental illness (e)	10.8	11.3	13.0	15.7	18.7	21.2	13.9
<i>Ecstasy</i>							
<i>Level of psychological distress</i>							
Low	69.9	70.1	70.0	49.5	55.9	51.4	69.3
Moderate	20.7	20.2	20.4	31.3	28.9	30.7	20.6
High	7.3	7.3	7.0	16.0	12.1	12.0	7.2
Very high	2.1	2.4	2.6	3.2	3.0	6.0	2.8
<i>Self-reported health condition (c)</i>							
Diabetes	5.7	5.5	6.5	1.1	**1.0	**0.8	6.3
Heart diseases (d)	18.9	19.5	20.9	3.2	*1.2	*1.7	20.4
Asthma	8.4	8.6	9.5	11.2	11.0	9.8	9.5
Cancer	2.8	2.9	3.0	0.3	**0.2	**0.6	2.9
Mental illness (e)	10.9	11.9	13.6	16.0	16.2	17.9	13.9

TABLE 12A.74

Table 12A.74 **Selected illicit drug use by people aged 18 years or over, by level of psychological distress and self-reported health conditions (per cent) (a), (b)**

	<i>Not used drug in last 12 months</i>			<i>Used drug in last 12 months</i>			<i>All people (18+)</i>
	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2013</i>
<i>Meth/amphetamines</i>							
<i>Level of psychological distress</i>							
Low	69.9	70.1	70.0	44.7	51.2	41.7	69.3
Moderate	20.9	20.3	20.4	31.7	28.0	31.8	20.6
High	7.2	7.3	7.0	19.0	13.3	15.6	7.2
Very high	2.0	2.3	2.6	4.6	7.5	10.9	2.8
<i>Self-reported health condition (c)</i>							
Diabetes	5.7	5.5	6.4	0.9	*1.5	*1.4	6.3
Heart diseases (d)	18.8	19.3	20.7	3.7	4.5	5.2	20.4
Asthma	8.4	8.6	9.5	11.4	11.2	11.5	9.5
Cancer	2.8	2.9	3.0	0.1	*0.7	*1.9	2.9
Mental illness (e)	10.9	11.7	13.5	20.3	25.6	29.0	13.9
<i>Cocaine</i>							
<i>Level of psychological distress</i>							
Low	69.6	70.0	69.8	45.8	55.0	55.3	69.3
Moderate	20.9	20.3	20.4	35.8	27.4	27.4	20.6
High	7.4	7.3	7.1	14.4	14.1	11.7	7.2
Very high	2.1	2.4	2.7	3.9	3.4	5.6	2.8
<i>Self-reported health condition (c)</i>							
Diabetes	5.6	5.5	6.4	0.9	**0.5	**0.8	6.3
Heart diseases (d)	18.6	19.4	20.7	4.4	*2.3	*3.5	20.4
Asthma	8.4	8.7	9.6	12.1	6.7	7.8	9.5
Cancer	2.7	2.9	3.0	0.2	**0.4	**1.0	2.9
Mental illness (e)	11.1	11.9	13.7	15.2	17.4	17.4	13.9

(a) Recent use means used in the previous 12 months.

(b) Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with relative standard errors greater than 50 per cent should be considered as unreliable for most practical purposes. Estimates that have RSEs greater than 50 per cent are marked with " ** " and those with RSEs of between 25 per cent and 50 per cent are marked with " * ".

(c) Respondents could select more than one condition in response to the question 'In the last 12 months have you been diagnosed or treated for...?'.
(d) Includes heart disease and hypertension (high blood pressure).

(e) Includes depression, anxiety disorder, schizophrenia, bipolar disorder, an eating disorder and other form of psychosis.

(e) Includes depression, anxiety disorder, schizophrenia, bipolar disorder, an eating disorder and other form of psychosis.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.75

Table 12A.75 **Illicit drug use, people aged 14 years or older, by social characteristics (per cent) (a)**

	<i>Never used</i>			<i>Ex-users (b)</i>			<i>Recent users (c)</i>		
	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>
<i>All illicit drugs</i>									
<i>Socioeconomic status</i>									
Quintile 1 (lowest)	65.1	64.0	61.6	21.1	20.9	22.5	13.8	15.1	15.9
Quintile 2	65.3	61.8	59.7	23.4	22.7	25.4	11.4	15.5	15.0
Quintile 3	64.2	60.4	57.4	22.2	26.0	27.7	13.6	13.6	14.9
Quintile 4	60.5	59.5	57.5	26.2	26.5	27.9	13.3	14.0	14.6
Quintile 5 (highest)	59.5	57.1	56.3	26.0	27.7	28.7	14.5	15.2	15.0
<i>Geography</i>									
Major cities	62.5	60.6	59.1	23.7	24.6	26.0	13.8	14.8	14.9
Inner regional	64.4	61.2	58.0	23.8	24.9	27.9	11.8	13.9	14.1
Outer regional	61.0	59.9	55.9	26.2	25.0	27.4	12.8	15.0	16.7
Remote/Very remote	51.8	50.8	49.9	27.6	31.9	31.4	20.6	17.2	18.7
<i>Indigenous status</i>									
Aboriginal and/or Torres Strait Islander	47.1	46.5	47.4	28.5	28.5	28.5	24.4	25.0	24.1
Non-Indigenous	62.7	60.8	58.6	24.2	25.1	26.6	13.1	14.2	14.8
<i>Cannabis</i>									
<i>Socioeconomic status</i>									
Quintile 1 (lowest)	70.9	68.7	69.7	20.0	21.0	20.0	9.0	10.3	10.3
Quintile 2	70.6	66.5	67.0	22.3	22.8	22.9	7.2	10.7	10.1
Quintile 3	69.1	64.5	63.1	21.9	25.4	26.2	8.9	10.1	10.7
Quintile 4	65.0	63.8	64.2	26.0	26.8	25.9	9.0	9.4	9.9
Quintile 5 (highest)	64.0	60.6	63.1	26.4	28.4	26.9	9.5	11.1	10.0
<i>Geography</i>									
Major cities	67.5	64.8	66.0	23.6	24.8	24.2	8.9	10.4	9.8
Inner regional	69.1	65.5	65.0	23.0	24.7	24.9	7.9	9.8	10.1
Outer regional	66.1	64.2	63.1	24.9	25.4	24.9	9.0	10.4	12.0
Remote/Very remote	58.3	55.2	56.1	27.6	33.4	30.3	14.1	11.4	13.6
<i>Indigenous status</i>									
Aboriginal and/or Torres Strait Islander	57.6	52.0	54.4	26.9	29.5	26.6	15.5	18.5	19.0
Non-Indigenous	67.4	64.8	65.4	23.8	25.2	24.6	8.7	10.0	10.0
<i>Ecstasy</i>									
<i>Socioeconomic status</i>									
Quintile 1 (lowest)	93.4	92.0	91.8	4.1	6.2	6.6	2.5	1.9	1.6
Quintile 2	93.8	90.8	91.5	4.0	6.6	6.6	2.3	2.5	2.0
Quintile 3	91.9	90.1	89.1	4.5	7.3	8.0	3.7	2.7	3.0
Quintile 4	90.6	88.7	88.9	5.8	8.1	8.3	3.6	3.1	2.8

TABLE 12A.75

Table 12A.75 **Illicit drug use, people aged 14 years or older, by social characteristics (per cent) (a)**

	<i>Never used</i>			<i>Ex-users (b)</i>			<i>Recent users (c)</i>		
	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>
Quintile 5 (highest)	88.7	87.6	88.6	6.7	8.0	8.5	4.6	4.4	2.9
<i>Geography</i>									
Major cities	90.4	88.9	89.3	5.7	7.8	7.9	3.9	3.3	2.9
Inner regional	93.8	91.9	91.9	3.8	6.1	6.6	2.5	2.0	1.5
Outer regional	93.7	91.9	91.7	4.5	5.9	6.7	1.8	2.2	1.6
Remote/Very remote	88.8	86.4	87.1	6.6	9.5	11.1	4.6	*4.1	*1.8
<i>Indigenous status</i>									
Aboriginal and/or Torres Strait Islander	90.2	89.9	89.3	6.1	*7.0	9.7	3.7	*3.0	**1.1
Non-Indigenous	91.4	89.7	89.9	5.2	7.3	7.6	3.4	3.0	2.5
<i>Meth/amphetamines</i>									
<i>Socioeconomic status</i>									
Quintile 1 (lowest)	93.8	92.9	93.4	3.9	4.7	4.4	2.3	2.4	2.2
Quintile 2	95.0	92.9	93.7	3.1	5.0	4.1	1.8	2.1	2.1
Quintile 3	94.3	93.4	92.3	3.2	4.5	5.3	2.5	2.1	2.4
Quintile 4	93.5	92.8	94.1	4.2	5.4	4.1	2.2	1.8	1.8
Quintile 5 (highest)	93.5	93.2	94.3	4.3	4.8	3.9	2.2	2.0	1.8
<i>Geography</i>									
Major cities	93.6	92.8	93.6	3.9	5.1	4.3	2.5	2.0	2.1
Inner regional	95.1	93.8	94.3	3.2	4.1	4.1	1.7	2.0	1.6
Outer regional	94.3	94.1	94.0	4.1	4.4	4.0	1.6	1.5	2.0
Remote/Very remote	91.3	88.8	87.0	5.7	7.2	8.6	3.0	*4.0	*4.4
<i>Indigenous status</i>									
Aboriginal and/or Torres Strait Islander	92.2	92.4	89.8	5.5	*4.0	7.0	2.3	*3.6	*3.1
Non-Indigenous	94.0	93.1	93.7	3.8	5.0	4.3	2.2	2.0	2.0
<i>Cocaine</i>									
<i>Socioeconomic status</i>									
Quintile 1 (lowest)	96.4	95.0	94.2	3.2	4.0	4.6	0.5	1.0	1.2
Quintile 2	96.1	94.7	94.2	3.2	3.9	4.4	0.7	1.4	1.4
Quintile 3	95.1	93.0	92.7	3.7	5.5	5.4	1.2	1.5	1.9
Quintile 4	93.8	92.2	91.2	4.4	5.6	6.3	1.8	2.2	2.5
Quintile 5 (highest)	90.7	89.5	90.0	6.0	6.2	6.5	3.3	4.3	3.5
<i>Geography</i>									
Major cities	93.1	91.8	91.6	4.8	5.6	5.8	2.1	2.6	2.6
Inner regional	96.8	94.7	94.7	2.6	4.3	4.5	0.6	1.0	0.8
Outer regional	95.3	95.7	94.1	4.3	3.4	4.8	0.4	*0.9	*1.1
Remote/Very remote	95.0	92.6	92.0	2.7	5.3	5.6	2.3	**2.0	*2.5

Table 12A.75 **Illicit drug use, people aged 14 years or older, by social characteristics (per cent) (a)**

	<i>Never used</i>			<i>Ex-users (b)</i>			<i>Recent users (c)</i>		
	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>	<i>2007</i>	<i>2010</i>	<i>2013</i>
<i>Indigenous status</i>									
Aboriginal and/or Torres Strait Islander	92.4	95.9	93.4	6.3	*3.2	4.8	1.3	**0.9	*1.9
Non-Indigenous	94.2	92.7	92.3	4.2	5.2	5.5	1.6	2.1	2.2

(a) Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with relative standard errors greater than 50 per cent should be considered as unreliable for most practical purposes. Estimates that have RSEs greater than 50 per cent are marked with " ** " and those with RSEs of between 25 per cent and 50 per cent are marked with " * ".

(b) Used, but not in the previous 12 months.

(c) Used in the previous 12 months.

Source: AIHW (2014) *National Drug Strategy Household Survey detailed report 2013*, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.

TABLE 12A.76

Table 12A.76 **Prevalence of lifetime mental disorders among adults aged 16–85 years, 2007 (per cent) (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
Any 12-month mental disorder (c)									
Anxiety disorders	14.4 ± 1.7	15.4 ± 2.0	13.1 ± 2.5	15.1 ± 3.7	14.4 ± 3.3	np	np	np	14.4 ± 0.9
Affective disorders	6.4 ± 1.2	6.6 ± 1.7	6.1 ± 1.6	6.2 ± 1.8	6.3 ± 2.3	np	np	np	6.2 ± 0.7
Substance use disorders	4.2 ± 1.1	5.5 ± 1.3	5.8 ± 1.8	6.0 ± 2.2	5.5 ± 2.0	np	np	np	5.1 ± 0.7
Any 12-month mental disorder (c), (d)	20.1 ± 2.2	20.7 ± 2.3	19.2 ± 2.6	21.4 ± 4.1	19.1 ± 3.4	14.1 ± 5.4	np	np	20.0 ± 1.1
Lifetime mental disorder, with no 12-month symptoms (e)	23.2 ± 1.9	26.3 ± 2.9	28.1 ± 3.4	23.6 ± 4.1	26.3 ± 4.1	30.7 ± 6.9	np	33.3 ± 12.9	25.5 ± 1.4
Without lifetime mental disorders (f)	56.7 ± 2.2	53.0 ± 3.6	52.6 ± 3.8	55.1 ± 5.2	54.6 ± 4.5	55.2 ± 8.2	53.1 ± 11.9	49.0 ± 18.8	54.5 ± 1.4

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) Estimates with RSEs greater than 25 per cent are considered unreliable. These estimates are not published.

(c) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(d) A person can have had more than one 12-month mental disorder. Therefore, the components may not add to the total.

(e) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(f) People who did not meet criteria for diagnosis of a lifetime mental disorder.

np Not published.

Source: ABS (unpublished) *2007 Survey of Mental Health and Wellbeing*, Cat. no. 4326.0.

Table 12A.77 Prevalence of lifetime mental disorders among adults aged 16–85 years, by sex, 2007 (per cent) (a)

	<i>Males</i>	<i>Females</i>	<i>People</i>
Any 12-month mental disorder (b), (c)			
Anxiety disorders			
Panic disorders	2.3 ± 0.7	2.8 ± 0.6	2.6 ± 0.5
Agoraphobia	2.1 ± 0.7	3.5 ± 0.7	2.8 ± 0.5
Social phobia	3.8 ± 1.0	5.7 ± 0.8	4.7 ± 0.6
Generalised anxiety disorder	2.0 ± 0.7	3.5 ± 0.8	2.7 ± 0.6
Obsessive compulsive disorder	1.6 ± 0.6	2.2 ± 0.5	1.9 ± 0.4
Post traumatic stress disorder	4.6 ± 1.0	8.3 ± 1.0	6.4 ± 0.6
<i>Any anxiety disorder (c)</i>	10.8 ± 1.4	17.9 ± 1.3	14.4 ± 0.9
Affective disorders			
Depression (d)	3.1 ± 0.8	5.1 ± 0.8	4.1 ± 0.6
Dysthymia	1.0 ± 0.4	1.5 ± 0.5	1.3 ± 0.3
Bipolar	1.8 ± 0.6	1.7 ± 0.4	1.8 ± 0.4
<i>Any affective disorder (c)</i>	5.3 ± 1.0	7.1 ± 1.0	6.2 ± 0.7
Substance use disorders			
Alcohol harmful use	3.8 ± 0.8	2.1 ± 0.6	2.9 ± 0.5
Alcohol dependence	2.2 ± 0.7	0.7 ± 0.2	1.4 ± 0.3
Drug use (e)	2.1 ± 0.6	0.8 ± 0.3	1.4 ± 0.3
<i>Any substance use disorder (c), (e)</i>	7.0 ± 1.2	3.3 ± 0.7	5.1 ± 0.7
Any 12-month mental disorder (c)	17.6 ± 1.9	22.3 ± 1.3	20.0 ± 1.1
Lifetime mental disorder, with no 12-month symptoms (f)	30.5 ± 2.2	20.7 ± 1.4	25.5 ± 1.4
No lifetime mental disorder (g)	51.9 ± 2.0	57.0 ± 1.7	54.5 ± 1.4

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(c) A person can have had more than one 12-month mental disorder. Therefore, the components may not add to the total.

(d) Includes severe depressive episode, moderate depressive episode and mild depressive episode.

(e) Includes harmful use and dependence.

(f) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(g) People who did not meet criteria for diagnosis of a lifetime mental disorder.

Source: ABS (unpublished) *2007 Survey of Mental Health and Wellbeing*, Cat. no. 4326.0.

TABLE 12A.78

Table 12A.78 **Prevalence of lifetime mental disorders among adults, by age, 2007 (per cent) (a), (b)**

	16–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	75–85 years
Any 12-month mental disorder (c), (d)							
Anxiety disorders	15.4 ± 2.0	16.3 ± 2.8	18.1 ± 3.0	17.6 ± 3.0	11.3 ± 1.9	6.3 ± 1.5	4.0 ± 1.8
Affective disorders	6.3 ± 1.5	7.9 ± 2.1	8.3 ± 2.1	7.1 ± 2.2	4.2 ± 1.3	2.8 ± 1.2	np
Substance use disorders	12.7 ± 2.0	7.3 ± 2.2	4.6 ± 1.6	3.8 ± 1.6	np	np	np
Any 12-month mental disorder (c), (d)	26.4 ± 2.7	24.8 ± 3.2	23.3 ± 3.3	21.5 ± 3.5	13.6 ± 2.1	8.6 ± 1.6	5.9 ± 2.1
Lifetime mental disorder, with no 12-month symptoms (e)	13.2 ± 2.0	29.0 ± 4.4	30.7 ± 3.3	30.4 ± 4.2	27.6 ± 3.6	23.1 ± 2.6	16.2 ± 4.1
No lifetime mental disorder (f)	60.5 ± 3.0	46.2 ± 3.9	46.0 ± 3.3	48.2 ± 4.6	58.8 ± 4.1	68.3 ± 3.0	77.8 ± 4.6

(a) Estimates with RSEs greater than 25 per cent are considered unreliable. These estimates are not published.

(b) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(c) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(d) A person can have had more than one 12-month mental disorder. Therefore, the components may not add to the total.

(e) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(f) People who did not meet criteria for diagnosis of a lifetime mental disorder.

np Not published.

Source: ABS (unpublished) *2007 Survey of Mental Health and Wellbeing*, Cat. no. 4326.0.

TABLE 12A.79

Table 12A.79 **Prevalence of lifetime mental disorders among adults, by disadvantage and section of state, 2007 (per cent) (a)**

	<i>Mental disorders with symptoms in last 12 months (b)</i>			<i>Total mental disorders with symptoms in last 12 months (b), (c)</i>	<i>Total without mental disorders with symptoms in last 12 months (d)</i>	<i>Total</i>
	<i>Anxiety disorders</i>	<i>Affective disorders</i>	<i>Substance use disorders</i>			
<i>Index of disadvantage</i>						
1st quintile	15.8 ± 2.6	7.5 ± 2.1	5.6 ± 1.8	21.5 ± 3.0	78.5 ± 2.9	100.0
5th quintile	10.9 ± 2.3	4.2 ± 1.1	4.0 ± 1.2	15.9 ± 2.7	84.1 ± 2.8	100.0
<i>Section of state</i>						
Major urban	14.6 ± 1.2	6.5 ± 0.8	5.5 ± 0.8	20.4 ± 1.4	79.6 ± 1.4	100.0
Other urban	13.3 ± 2.1	6.1 ± 1.4	4.8 ± 1.3	19.2 ± 2.7	80.8 ± 2.7	100.0
Balance of state	14.9 ± 2.7	5.1 ± 1.7	3.5 ± 1.2	19.2 ± 2.6	80.8 ± 2.7	100.0
All people aged 16–85 years	14.4 ± 0.9	6.2 ± 0.7	5.1 ± 0.6	20.0 ± 1.1	80.0 ± 1.1	100.0

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(c) A person can have had more than one 12-month mental disorder. Therefore, the components may not add to the total.

(d) People who did not meet criteria for diagnosis of a lifetime mental disorder and people who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

Source: ABS (2008) *2007 Survey of Mental Health and Wellbeing*, Cat. no. 4326.0.

Table 12A.80 **Suicides and mortality rate, by sex, Australia (a), (b), (c)**

	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Suicides (no.)										
Males	1 661	1 658	1 624	1 699	1 833	1 785	1 914	1 812	1 925	1 885
Females	437	444	494	530	508	552	566	581	649	637
People	2 098	2 102	2 118	2 229	2 341	2 337	2 480	2 393	2 574	2 522
Suicide death rate (per 100 000 people) (d)										
Males	16.8	16.5	16.0	16.4	17.3	16.5	17.5	16.3	17.0	16.4
Females	4.4	4.4	4.8	5.1	4.8	5.1	5.1	5.2	5.7	5.5
People	10.5	10.4	10.4	10.7	11.0	10.8	11.3	10.7	11.3	10.9

- (a) Suicide deaths include ICD-10 codes X60–X84 and Y87.0. Care needs to be taken in interpreting figures relating to suicide. See ABS *Causes of Death, 2013* (Cat. no. 3303.0) Explanatory Notes 87–93.
- (b) By year of registration. Year-to-year variation can be influenced by coronial workloads.
- (c) Data for 2006 to 2011 have undergone revisions and are now considered final. Data for 2012 have been revised and are subject to further revisions. Data for 2013 are preliminary and subject to a revisions process. See ABS' *Causes of Death, Australia 2013*, publication for more information.
- (d) Crude death rate per 100 000 people using estimated resident populations (ERPs) for Australia (people) at 30 June of relevant year.

Source: ABS (2015) *Causes of Death, Australia 2013*, Cat. no. 3303.0, Canberra.

TABLE 12A.81

Table 12A.81 **Suicides and mortality rate, by age and sex, Australia (a), (b)**

	15–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	75–84 years	85 years +	All ages (c)
2013									
Suicides (no.)									
Males	256	314	373	364	257	149	96	60	1 885
Females	92	93	137	124	92	45	28	18	637
People	348	407	510	488	349	194	124	78	2 522
Suicide death rate (per 100 000 people) (d), (e)									
Males	16.1	18.4	23.3	23.9	19.7	16.2	20.4	38.3	16.4
Females	6.1	5.5	8.5	8.0	6.9	4.8	5.0	6.4	5.5
People	11.2	12.0	15.9	15.9	13.2	10.4	12.0	17.7	10.9
2009–2013									
Suicides (no.)									
Males	1 151	1 664	1 985	1 847	1 224	664	510	239	9 321
Females	442	476	615	586	409	192	154	75	2 985
People	1 593	2 140	2 600	2 433	1 633	856	664	314	12 306
Suicide death rate (per 100 000 people) (e)									
Males	14.7	20.7	25.3	24.6	19.2	16.0	22.7	34.4	16.8
Females	5.9	6.0	7.7	7.6	6.3	4.5	5.6	5.7	5.3
People	10.4	13.4	16.4	16.0	12.7	10.2	13.2	15.6	11.0

(a) Suicide deaths include ICD-10 codes X60–X84 and Y87.0. Care needs to be taken in interpreting data relating to suicide. See ABS Causes of Death, 2013 (Cat. no. 3303.0) Explanatory Notes 87–93.

(b) Data for 2006 to 2011 have undergone revisions and are now considered final. Data for 2012 have been revised and are subject to further revisions. Data for 2013 are preliminary and subject to a revisions process. See ABS' Causes of Death, Australia 2013, publication for more information.

(c) All ages includes deaths of people aged under 15 years and age not stated.

(d) Crude death rate per 100 000 estimated resident population as at 30 June 2013 for each age group and sex.

(e) Rate per 100 000 estimated resident population at 30 June of the relevant mid point year (for 2009–2013 it is 2011). The total death rate per 100 000 people for 2009–2013 does not match that in tables 12A.82–84 as it is a crude rate and the rates in tables 12A.82–84 are age standardised.

Table 12A.81 **Suicides and mortality rate, by age and sex, Australia (a), (b)**

	<i>15–24 years</i>	<i>25–34 years</i>	<i>35–44 years</i>	<i>45–54 years</i>	<i>55–64 years</i>	<i>65–74 years</i>	<i>75–84 years</i>	<i>85 years +</i>	<i>All ages (c)</i>
--	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------	-------------------	---------------------

Source: ABS (2015) *Causes of Death, Australia 2013*, Cat. no. 3303.0, Canberra; ABS (2015) *Australian Demographic Statistics*, Cat. no. 3101.0.

Table 12A.82 **Suicide deaths and death rate (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas (c)</i>	<i>ACT (c)</i>	<i>NT (c)</i>	<i>Aust</i>
Suicide deaths (no.)									
2004	587	521	453	194	178	88	26	51	2 098
2005	549	506	459	203	231	74	35	45	2 102
2006 (d)	577	485	494	245	180	72	32	33	2 118
2007 (d)	611	474	520	266	205	66	32	55	2 229
2008 (d)	620	545	553	300	175	73	36	38	2 341
2009 (d)	623	576	525	279	185	79	32	37	2 337
2010 (d)	674	558	588	313	197	64	41	45	2 480
2011 (d)	617	526	578	309	212	74	33	44	2 393
2012 (e)	728	511	629	366	198	70	24	48	2 574
2013 (f)	696	488	663	332	199	74	37	33	2 522
2009–2013	3 338	2 659	2 983	1 599	991	361	167	207	12 306
Suicide death rate per 100 000 people (g), (h), (i)									
2004	8.7	10.5	11.7	9.8	11.6	18.2	8.0	25.5	10.4
2005	8.0	9.8	11.6	10.1	14.9	15.8	10.5	21.7	10.3
2006 (d)	8.4	9.5	12.4	12.1	11.5	14.8	9.7	14.2	10.3
2007 (d)	8.8	9.0	12.5	12.6	12.9	14.1	9.1	26.5	10.5
2008 (d)	8.9	10.3	13.3	14.2	11.0	15.2	10.3	17.9	11.1
2009 (d)	8.7	10.5	12.1	12.3	11.5	15.4	8.9	17.4	10.7
2010 (d)	9.3	10.1	13.4	13.6	11.8	13.0	11.3	18.8	11.2
2011 (d)	8.4	9.2	12.9	12.9	12.9	14.1	9.3	18.5	10.5
2012 (e)	9.8	8.9	13.8	14.9	11.7	13.4	6.2	19.2	11.2
2013 (f)	9.2	8.2	14.3	13.1	11.6	14.3	9.6	14.3	10.7
2009–2013 (g)	9.1	9.4	13.3	13.4	11.9	14.0	9.1	17.6	10.9

- (a) By year of registration. Year-to-year variation can be influenced by coronial workloads.
- (b) Suicide deaths include ICD-10 codes X60–X84 and Y87.0. Care needs to be taken in interpreting figures relating to suicide due to limitations of data.
- (c) Low population results in small variations in the number of suicides appearing as large changes across the single year rates.
- (d) Data for 2006 to 2011 have undergone revisions and are now considered final.
- (e) Data for 2012 have been revised and are subject to further revisions.
- (f) Data for 2013 are preliminary and subject to a revisions process.
- (g) Rate per 100 000 estimated resident population at 30 June of the relevant single year or for five year average the mid-point year (2009–2013). 2009–2013 rate includes final 2009, 2010 and 2011 data, revised 2012 data and preliminary 2013 data.
- (h) Death rates standardised to the mid-year 2001 population. The total death rate per 100 000 people for 2009–2013 does not match that in table 12A.81 as it reports the crude rate.
- (i) The ERPs used to derived these rates differ across years. For data up to 2005 the rates are derived using ERPs based on the 2001 Census. For data from 2006 the rates are derived using ERPs based on the 2006 Census. For data from 2009 (and for the five year averages 2009–2013) the rates are derived using the ERPs based on the 2011 Census. Rates derived using ERPs based on different Censuses are not comparable.

Table 12A.82 **Suicide deaths and death rate (a), (b)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas (c)</i>	<i>ACT (c)</i>	<i>NT (c)</i>	<i>Aust</i>
Source:	ABS (2015) <i>Causes of Death, Australia 2013</i> , Cat. no. 3303.0, Canberra; ABS (unpublished) <i>Causes of Death, Australia</i> , Cat. no. 3303.0.								

Table 12A.83 Suicide deaths and death rate of people aged 15–24 years (a), (b), (c), (d), (e)

	NSW	Vic	Qld	WA	SA	Tas (f)	ACT (f)	NT (f)	Aust (g)
Number of suicide deaths of people aged 15–24 years									
2004	75	66	54	23	22	np	3	np	265
2005	66	61	67	30	37	9	5	15	290
2006	74	61	74	41	25	9	6	8	298
2007	54	74	81	46	19	4	3	21	300
2008	62	63	80	44	21	np	6	9	288
2009	63	60	63	47	21	8	np	11	276
2010	65	79	76	37	22	7	3	11	299
2011	61	65	89	53	36	10	5	16	335
2012	75	64	81	56	20	10	1	18	327
2013	84	66	93	51	28	9	5	12	348
2009–2013	348	334	402	244	127	44	14	68	1 585
Suicide death rate per 100 000 people aged 15–24 years (h), (i)									
2004	8.3	9.7	9.9	8.2	10.8	np	5.8	np	9.5
2005	7.2	8.9	11.9	10.5	17.9	13.9	9.7	48.1	10.2
2006	8.0	8.5	12.8	13.8	11.7	13.8	11.1	24.5	10.3
2007	5.7	10.1	13.6	15.1	8.8	4.6	3.6	62.6	10.1
2008	6.4	8.3	13.0	14.0	9.6	np	11.0	26.1	9.5
2009	6.5	7.8	10.3	14.5	9.6	12.1	7.0	30.8	9.1
2010	6.7	10.3	12.2	11.3	9.9	10.5	np	30.4	9.8
2011	6.4	8.5	14.2	16.0	16.3	15.1	8.5	45.1	10.9
2012	7.8	8.3	12.8	16.6	9.1	15.2	np	50.9	10.6
2013	8.6	8.6	14.5	14.8	12.7	13.8	8.8	33.7	11.2
2009–2013 (i)	7.2	8.7	13.1	14.7	11.5	13.3	6.2	38.3	10.4

(a) By year of registration. Year-to-year variation can be influenced by coronial workloads.

(b) Suicide deaths include ICD-10 codes X60–X84 and Y87.0. Care needs to be taken in interpreting data relating to suicide. See ABS Causes of Death, 2013 (Cat. no. 3303.0) Explanatory Notes 87–93.

(c) From 2006 data onwards, data cells with small values have been randomly assigned to protect the confidentiality of individuals. As a result, some totals will not equal the sum of their components. Rates use the actual count and not the randomly assigned value. Cells with a zero value have not been affected by confidentialisation.

(d) All footnotes and caveats, including this notice, must remain attached to data at all times.

(e) All causes of death data from 2006 onward are subject to a revisions process – once data for a reference year are 'final', they are no longer revised. Revised data for this year's report are: 2011 (final), 2012 (revised) and the data for 2013 are preliminary. See Explanatory Notes 52–54 and Technical Note, Causes of Death Revisions 2011 and 2012 in *Causes of Death, Australia, 2013* (Cat. no. 3303.0).

(f) Low population results in small variations in the number of suicides appearing as large changes across the single year rates.

(g) Includes 'Other Territories'.

Table 12A.83 Suicide deaths and death rate of people aged 15–24 years (a), (b), (c), (d), (e)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i> (f)	<i>ACT</i> (f)	<i>NT</i> (f)	<i>Aust</i> (g)
(h) Rate per 100 000 ERP at 30 June of the relevant single year or for five year average the mid-point year (2009–2013). 2009–2013 rate includes final 2009, 2010 and 2011 data, revised 2012 data and preliminary 2013 data.									
(i) The ERPs used to derived these rates differ across years. For data up to 2005 the rates are derived using ERPs based on the 2001 Census. For data from 2006 the rates are derived using ERPs based on the 2006 Census. For data from 2009 (and for the five year averages 2009–2013) the rates are derived using the ERPs based on the 2011 Census. Rates derived using ERPs based on different Censuses are not comparable.									
np not published									

Source: ABS (2015) *Causes of Death, Australia 2013*, Cat. no. 3303.0, Canberra; ABS (unpublished) *Causes of Death, Australia*, Cat. no. 3303.0.

TABLE 12A.84

Table 12A.84 **Suicide deaths and suicide death rate, by area (a), (b), (c), (d), (e), (f), (g), (h), (i), (j)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas (k)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (c)</i>
<i>Number of suicide deaths by area</i>									
2004									
Capital city	358	345	194	141	125	29	26	22	1 240
Other urban	192	122	199	38	np	37	629
Rural	32	50	55	15	21	22	..	16	211
2005									
Capital city	342	332	179	142	173	29	35	23	1 255
Other urban	186	124	204	45	33	31	..	11	634
Rural	19	49	69	14	25	12	..	11	199
2006									
Capital city	340	330	187	157	133	28	32	14	1 221
Urban centres	129	64	171	19	..	20	403
Rural	108	91	136	69	47	24	–	19	494
2007									
Capital city	393	327	189	180	148	22	32	27	1 318
Urban centres	140	63	191	20	–	25	–	..	439
Rural	76	84	137	65	57	18	–	27	464
2008									
Capital city	362	374	216	219	125	27	36	23	1 382
Urban centres	127	76	215	27	..	26	471
Rural	131	95	122	54	50	20	–	15	487
2009									
Capital city	326	385	198	194	145	35	32	15	1 330
Urban centres	208	107	198	35	18	22	..	2	591
Rural	87	81	124	44	20	22	–	19	398
2010									
Capital city	352	366	220	225	151	33	41	20	1 408
Urban centres	204	100	227	41	17	15	..	3	608
Rural	115	89	133	45	26	16	–	21	445
2011									
Capital city	317	355	252	196	155	31	32	12	1 350
Urban centres	206	87	213	47	18	25	na	4	600
Rural	91	82	111	65	39	18	3	27	434
2012									
Capital city	349	336	247	253	141	31	24	14	1 395
Urban centres	227	81	251	48	20	17	..	7	651
Rural	149	90	127	62	37	21	–	26	512
2013									
Capital city	336	318	288	229	147	30	37	9	1 394
Urban centres	217	79	220	36	18	20	..	3	593

TABLE 12A.84

Table 12A.84 **Suicide deaths and suicide death rate, by area (a), (b), (c), (d), (e), (f), (g), (h), (i), (j)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas (k)</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (c)</i>
Rural	140	89	151	64	34	24	–	21	523
2009–2013									
Capital city	1 682	1 758	1 202	1 093	737	158	166	70	6 866
Urban centres	1 051	454	1 113	212	92	99	..	21	3 042
Rural	593	435	646	280	162	103	2	114	2 334
<i>Suicide death rate per 100 000 people by area (l)</i>									
2004									
Capital city	8.5	9.6	10.9	9.7	11.1	14.3	8.0	20.1	9.7
Other urban	8.8	11.8	12.3	9.9	np	19.9	..	np	11.1
Rural	10.2	14.7	11.1	10.9	12.0	23.4	–	37.1	13.2
Total	8.7	10.5	11.7	9.8	11.6	18.2	8.0	25.5	10.4
2005									
Capital city	7.8	8.9	9.8	9.5	15.0	14.5	10.5	19.6	9.5
Other urban	8.6	12.2	12.3	11.6	14.8	17.0	..	22.2	11.2
Rural	6.5	14.7	13.9	9.6	13.5	12.9	..	27.2	12.5
Total	8.0	9.8	11.6	10.1	14.9	15.8	10.5	21.7	10.3
2006									
Capital city	7.8	8.8	10.3	10.5	11.5	13.8	9.7	np	9.2
Urban centres	10.0	11.0	12.8	np	..	10.9	11.3
Rural	9.3	11.7	16.6	20.9	11.5	23.8	..	np	13.2
Total	8.4	9.5	12.4	12.1	11.5	14.8	9.7	14.2	10.3
2007									
Capital city	8.8	8.3	10.1	11.4	12.5	10.7	9.1	22.8	9.6
Urban centres	10.8	10.7	12.9	10.3	..	13.9	11.7
Rural	6.4	10.9	17.0	18.9	13.8	18.5	..	29.1	12.5
Total	8.8	9.0	12.5	12.6	12.9	14.1	9.1	26.5	10.5
2008									
Capital city	8.3	9.5	11.5	14.0	10.6	13.3	10.3	20.2	10.2
Urban centres	9.8	12.9	14.4	13.9	..	13.6	12.6
Rural	10.9	11.9	15.2	16.5	12.6	22.4	..	np	13.2
Total	8.9	10.3	13.3	14.2	11.0	15.2	10.3	17.9	11.1
2009									
Capital city	7.8	9.6	9.9	11.2	11.9	17.0	8.9	np	9.6
Urban centres	11.1	13.8	12.3	13.7	np	13.9	..	np	12.2
Rural	9.2	12.0	17.5	15.6	8.1	14.7	–	np	12.8
Total	8.7	10.5	12.1	12.3	11.5	15.4	8.9	17.4	10.7
2010									
Capital city	8.3	9.1	10.9	12.9	11.9	16.5	11.4	17.7	10.0
Urban centres	10.6	12.6	14.1	15.7	np	np	..	np	12.4
Rural	11.9	12.6	17.9	15.5	10.0	np	–	21.1	14.1

Table 12A.84 **Suicide deaths and suicide death rate, by area (a), (b), (c), (d), (e), (f), (g), (h), (i), (j)**

	NSW	Vic	Qld	WA	SA	Tas (k)	ACT	NT	Aust (c)
Total	9.3	10.1	13.4	13.6	11.8	13.0	11.3	18.8	11.2
2011									
Capital city	7.3	8.5	12.2	10.8	12.4	14.4	9.0	10.3	9.4
Urban centres	10.7	10.5	12.9	17.2	np	16.1	na	np	12.0
Rural	9.5	11.5	15.3	22.0	15.2	np	np	30.7	13.9
Total	8.4	9.2	12.9	12.9	12.9	14.1	9.3	18.5	10.5
2012									
Capital city	7.9	8.0	11.8	13.5	11.0	14.9	6.2	np	9.6
Urban centres	11.8	9.8	14.8	17.0	16.8	np	..	np	12.8
Rural	15.3	13.2	17.3	20.9	13.4	12.9	–	26.3	16.0
Total	9.8	8.9	13.8	14.9	11.7	13.4	6.2	19.2	11.2
2013									
Capital city	7.5	7.3	13.5	12.0	11.2	15.1	9.7	np	9.6
Urban centres	11.0	8.9	12.7	12.3	np	11.9	na	np	11.4
Rural	14.0	12.9	19.9	20.7	12.4	15.7	–	21.8	16.0
Total	9.2	8.2	14.3	13.1	11.6	14.3	9.6	14.3	10.7
2009–2013									
Capital city	7.8	8.5	11.7	12.1	11.7	15.3	9.1	12.1	9.6
Urban centres	11.0	11.1	13.4	15.6	14.3	12.3	..	14.5	12.2
Rural	12.1	12.7	17.6	19.1	12.3	13.6	np	25.5	14.8
Total	9.1	9.4	13.3	13.4	11.9	14.0	9.1	17.6	10.9

- (a) Suicide deaths include ICD-10 codes X60–X84 and Y87.0. Care needs to be taken in interpreting data relating to suicide. See *ABS Causes of Death, 2013* (Cat. no. 3303.0) Explanatory Notes 87–93.
- (b) The total for each state and territory includes deaths registered to that state but which had a usual address which was undefined, overseas, of no fixed abode or off-shore and migratory. Such 'special purpose' Statistical Area 2s are only included in the state total.
- (c) The Australian total includes the 'Other Territories' — Jervis Bay, Christmas Island and the Cocos (Keeling) Islands.
- (d) For single year data prior to 2006, the categories were as follows: 'capital city' comprises capital city statistical divisions; 'other urban' comprises centres with more than 20 000 people; 'rural' comprises all areas except capital cities and other urban. 'Other urban' comprises statistical local areas with 50 per cent or greater of their 2001 census enumerated population contained in urban centres, based on Australian Standard Geographical Classification (ASGC) 2001 boundaries. 'Rural' comprises statistical local areas with 50 per cent or greater of their 2001 census enumerated population contained in rural areas. Changes in the population within geographical areas may not be reflected in the rates provided. There is some risk that urban growth areas have been classified as rural as the geography was based on the population in those areas in 2001. Therefore, analysis of data should be undertaken with caution.

Table 12A.84 Suicide deaths and suicide death rate, by area (a), (b), (c), (d), (e), (f), (g), (h), (i), (j)

	NSW	Vic	Qld	WA	SA	Tas (k)	ACT	NT	Aust (c)
(e)	For single year 2006, 2007 and 2008, the categories were derived as follows: 'capital cities' — comprising capital city statistical divisions, 'urban centres' — based on 'statistical districts' that are urban centres with population >25 000 people, excluding capital city statistical divisions, (three statistical districts cross state boundaries and have to be split across the relevant states/territories — Albury–Wodonga, Canberra–Queanbeyan and Gold Coast–Tweed); 'rural' — balance of state, that is all areas other than capital cities and urban centres.								
(f)	For the single years from 2009 and the five year sum and averages (2009–2013), the capital city, urban centres and rural groupings are based on the ABS' Significant Urban Areas classification (Cat. no. 1270.0.55.004). Capital cities are comprised of those Statistical Area 2s classified as capital cities. Urban centres are comprised of all Statistical Area 2s within a state which are classified as having or contributing to an urban area with a population of 10 000 or greater, excluding capital cities. Rural areas are those Statistical Area 2s which are not within a capital city or urban centre. For further information, see Cat. no. 1270.0.55.004 — Australian Statistical Geography Standard (ASGS): Volume 4 — Significant Urban Areas, Urban Centres and Localities, Section of State, July 2011. Some Significant Urban Areas cross state boundaries: Canberra – Queanbeyan (ACT/NSW); Albury – Wodonga (NSW/Vic); and Gold Coast – Tweed Heads (Qld/NSW). In these cases, deaths have been included in the Urban Centre category in the relevant state. The exception is Canberra – Queanbeyan: the Canberra portion forms the Capital City area for ACT, while the Queanbeyan portion has been included in the Urban Centres data for NSW.								
(g)	All causes of death data from 2006 onward are subject to a revisions process — once data for a reference year are 'final', they are no longer revised. Revised data for this year's report are: 2011 (final), 2012 (revised) and the data for 2013 are preliminary.								
(h)	For data from 2006, cells with small values have been randomly assigned to protect the confidentiality of individuals. As a result, some totals will not equal the sum of their components. Cells with a zero value have not been affected by confidentialisation.								
(i)	All footnotes and caveats, including this notice, must remain attached to data at all times.								
(j)	The total for each state and territory includes deaths registered to that state, but which had a usual address which was undefined, overseas, of no fixed abode or off-shore and migratory. Such 'special purpose' Statistical Area 2s are only included in the state total.								
(k)	The three criteria for this data tend to distort the Tasmanian picture due to the low level of urbanisation.								
(l)	Age-standardised death rates per 100 000 are standardised to Australian 30 June 2001 population. Rates for 2009–2013 and 2013 are based on 2013 revised substate estimated resident population data.								

.. Not applicable. – Nil or rounded to zero. **np** Not published.

Source: ABS (unpublished) *Causes of Death, Australia*, Cat. no. 3303.0.

Table 12A.85 Suicide deaths, by Indigenous status, 2009–2013 (a), (b), (c), (d), (e), (f), (g)

	<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (h)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total (i)</i>
<i>Number</i>										
Aboriginal and Torres Strait Islander	no.	101	np	191	157	37	np	np	115	601
Non-Indigenous (g)	no.	3 237	np	2 780	1 442	954	np	np	92	8 505
Total		3 338	np	2 971	1 599	991	np	np	207	9 106
<i>Suicide rate per 100 000 (i), (j), (k)</i>										
Aboriginal and Torres Strait Islander	no.	12.3	np	21.4	35.9	20.8	np	np	29.4	21.4
Non-Indigenous (g)	no.	9.0	np	12.8	12.5	11.7	np	np	11.6	10.9
<i>Variability bands (rate per 100 000)</i>										
Aboriginal and Torres Strait Islander	±	6.0	np	7.8	14.4	16.0	np	np	12.8	4.3
Non-Indigenous (k)	±	0.7	np	1.1	1.4	1.7	np	np	5.9	0.5

- (a) All causes of death data from 2006 onward are subject to a revisions process — once data for a reference year are 'final', they are no longer revised. Revised data for this year's report are: 2011 (final), 2012 (revised) and the data for 2013 are preliminary. See Explanatory Notes 52–54 and Technical Note, Causes of Death Revisions 2011 and 2012 in *Causes of Death, Australia, 2013* (Cat. no. 3303.0). See also Explanatory Notes 70–95 for further information on specific issues relating to 2013 data.
- (b) Data are based on State or Territory of usual residence.
- (c) Suicide deaths include ICD-10 codes X60–X84 and Y87.0. Care needs to be taken in interpreting data relating to suicide. See ABS Causes of Death, 2013 (Cat. no. 3303.0) Explanatory Notes 87–93.
- (d) Data are presented in a five-year aggregation (2009–2013) due to volatility of the small numbers involved.
- (e) All footnotes and caveats, including this notice, must remain attached to data at all times.
- (f) Data are reported for NSW, Qld, WA, SA and the NT only. Only these five states and territories have evidence of a sufficient level of Indigenous identification and sufficient numbers of Indigenous deaths to support mortality analysis. See Explanatory Notes 58–67 for further information on interpreting data relating to deaths of Indigenous persons.
- (g) Deaths where the Indigenous status of the deceased was not stated are included with the non-Indigenous data.
- (h) Care should be taken when interpreting deaths data for Queensland as they were affected by changes in the timeliness of birth and death registrations. Queensland deaths data for 2010 were adjusted to minimise the impact of late registration of deaths on mortality indicators. See Retrospective deaths by Causes of Death, Queensland, 2010 (Technical Note) in *Causes of Death, Australia, 2010* (Cat. no. 3303.0) for a more detailed explanation.
- (i) Total includes data for NSW, Queensland, WA, SA and the NT only.
- (j) Age standardised death rates. Deaths are per 100 000 of estimated mid-year population. The current ABS standard population is all persons in the Australian population at 30 June 2001. See Glossary of *Causes of Death, Australia, 2013* (Cat. no. 3303.0) for further information.
- (k) The Aboriginal and Torres Strait Islander population denominator used for calculating death rates in this table is from Estimates and Projections, Aboriginal and Torres Strait Islander Australians, 2001 to 2026 (Cat. no. 3238.0), Series B. These are 2011-census-based population projections. The non-Indigenous denominator has been derived by subtracting the Indigenous population projections from the total persons 2011-census-based population estimates.

Table 12A.85 **Suicide deaths, by Indigenous status, 2009–2013 (a), (b), (c), (d), (e), (f), (g)**

<i>Unit</i>	<i>NSW</i>	<i>Vic</i>	<i>Qld (h)</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total (i)</i>
-------------	------------	------------	----------------	-----------	-----------	------------	------------	-----------	------------------

np Not published.

Source: ABS (unpublished) *Causes of Death, Australia*, Cat. no. 3303.0; ABS (2015) *Estimates and Projections, Aboriginal and Torres Strait Islander Australians*, 2001 to 2026, Cat. no. 3238.0.

TABLE 12A.86

Table 12A.86 **Age-standardised proportions of adults by health risk factors and mental illness status, 2011-12 (a), (b), (c)**

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (d)	Total
<i>Overweight/obese</i>									
People with mental or behavioural problems (e), (f)	64.7 ± 5.9	66.3 ± 5.9	65.4 ± 6.4	73.6 ± 5.3	69.9 ± 6.2	65.4 ± 8.4	61.8 ± 6.7	68.1 ± 13.3	67.0 ± 2.5
People without mental or behavioural problems	60.2 ± 2.4	61.0 ± 2.3	65.3 ± 2.5	64.7 ± 2.3	65.5 ± 2.5	63.5 ± 2.7	63.9 ± 4.2	63.1 ± 4.5	62.4 ± 1.2
All people	61.1 ± 2.1	61.9 ± 2.2	65.4 ± 2.3	66.0 ± 2.1	66.1 ± 2.2	64.1 ± 2.5	63.6 ± 3.9	63.7 ± 3.9	63.2 ± 1.1
<i>Daily smoker</i>									
People with mental or behavioural problems (e), (f)	23.6 ± 4.5	28.9 ± 6.4	25.7 ± 4.6	26.0 ± 5.8	26.7 ± 4.9	32.4 ± 5.7	20.0 ± 5.6	29.1 ± 10.1	26.1 ± 2.4
People without mental or behavioural problems	13.4 ± 1.5	14.7 ± 1.7	15.8 ± 2.1	15.0 ± 1.9	15.5 ± 2.1	21.5 ± 2.3	11.7 ± 2.7	21.8 ± 3.0	14.7 ± 0.8
All people	14.8 ± 1.4	16.8 ± 1.8	17.5 ± 1.9	16.9 ± 2.1	17.4 ± 1.8	23.2 ± 2.2	13.4 ± 2.6	22.6 ± 2.8	16.5 ± 0.7
<i>At risk of long term harm from alcohol (g)</i>									
People with mental or behavioural problems (e), (f)	21.7 ± 4.9	20.5 ± 3.8	20.4 ± 4.4	25.1 ± 4.7	17.8 ± 5.3	22.2 ± 6.0	22.4 ± 6.9	19.8 ± 9.0	21.3 ± 2.0
People without mental or behavioural problems	17.9 ± 1.7	17.0 ± 1.8	19.8 ± 2.0	25.2 ± 2.4	18.2 ± 1.9	23.0 ± 2.7	20.6 ± 2.3	25.0 ± 3.8	19.0 ± 0.9
All people	18.5 ± 1.5	17.5 ± 1.6	19.9 ± 1.8	25.3 ± 2.1	18.2 ± 1.8	22.8 ± 2.4	21.0 ± 2.4	24.2 ± 3.5	19.4 ± 0.8

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) Numerators — number of adults (aged 18 years or over) who are overweight or obese or a daily smoker or at risk of long term harm from alcohol (by mental health status). Denominators — number of adults (aged 18 years or over) in the population (by mental health status).

(c) As State and Territory comparisons are affected by age, estimates have been age standardised to the 2001 estimated resident population.

(d) Data for NT should be interpreted with caution as the *Australian Health Survey (2011-12 NHS component)* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

(e) People with a mental or behavioural condition are defined as having a current self-reported mental and behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.

Table 12A.86 **Age-standardised proportions of adults by health risk factors and mental illness status, 2011-12 (a), (b), (c)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (d)</i>	<i>Total</i>
--	------------	------------	------------	-----------	-----------	------------	------------	---------------	--------------

(f) Includes organic mental conditions, alcohol and drug conditions, mood conditions and other mental and behavioural conditions.

(g) 'At risk of long term harm' is based on the 2009 National Health and Medical Research Council (NHMRC) guidelines.

Source: ABS (unpublished) *Australian Health Survey 2011-13 (2011-12 NHS component)*, Cat. no. 4364.0.

TABLE 12A.87

Table 12A.87 **Age-standardised proportions of adults by long-term health conditions and mental illness status, 2011-
(a), (b), (c)**

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (d)	Total
<i>Cancer</i>									
People with mental or behavioural problems (e), (f)	3.0* ± 1.9	3.9* ± 2.2	3.2* ± 1.8	6.6 ± 3.0	1.5* ± 1.2	4.4* ± 2.7	3.4* ± 2.7	13.4* ± 7.3	3.5 ± 0.9
People without mental or behavioural problems	1.1 ± 0.4	1.6 ± 0.5	2.1 ± 0.6	1.5 ± 0.5	1.4 ± 0.5	1.6 ± 0.7	2.1* ± 1.0	1.7* ± 0.9	1.5 ± 0.2
All people	1.4 ± 0.4	2.0 ± 0.6	2.2 ± 0.6	2.3 ± 0.6	1.4 ± 0.5	2.2 ± 0.7	2.3 ± 1.0	2.8 ± 1.3	1.8 ± 0.2
<i>Diabetes</i>									
People with mental or behavioural problems (e), (f)	7.5 ± 2.9	5.9 ± 2.1	7.2 ± 2.5	6.4 ± 2.5	6.1 ± 2.5	6.4 ± 3.0	3.4* ± 2.5	10.5* ± 9.9	6.6 ± 1.1
People without mental or behavioural problems	5.5 ± 0.8	5.0 ± 0.8	4.7 ± 1.0	5.5 ± 1.2	5.7 ± 1.2	5.3 ± 1.3	5.8 ± 1.6	6.9 ± 2.6	5.3 ± 0.4
All people	5.8 ± 0.8	5.2 ± 0.8	5.2 ± 1.0	5.6 ± 1.1	5.8 ± 1.0	5.6 ± 1.2	5.4 ± 1.3	7.5 ± 2.1	5.5 ± 0.4
<i>Arthritis</i>									
People with mental or behavioural problems (e), (f)	29.1 ± 5.5	25.4 ± 4.1	25.1 ± 4.2	24.2 ± 5.1	26.3 ± 4.2	29.4 ± 4.9	31.9 ± 4.6	26.2 ± 11.5	26.9 ± 2.4
People without mental or behavioural problems	17.0 ± 1.3	15.9 ± 1.4	16.1 ± 1.6	17.3 ± 1.8	17.7 ± 1.9	19.8 ± 2.0	16.8 ± 2.5	14.1 ± 3.2	16.7 ± 0.7
All people	18.9 ± 1.1	17.4 ± 1.4	17.6 ± 1.6	18.6 ± 1.8	19.2 ± 1.9	21.6 ± 2.0	19.3 ± 2.5	15.5 ± 3.2	18.3 ± 0.7
<i>Cardiovascular disease</i>									
People with mental or behavioural problems (e), (f)	7.6 ± 2.4	9.6 ± 2.9	12.9 ± 3.4	8.2 ± 2.8	9.9 ± 2.6	11.9 ± 4.0	15.6 ± 4.4	14.5* ± 10.7	9.5 ± 1.2
People without mental or behavioural problems	5.8 ± 0.9	4.3 ± 0.8	5.8 ± 0.9	4.6 ± 0.9	4.9 ± 1.0	6.2 ± 1.2	5.6 ± 1.4	4.3 ± 1.9	5.2 ± 0.4
All people	6.1 ± 0.9	5.1 ± 0.8	6.8 ± 0.9	5.2 ± 0.9	5.7 ± 0.9	7.0 ± 1.1	7.4 ± 1.4	5.5 ± 2.4	5.9 ± 0.4

Table 12A.87 **Age-standardised proportions of adults by long-term health conditions and mental illness status, 2011-
(a), (b), (c)**

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (d)	Total
<i>Asthma</i>									
People with mental or behavioural problems (e), (f)	16.5 ± 3.9	18.8 ± 4.0	15.5 ± 3.4	16.0 ± 5.0	14.7 ± 4.3	17.0 ± 5.6	18.8 ± 5.7	17.6* ± 10.9	16.7 ± 1.8
People without mental or behavioural problems	8.8 ± 1.5	9.4 ± 1.2	9.3 ± 1.3	9.7 ± 1.5	9.8 ± 1.7	9.8 ± 2.0	8.3 ± 1.7	7.7 ± 2.5	9.2 ± 0.7
All people	9.9 ± 1.5	10.8 ± 1.2	10.3 ± 1.3	10.8 ± 1.4	10.7 ± 1.6	11.1 ± 2.0	10.0 ± 1.7	8.7 ± 2.3	10.4 ± 0.7

- (a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent). A '**' indicates a RSE of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution. Estimates with RSEs greater than 50 per cent are considered too unreliable for general use.
- (b) Numerators — number of adults who are have the specific long-term health condition for example, cancer (by mental health status). Denominators — number of adults in the population (by mental health status).
- (c) As State and Territory comparisons are affected by age, estimates have been age standardised to the 2001 estimated resident population.
- (d) Data for NT should be interpreted with caution as the *Australian Health Survey (2011-12 NHS component)* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.
- (e) People with a mental or behavioural condition are defined as having a current self-reported mental and behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.
- (f) Includes organic mental conditions, alcohol and drug conditions, mood conditions and other mental and behavioural conditions.

Source: ABS (unpublished) *Australian Health Survey 2011-13 (2011-12 NHS component)*, Cat. no. 4364.0.

TABLE 12A.88

Table 12A.88 **Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2011-12 (per cent) (a), (b), (c)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (d)</i>	<i>Total</i>
People aged 16–64 years who are employed									
People with mental or behavioural problems (e), (f)	65.2 ± 7.7	59.4 ± 6.4	57.7 ± 6.7	65.0 ± 5.9	61.2 ± 7.2	51.6 ± 8.7	72.5 ± 8.2	63.2 ± 10.3	61.7 ± 3.1
People without mental or behavioural problems	78.7 ± 1.7	81.0 ± 1.8	81.8 ± 2.0	81.5 ± 1.9	78.7 ± 2.4	76.1 ± 2.9	85.6 ± 2.1	84.8 ± 3.1	80.3 ± 0.9
All people	76.6 ± 2.0	77.7 ± 1.8	77.7 ± 2.1	78.7 ± 1.9	76.0 ± 2.5	71.8 ± 3.2	83.4 ± 2.3	81.9 ± 3.1	77.4 ± 1.0
People aged 16–64 years who are unemployed									
People with mental or behavioural problems (e), (f)	4.3* ± 2.7	6.0* ± 3.2	9.6 ± 3.3	5.5* ± 3.6	7.0* ± 3.6	8.7* ± 4.6	2.9* ± 2.7	5.6** ± 7.0	6.3 ± 1.4
People without mental or behavioural problems	2.8 ± 0.9	2.8 ± 1.0	3.2 ± 1.0	2.8 ± 1.1	3.8 ± 1.3	3.6 ± 1.2	1.4* ± 0.9	2.0* ± 1.2	3.0 ± 0.4
All people	3.0 ± 0.8	3.4 ± 1.0	4.3 ± 1.0	3.3 ± 1.0	4.3 ± 1.2	4.4 ± 1.3	1.8* ± 0.9	2.4* ± 1.2	3.5 ± 0.4
People aged 16–64 years who are in the labour force									
People with mental or behavioural problems (e), (f)	69.5 ± 7.3	65.4 ± 6.5	67.3 ± 6.6	70.6 ± 6.2	68.2 ± 7.2	60.3 ± 8.7	75.4 ± 7.9	68.7 ± 11.2	68.0 ± 3.2
People without mental or behavioural problems	81.5 ± 1.6	83.8 ± 1.7	85.1 ± 1.8	84.4 ± 1.8	82.5 ± 2.1	79.6 ± 2.9	87.0 ± 2.0	86.8 ± 2.7	83.3 ± 0.9
All people	79.7 ± 1.8	81.1 ± 1.7	82.0 ± 1.8	81.9 ± 1.6	80.3 ± 2.2	76.2 ± 3.0	85.1 ± 2.0	84.3 ± 2.7	80.8 ± 0.9
People aged 16–64 years who are not in the labour force									
People with mental or behavioural problems (e), (f)	30.5 ± 7.3	34.6 ± 6.5	32.7 ± 6.6	29.4 ± 6.2	31.8 ± 7.2	39.7 ± 8.8	24.6 ± 7.9	31.3 ± 11.1	32.0 ± 3.2
People without mental or behavioural problems	18.5 ± 1.6	16.2 ± 1.7	14.9 ± 1.8	15.6 ± 1.8	17.5 ± 2.1	20.4 ± 2.9	13.0 ± 2.0	13.2 ± 2.7	16.7 ± 0.9
All people	20.3 ± 1.8	18.9 ± 1.7	18.0 ± 1.8	18.1 ± 1.6	19.7 ± 2.2	23.8 ± 3.0	14.9 ± 2.0	15.7 ± 2.7	19.2 ± 0.9

Table 12A.88 Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2011-12 (per cent) (a), (b), (c)

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (d)	Total
(a)	The rates reported in this table include 95 per cent confidence intervals (for example, X per cent \pm X per cent). A '**' indicates a RSE of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution. A '***' indicates a RSE of greater than 50 per cent. Estimates with RSEs greater than 50 per cent are considered too unreliable for general use.								
(b)	Numerators — number of people aged 16–64 years who are employed/unemployed/in the labour force/not in the labour force (by mental health status). Denominators — number of people aged 16–64 years in the population (by mental health status).								
(c)	As State and Territory comparisons are affected by age, estimates have been age standardised to the 2001 estimated resident population.								
(d)	Data for NT should be interpreted with caution as the <i>Australian Health Survey (2011-12 NHS component)</i> excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.								
(e)	People with a mental or behavioural condition are defined as having a current self-reported mental and behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.								
(f)	Includes organic mental conditions, alcohol and drug conditions, mood conditions and other mental and behavioural conditions.								
<i>Source:</i> ABS (unpublished) <i>Australian Health Survey 2011-13 (2011-12 NHS component)</i> , Cat. no. 4364.0.									

TABLE 12A.89

Table 12A.89 **Age-standardised proportion of the population aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status, 2011-12 (per cent) (a), (b), (c)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (d)</i>	<i>Total</i>
People with mental or behavioural problems (e), (f)	80.8 ± 9.1	80.4 ± 12.2	79.4 ± 8.3	70.9 ± 11.9	84.2 ± 10.2	74.8 ± 11.0	82.2 ± 12.3	55.2* ± 27.9	79.2 ± 4.2
People without mental or behavioural problems	93.2 ± 2.4	90.5 ± 2.7	87.0 ± 3.4	88.7 ± 3.7	85.5 ± 4.7	86.6 ± 4.8	97.2 ± 2.1	87.5 ± 5.4	90.2 ± 1.2
All people	91.8 ± 2.3	89.2 ± 2.8	85.8 ± 3.3	85.7 ± 4.0	85.4 ± 4.4	84.5 ± 4.3	94.9 ± 2.6	83.2 ± 6.1	88.7 ± 1.1

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent). A '*' indicates a RSE of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution.

(b) Numerators – number of people aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status. Denominators – number of people aged 16–30 years, by mental health status.

(c) As State and Territory comparisons are affected by age, estimates have been age standardised to the 2001 estimated resident population.

(d) Data for NT should be interpreted with caution as the *Australian Health Survey (2011-12 NHS component)* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

(e) People with a mental or behavioural condition are defined as having a current self-reported mental and behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.

(f) Includes organic mental conditions, alcohol and drug conditions, mood conditions and other mental and behavioural conditions.

Source: ABS (unpublished) *Australian Health Survey 2011-13 (2011-12 NHS component)*, Cat. no. 4364.0.

TABLE 12A.90

Table 12A.90 **Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2007–08 (per cent) (a), (b), (c)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (d)</i>	<i>Total</i>
People aged 16–64 years who are employed									
People with mental or behavioural problems (e), (f)	59.3 ± 6.2	68.2 ± 5.8	65.4 ± 6.9	70.8 ± 7.5	48.6 ± 6.9	55.7 ± 8.3	75.4 ± 5.9	57.2 ± 23.7	63.8 ± 3.2
People without mental or behavioural problems	78.0 ± 2.3	79.8 ± 2.0	79.0 ± 2.3	83.1 ± 2.3	79.3 ± 2.6	74.2 ± 3.1	85.9 ± 2.1	83.4 ± 11.1	79.4 ± 1.0
All people	75.6 ± 2.2	78.4 ± 1.8	77.0 ± 2.2	81.3 ± 2.4	75.1 ± 2.6	71.6 ± 3.1	84.5 ± 2.0	83.9 ± 8.8	77.3 ± 1.0
People aged 16–64 years who are unemployed									
People with mental or behavioural problems (e), (f)	7.2 ± 3.3	4.2 ± 2.0	4.2* ± 3.2	3.1* ± 2.5	8.7 ± 3.5	6.6* ± 5.9	3.6* ± 3.5	–	5.3 ± 1.2
People without mental or behavioural problems	2.4 ± 0.8	2.3 ± 0.8	2.9 ± 1.1	2.3 ± 1.1	3.1 ± 1.0	4.1 ± 2.0	np	np	2.5 ± 0.4
All people	3.1 ± 0.8	2.5 ± 0.7	3.1 ± 1.0	2.4 ± 1.0	3.9 ± 1.0	4.3 ± 1.7	np	np	2.9 ± 0.4
People aged 16–64 years who are in the labour force									
People with mental or behavioural problems (e), (f)	66.4 ± 5.7	72.4 ± 6.1	69.6 ± 6.2	73.9 ± 7.2	57.3 ± 7.2	62.3 ± 9.5	79.1 ± 5.9	57.2 ± 23.7	69.1 ± 2.8
People without mental or behavioural problems	80.4 ± 2.2	82.1 ± 2.0	81.9 ± 2.1	85.4 ± 2.1	82.4 ± 2.2	78.3 ± 2.8	87.4 ± 2.0	85.1 ± 10.5	81.9 ± 1.0
All people	78.7 ± 2.1	80.9 ± 1.8	80.1 ± 1.9	83.7 ± 2.2	79.0 ± 2.1	75.9 ± 3.1	86.2 ± 1.9	85.6 ± 8.1	80.2 ± 1.0
People aged 16–64 years who are not in the labour force									
People with mental or behavioural problems (e), (f)	33.6 ± 5.7	27.6 ± 6.1	30.4 ± 6.2	26.1 ± 7.2	42.7 ± 7.2	37.7 ± 9.5	np	np	30.9 ± 2.8
People without mental or behavioural problems	19.6 ± 2.2	17.9 ± 2.0	18.1 ± 2.1	14.6 ± 2.1	17.6 ± 2.2	21.7 ± 2.8	np	np	18.1 ± 1.0
All people	21.3 ± 2.1	19.1 ± 1.8	19.9 ± 1.9	16.3 ± 2.2	21.0 ± 2.1	24.1 ± 3.1	13.8 ± 1.9	14.4* ± 8.1	19.8 ± 1.0

Table 12A.90 **Age-standardised proportion of people aged 16–64 years who are employed, by mental illness status, 2007-08 (per cent) (a), (b), (c)**

	NSW	Vic	Qld	WA	SA	Tas	ACT	NT (d)	Total
(a)	The rates reported in this table include 95 per cent confidence intervals (for example, X per cent \pm X per cent). A ^{†††} indicates a relative standard error (RSE) of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution. A ^{††††} indicates a RSE of greater than 50 per cent. Estimates with RSEs greater than 50 per cent are considered too unreliable for general use.								
(b)	Numerators — number of people aged 16–64 years who are employed/unemployed/in the labour force/not in the labour force (by mental health status). Denominators — number of people aged 16–64 years in the population (by mental health status).								
(c)	As State and Territory comparisons are affected by age, estimates have been age standardised to the 2001 estimated resident population.								
(d)	Data for NT should be interpreted with caution as the <i>National Health Survey 2007-08</i> excluded very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.								
(e)	People with a mental or behavioural condition are defined as having a current self-reported mental and behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.								
(f)	Includes organic mental conditions, alcohol and drug conditions, mood conditions and other mental and behavioural conditions. – Nil or rounded to zero. np Not published.								

Source: ABS (unpublished) *National Health Survey 2007-08*, Cat. no. 4364.0.

TABLE 12A.91

Table 12A.91 **Population aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status, 2007-08 (per cent) (a), (b), (c), (d)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Total</i>
People with a mental illness (e), (f)	78.1 ± 11.8	80.7 ± 10.0	83.6 ± 11.3	84.0 ± 10.6	66.1 ± 9.8	63.0 ± 17.5	88.3 ± 7.2	np	79.6 ± 5.7
People without a mental illness	89.8 ± 2.9	91.8 ± 2.7	86.9 ± 4.4	89.8 ± 3.9	89.1 ± 3.1	87.0 ± 5.1	94.7 ± 2.3	88.0 ± 24.9	89.7 ± 1.7
All people	88.4 ± 2.8	90.3 ± 2.6	86.4 ± 3.9	88.9 ± 4.0	85.9 ± 3.3	83.3 ± 6.0	93.8 ± 2.1	88.0 ± 24.9	88.4 ± 1.6

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent). A '*' indicates a relative standard error (RSE) of between 25 per cent and 50 per cent. Estimates with RSEs greater than 25 per cent should be used with caution.

(b) Numerators – number of people aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (full or part-time), by mental health status. Denominators – number of people aged 16–30 years, by mental health status.

(c) As State and Territory comparisons are affected by age, estimates have been age standardised to the 2001 estimated resident population.

(d) Data for NT should be interpreted with caution as the *National Health Survey 2007-08* excluded very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

(d) People with a mental or behavioural condition are defined as having a current self-reported mental and behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.

(e) Includes organic mental conditions, alcohol and drug conditions, mood conditions and other mental and behavioural conditions.

np Not published.

Source: ABS (unpublished) *2007-08 National Health Survey*, Cat. no. 4364.0.

TABLE 12A.92

Table 12A.92 **Labour force and employment participation among adults aged 16–64 years, by mental disorder status, 2007 (per cent) (a)**

	<i>Employed (b)</i>			<i>Unemployed (b)</i>	<i>In labour force</i>	<i>Not in the labour force</i>
	<i>Full-time</i>	<i>Part-time</i>	<i>Total</i>			
Any 12-month mental disorder (c)						
Anxiety disorders	59.9 ± 5.5	35.4 ± 5.3	95.3 ± 2.0	4.7 ± 2.0	71.0 ± 3.4	29.0 ± 3.4
Affective disorders	57.4 ± 6.8	32.6 ± 7.0	90.0 ± 4.3	10.0 ± 4.3	69.8 ± 4.3	30.2 ± 4.3
Substance use disorders	62.3 ± 6.9	30.8 ± 7.3	93.1 ± 3.3	6.9 ± 3.3	83.0 ± 5.4	17.0 ± 5.4
Any 12-month mental disorder (c), (d)	59.8 ± 4.7	34.7 ± 4.4	94.5 ± 1.7	5.5 ± 1.7	73.6 ± 2.7	26.4 ± 2.7
Lifetime mental disorder, with no 12-month symptoms (e)	68.7 ± 3.8	27.4 ± 3.7	96.1 ± 1.7	3.9 ± 1.7	80.9 ± 2.4	19.1 ± 2.4
No lifetime mental disorder (f)	63.7 ± 2.3	33.1 ± 2.3	96.8 ± 0.9	3.2 ± 0.9	78.4 ± 1.6	21.6 ± 1.6

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) The employed and unemployed rates are as a proportion of those in the labour force.

(c) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(d) A person may have more than one mental disorder. Therefore the components may not add to the total.

(e) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(f) People who did not meet criteria for diagnosis of a lifetime mental disorder.

Source: ABS (unpublished) 2007 Survey of Mental Health and Wellbeing, Cat. no. 4326.0.

Table 12A.93 **Education, training and employment participation among adults aged 16–30 years, by mental disorder status, 2007 (per cent) (a), (b)**

	Studying (c)	Not studying			Total
		Employed	Unemployed/Not in the labour force	Total	
Any 12-month mental disorder (d)	42.0 ± 4.9	44.3 ± 5.0	13.7 ± 3.0	58.0 ± 4.9	100.0
Lifetime mental disorder, with no 12-month symptoms (e)	29.5 ± 6.6	55.9 ± 7.3	np	70.5 ± 6.6	100.0
No lifetime mental disorder (f)	51.6 ± 3.8	39.2 ± 3.3	9.2 ± 2.2	48.4 ± 3.8	100.0

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(b) Estimates with RSEs greater than 25 per cent are considered unreliable. These estimates are not published.

(c) Includes people studying full-time and part-time and people still at school.

(d) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(e) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(f) People who did not meet criteria for diagnosis of a lifetime mental disorder.

np Not published.

Source: ABS (unpublished) 2007 Survey of Mental Health and Wellbeing, Cat. no. 4326.0.

Table 12A.94 Labour force and employment participation among adults aged 16–30 years, by mental disorder status, 2007 (per cent) (a)

	<i>Employed (b)</i>	<i>Unemployed (b)</i>	<i>Not in the labour force</i>
Any 12-month mental disorder (c)	92.1 ± 3.2	7.9 ± 3.2	19.2 ± 3.4
Lifetime mental disorder, with no 12-month symptoms (d)	92.2 ± 9.0	np	17.6 ± 6.2
No lifetime mental disorder (e)	93.6 ± 1.9	6.4 ± 1.9	22.1 ± 2.9

(a) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent). Estimates with RSEs greater than 25 per cent are not published.

(b) The employed and unemployed rates are as a proportion of those in the labour force.

(c) People who met criteria for diagnosis of a lifetime mental disorder (with hierarchy) and had symptoms in the 12 months prior to interview.

(d) People who had experienced a mental disorder at some point in their life, but who did not have symptoms in the previous 12 months.

(e) People who did not meet criteria for diagnosis of a lifetime mental disorder.

np Not published.

Source: ABS (unpublished) *2007 Survey of Mental Health and Wellbeing*, Cat. no. 4326.0.

Table 12A.95 Proportion of people who had face-to-face contact with family or friends living outside the household in the last week, by mental illness status, 2014 (per cent) (a), (b), (c), (d)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT (e)</i>	<i>Total</i>
People with mental illness a	75.4 ± 7.3	79.0 ± 5.3	72.5 ± 5.2	77.0 ± 6.6	81.7 ± 8.1	78.1 ± 5.2	76.3 ± 5.9	54.8 ± 11.5	76.5 ± 3.1
People without mental illness a	75.1 ± 2.8	77.9 ± 3.1	75.6 ± 3.3	77.4 ± 3.1	85.1 ± 2.1	86.5 ± 2.8	75.8 ± 2.6	69.6 ± 4.6	77.1 ± 1.4
All people	75.2 ± 2.4	78.2 ± 2.4	75.0 ± 2.9	77.3 ± 3.0	84.2 ± 2.0	84.6 ± 2.2	75.6 ± 2.7	68.2 ± 4.5	77.0 ± 1.2

(a) People with a mental illness is a self-reported data item. The data item refers to clinically recognised emotional and behavioural disorders, and perceived mental health problems such as feeling depressed, feeling anxious, stress and sadness.

(b) People who had face-to-face contact with family or friends living outside the household in the last week refers to those who reported having contact everyday or at least weekly.

(c) The rates reported in this table include 95 per cent confidence intervals (for example, X per cent ± X per cent).

(d) Cells in this table may have been randomly adjusted to avoid the release of confidential data.

(e) Data for NT should be interpreted with caution as the *General Social Survey 2014* excluded discrete Aboriginal and Torres Strait Islander communities and very remote areas, which comprise around 25 per cent of the estimated resident population of the NT.

Source: ABS (unpublished) *General Social Survey 2014*, Cat. no. 4159.0 (derived using Table Builder product).

Table 12A.96 Clinical outcomes of people receiving various types of mental health care provided by State and Territory public mental health services (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
<i>2007-08</i>									
Group A: People discharged from hospital (g)									
Significant improvement	75.6	76.1	71.3	74.8	66.7	72.2	np	np	73.3
No significant change	20.2	20.5	22.7	20.4	29.0	21.6	np	np	22.1
Significant deterioration	4.2	3.5	6.0	4.8	4.4	6.2	np	np	4.6
Group B: People discharged from community ambulatory care (h)									
Significant improvement	55.6	53.6	55.1	47.7	47.4	47.0	np	np	53.3
No significant change	42.0	42.5	38.9	44.7	47.0	46.4	np	np	41.7
Significant deterioration	2.4	3.9	6.0	7.6	5.6	6.6	np	np	5.0
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	24.5	27.9	29.3	28.5	24.9	27.7	np	23.3	27.1
No significant change	60.7	58.0	52.2	56.4	58.7	51.8	np	56.4	56.8
Significant deterioration	14.8	14.0	18.5	15.1	16.4	20.6	np	20.3	16.1
<i>2008-09</i>									
Group A: People discharged from hospital (g)									
Significant improvement	74.7	76.2	73.9	75.8	70.3	76.9	np	np	74.7
No significant change	21.2	20.1	21.2	20.2	25.4	20.2	np	np	21.2
Significant deterioration	4.0	3.7	4.9	4.0	4.4	2.8	np	np	4.0
Group B: People discharged from community ambulatory care (h)									
Significant improvement	55.9	50.3	57.8	52.9	46.3	45.9	np	np	52.6
No significant change	41.6	44.2	36.3	39.8	48.9	46.9	np	np	42.1
Significant deterioration	2.6	5.5	5.9	7.2	4.8	7.2	np	np	5.3
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	23.6	29.4	29.4	25.6	27.1	27.2	np	27.2	27.3
No significant change	61.9	56.2	53.3	58.7	57.7	58.0	np	49.9	57.2
Significant deterioration	14.5	14.4	17.3	15.7	15.2	14.7	np	23.0	15.5
<i>2009-10</i>									
Group A: People discharged from hospital (g)									
Significant improvement	68.7	73.5	74.1	72.9	70.0	77.2	np	np	71.7
No significant change	26.2	22.6	21.4	22.5	26.0	19.9	np	np	23.9
Significant deterioration	5.1	3.9	4.5	4.6	4.0	2.8	np	np	4.4
Group B: People discharged from community ambulatory care (h)									
Significant improvement	54.6	50.0	58.3	52.7	47.7	47.4	np	np	52.0
No significant change	42.1	43.8	35.7	42.3	48.2	48.5	np	np	42.6
Significant deterioration	3.3	6.1	5.9	5.0	4.0	4.1	np	np	5.4
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	22.6	28.3	31.9	27.2	25.2	27.4	18.5	25.5	27.3
No significant change	61.8	56.8	52.7	58.2	58.7	56.6	68.7	52.0	57.4
Significant deterioration	15.5	14.9	15.4	14.5	16.1	15.9	12.8	22.5	15.3

Table 12A.96 Clinical outcomes of people receiving various types of mental health care provided by State and Territory public mental health services (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
2010-11									
Group A: People discharged from hospital (g)									
Significant improvement	69.4	73.5	73.8	74.7	72.2	75.6	np	77.0	72.5
No significant change	25.1	22.8	20.1	21.6	24.1	20.1	np	19.5	23.1
Significant deterioration	5.4	3.7	6.2	3.7	3.8	4.3	np	3.5	4.5
Group B: People discharged from community ambulatory care (h)									
Significant improvement	56.6	45.5	59.2	51.7	46.0	52.7	np	np	50.0
No significant change	40.5	43.8	35.5	42.4	49.6	43.9	np	np	42.2
Significant deterioration	2.9	10.7	5.3	5.8	4.3	3.4	np	np	7.7
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	22.8	27.4	30.6	24.7	24.6	25.9	18.7	28.5	26.4
No significant change	62.2	57.3	53.5	59.3	61.1	57.3	67.8	50.3	58.1
Significant deterioration	15.0	15.3	15.9	16.0	14.3	16.8	13.5	21.2	15.4
2011-12									
Group A: People discharged from hospital (g)									
Significant improvement	68.1	na	73.4	72.1	71.3	73.0	np	77.6	70.8
No significant change	27.0	na	19.7	22.8	24.7	22.1	np	16.1	24.0
Significant deterioration	4.9	na	6.9	5.1	4.0	4.9	np	6.3	5.2
Group B: People discharged from community ambulatory care (h)									
Significant improvement	54.3	na	54.5	45.7	47.1	43.2	np	np	51.5
No significant change	42.4	na	39.5	48.7	48.8	51.7	np	np	43.7
Significant deterioration	3.3	na	5.9	5.6	4.0	5.1	np	np	4.8
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	23.0	na	30.4	24.6	23.8	27.5	29.0	27.4	26.0
No significant change	61.1	na	54.0	60.4	60.9	50.8	56.5	53.5	58.3
Significant deterioration	15.8	na	15.6	15.0	15.3	21.6	14.5	19.2	15.7
2012-13									
Group A: People discharged from hospital (g)									
Significant improvement	70.0	na	72.7	74.3	72.6	76.4	np	77.3	72.1
No significant change	25.6	na	20.7	21.5	22.6	19.5	np	16.0	22.9
Significant deterioration	4.4	na	6.6	4.2	4.8	4.1	np	6.7	5.0
Group B: People discharged from community ambulatory care (h)									
Significant improvement	51.7	na	53.7	47.3	42.9	51.3	np	np	50.8
No significant change	45.3	na	40.6	48.1	52.6	45.4	np	np	44.6
Significant deterioration	3.0	na	5.6	4.6	4.5	3.3	np	np	4.6
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	22.3	na	31.3	24.4	23.5	21.2	22.8	29.9	26.1
No significant change	61.8	na	53.5	59.3	61.2	59.9	61.9	51.9	58.1
Significant deterioration	15.9	na	15.2	16.3	15.3	18.9	15.2	18.2	15.7

Table 12A.96 Clinical outcomes of people receiving various types of mental health care provided by State and Territory public mental health services (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
<i>2013-14</i>									
Group A: People discharged from hospital (g)									
Significant improvement	69.1	73.3	74.8	75.6	68.9	75.5	39.1	77.5	72.4
No significant change	26.1	21.5	19.8	21.1	26.1	20.0	54.7	19.5	22.8
Significant deterioration	4.8	5.2	5.4	3.3	5.0	4.5	6.2	3.1	4.8
Group B: People discharged from community ambulatory care (h)									
Significant improvement	50.1	45.2	54.6	50.7	44.3	39.3	np	np	48.4
No significant change	45.9	48.0	38.9	44.9	51.6	57.0	np	np	45.7
Significant deterioration	4.0	6.8	6.5	4.4	4.1	3.6	np	np	5.9
Group C: People in ongoing community ambulatory care (i)									
Significant improvement	22.4	28.8	27.9	26.6	26.9	26.6	26.9	26.8	26.6
No significant change	62.2	55.1	54.1	58.1	60.0	55.8	53.1	49.2	57.2
Significant deterioration	15.4	16.1	18.0	15.3	13.1	17.7	20.1	24.0	16.2

- (a) These data were prepared by the Australian Mental Health Outcomes and Classification Network, using data submitted by State and Territory governments to the Australian Government Department of Health. Assessment of clinical outcomes is based on the changes reported in a consumer's score on a rating scale known as the Health of the Nation Outcomes Scale (HoNOS), or in the case of children and adolescent consumers, the Health of the Nation Outcome Scales for Children and Adolescents (HoNOSCA). Developed originally in England in the 1990s, these ratings scales comprise standard items that are rated by a clinician to measure the severity of the consumer's symptoms or disability across a range of domains (for example, depressed mood, hallucinations, substance use, suicidality, overactivity, activities of daily living, cognitive impairment). The HoNOS/HoNOSCA form part of small suite of standardised rating scales used to monitor outcomes across state and territory public sector mental health services and private hospitals with a specialised psychiatric unit. To be considered valid, HoNOS, or the HoNOSCA data needs to be completed correctly (a specified minimum number of items completed) and have a "matching pair" — that is, a beginning and end rating are needed to enable an outcome score to be determined.
- (b) Proportions may not add to 100 per cent due to rounding.
- (c) For all consumer groups, outcome scores for each episode are classified as either 'significant improvement', 'significant deterioration' or 'no significant change', based on Effect Size. Effect size is a statistic used to assess the magnitude of a treatment effect. It is based on the ratio of the difference between pre- and post- scores to the standard deviation of the pre- score. As a rule of thumb, effect sizes of 0.2 are considered small, 0.5 considered medium and 0.8 considered large. Based on this rule, a medium effect size of 0.5 was used to assign outcome scores to the three outcome categories. Thus individual episodes were classified as either: 'significant improvement' if the Effect Size index was greater than or equal to positive 0.5; 'significant deterioration' if the Effect Size index was less than or equal to negative 0.5; or 'no change' if the index was between -0.5 and 0.5.
- (d) Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. All national averages for 2011-12 and 2012-13 exclude Victoria.
- (e) Industrial action in Tasmania has limited the available data quality and quantity of data for 2011-12 and 2012-13.

Table 12A.96 Clinical outcomes of people receiving various types of mental health care provided by State and Territory public mental health services (per cent) (a), (b), (c)

	<i>NSW</i>	<i>Vic (d)</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas (e)</i>	<i>ACT (f)</i>	<i>NT (f)</i>	<i>Aust (d)</i>
(f)	Some data for the ACT and the NT are np (not published) due to insufficient observations. The number of observations of consumer outcomes for some care types is too low to publish because conclusions based on such low numbers are known to have high levels of unreliability. For the purposes of this indicator, the threshold for the minimum number of observations to be reached was set at 200.								
(g)	Group A covers people who received a discrete episode of inpatient care within a state/territory designated psychiatric inpatient unit during the reference year. The defining characteristic of the group is that the episode of inpatient care commenced, and was completed, within the year. Outcome scores were calculated as the difference between the total score recorded at admission and discharge. The analysis excludes episodes where length of stay was three days or less because it is not meaningful to compare admission and discharge ratings for short duration episodes.								
(h)	Group B covers people who received relatively short term community care from a state/territory mental health service during the reference year. The defining characteristic of the group is that the episode of community care commenced, and was completed, within the year. Outcome scores were calculated as the difference between the total score recorded at admission to, and discharge from, community care. A subgroup of people whose episode of community care completed because they were admitted to hospital is not included in this analysis.								
(i)	Group C covers people receiving relatively long term community care from a state/territory mental health service. It includes people who were receiving care for the whole of the reference year, and those who commenced community care sometime after 1 July who continued under care for the rest of the year. The defining characteristic of the group is that all remained in ongoing care when the year ended (30 June). Outcome scores were calculated as the difference between the total score recorded on the first occasion rated and the last occasion rated in the year.								

na Not available. **np** Not published.

Source: AIHW (unpublished) from data provided by the Australian Mental Health Outcomes and Classification Network.

Table 12A.97 People who received mental health care provided by State and Territory public mental health services and who significantly improved, by service type and age group (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
<i>2007-08</i>									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	61.7	72.3	61.8	np	np	np	np	np	63.2
Aged 18–64 years	77.1	78.3	72.5	78.0	68.4	72.8	np	np	75.0
Aged 65 years or over	68.8	67.0	67.2	58.3	69.8	np	np	np	66.4
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	59.4	53.9	59.8	np	41.1	np	np	np	53.7
Aged 18–64 years	55.0	56.0	55.6	46.6	57.1	np	np	np	55.3
Aged 65 years or over	52.0	49.3	47.4	42.6	np	np	np	np	47.8
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	36.3	37.7	41.1	39.8	28.7	np	np	np	35.8
Aged 18–64 years	23.2	26.6	27.3	28.4	23.0	28.4	np	23.0	25.8
Aged 65 years or over	23.1	26.2	26.3	20.7	22.0	np	np	np	23.7
<i>2008-09</i>									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	59.4	74.3	74.3	74.2	np	np	np	np	69.1
Aged 18–64 years	76.2	77.0	74.7	78.2	71.5	77.5	np	np	76.0
Aged 65 years or over	69.5	72.0	np	64.1	70.7	np	np	np	69.2
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	57.2	48.4	60.0	np	40.5	np	np	np	51.8
Aged 18–64 years	59.6	51.5	58.6	55.8	57.0	np	np	np	54.8
Aged 65 years or over	47.4	47.9	50.3	44.0	np	np	np	np	47.5
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	37.7	41.5	40.3	38.7	28.9	np	np	np	36.9
Aged 18–64 years	22.6	27.6	27.7	24.0	26.1	24.5	np	27.0	25.8
Aged 65 years or over	19.4	29.1	25.0	21.2	26.6	np	np	np	24.2
<i>2009-10</i>									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	56.3	67.4	np	67.1	np	np	np	np	62.2
Aged 18–64 years	72.4	74.6	76.1	75.9	71.7	77.1	np	np	73.9
Aged 65 years or over	45.9	69.9	67.7	60.4	69.0	np	np	np	61.6
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	61.1	48.7	55.9	64.8	41.7	np	np	np	50.7
Aged 18–64 years	58.7	51.1	61.6	54.5	58.0	np	np	np	54.3
Aged 65 years or over	39.6	47.3	48.5	44.0	np	np	np	np	46.5
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	32.9	38.2	41.5	39.3	28.4	np	np	np	36.4
Aged 18–64 years	21.6	26.9	30.2	26.5	24.5	27.4	16.4	24.7	26.1
Aged 65 years or over	22.6	25.9	25.6	20.4	20.2	np	np	np	23.1

Table 12A.97 People who received mental health care provided by State and Territory public mental health services and who significantly improved, by service type and age group (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
2010-11									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	59.0	62.1	np	67.0	np	np	np	np	59.8
Aged 18–64 years	71.5	74.3	76.0	77.6	73.8	75.8	np	np	74.1
Aged 65 years or over	60.1	74.5	64.5	61.4	80.7	np	np	np	68.1
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	59.1	51.3	57.1	63.8	38.7	np	np	np	51.2
Aged 18–64 years	61.4	45.7	62.8	52.9	62.6	58.7	np	np	51.8
Aged 65 years or over	45.2	42.4	47.1	44.7	np	np	np	np	44.1
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	37.8	38.4	46.0	39.4	25.2	np	np	np	37.0
Aged 18–64 years	20.6	24.9	28.0	23.1	25.2	24.3	15.4	27.3	24.4
Aged 65 years or over	22.0	28.6	24.3	20.0	19.7	np	np	np	23.8
2011-12									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	54.0	na	53.5	np	np	np	np	np	53.2
Aged 18–64 years	70.5	na	76.5	73.9	73.5	73.3	np	78.6	73.1
Aged 65 years or over	59.1	na	70.6	60.6	77.2	np	np	np	64.2
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	57.9	na	58.9	63.1	40.0	np	np	np	51.1
Aged 18–64 years	58.0	na	55.1	50.8	62.5	np	np	np	55.9
Aged 65 years or over	45.2	na	46.5	37.2	np	np	np	np	44.1
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	38.1	na	43.0	36.6	27.4	np	np	np	36.5
Aged 18–64 years	21.0	na	27.1	23.5	21.0	22.5	np	25.9	23.6
Aged 65 years or over	20.6	na	30.4	20.4	23.9	np	np	np	23.5
2012-13									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	57.7	na	52.2	np	37.0	np	np	np	53.2
Aged 18–64 years	72.2	na	75.6	76.7	76.4	77.3	np	77.7	74.6
Aged 65 years or over	61.1	na	74.3	60.7	np	np	np	np	65.3
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	57.5	na	60.2	57.9	38.3	np	np	np	51.4
Aged 18–64 years	53.8	na	54.0	51.0	54.3	np	np	np	53.6
Aged 65 years or over	44.0	na	44.7	39.1	np	np	np	np	44.2
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	34.5	na	42.0	38.7	29.2	np	np	np	36.3
Aged 18–64 years	20.5	na	28.4	23.1	18.8	17.7	np	27.7	23.7
Aged 65 years or over	19.4	na	27.1	19.3	23.8	np	np	np	22.2

Table 12A.97 People who received mental health care provided by State and Territory public mental health services and who significantly improved, by service type and age group (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
<i>2013-14</i>									
Group A: People discharged from hospital who significantly improved (g)									
Aged 0–17 years	60.1	58.4	55.5	np	44.1	np	np	np	56.5
Aged 18–64 years	71.2	75.2	77.6	78.0	71.9	76.0	40.1	78.8	74.6
Aged 65 years or over	59.2	72.3	66.9	60.5	np	np	np	np	66.6
Group B: People discharged from community care who significantly improved (h)									
Aged 0–17 years	55.6	42.6	56.6	63.9	41.5	np	np	np	48.7
Aged 18–64 years	52.2	46.3	55.9	52.1	53.0	np	np	np	50.1
Aged 65 years or over	39.9	43.2	45.6	41.3	np	np	np	np	42.9
Group C: People in ongoing community care who significantly improved (i)									
Aged 0–17 years	34.2	36.2	41.7	41.4	31.5	np	np	np	37.0
Aged 18–64 years	20.8	27.4	24.3	25.4	23.0	23.3	np	26.0	24.2
Aged 65 years or over	19.0	27.5	23.6	18.7	30.3	np	np	np	23.6

- (a) These data were prepared by the Australian Mental Health Outcomes and Classification Network, using data submitted by State and Territory governments to the Australian Government Department of Health. Assessment of clinical outcomes is based on the changes reported in a consumer's score on a rating scale known as the Health of the Nation Outcomes Scale (HoNOS), or in the case of children and adolescent consumers, the Health of the Nation Outcome Scales for Children and Adolescents (HoNOSCA). Developed originally in England in the 1990s, these ratings scales comprise standard items that are rated by a clinician to measure the severity of the consumer's symptoms or disability across a range of domains (for example, depressed mood, hallucinations, substance use, suicidality, overactivity, activities of daily living, cognitive impairment). The HoNOS/HoNOSCA form part of small suite of standardised rating scales used to monitor outcomes across state and territory public sector mental health services and private hospitals with a specialised psychiatric unit. To be considered valid, HoNOS, or the HoNOSCA data needs to be completed correctly (a specified minimum number of items completed) and have a "matching pair" — that is, a beginning and end rating are needed to enable an outcome score to be determined.
- (b) Proportions may not add to 100 per cent due to rounding.
- (c) For all consumer groups, outcome scores for each episode are classified as either 'significant improvement', 'significant deterioration' or 'no significant change', based on Effect Size. Effect size is a statistic used to assess the magnitude of a treatment effect. It is based on the ratio of the difference between pre- and post- scores to the standard deviation of the pre- score. As a rule of thumb, effect sizes of 0.2 are considered small, 0.5 considered medium and 0.8 considered large. Based on this rule, a medium effect size of 0.5 was used to assign outcome scores to the three outcome categories. Thus individual episodes were classified as either: 'significant improvement' if the Effect Size index was greater than or equal to positive 0.5; 'significant deterioration' if the Effect Size index was less than or equal to negative 0.5; or 'no change' if the index was between -0.5 and 0.5.
- (d) Victorian 2011-12 and 2012-13 data are unavailable due to service level collection gaps resulting from protected industrial action during this period. All national averages for 2011-12 and 2012-13 exclude Victoria.

Table 12A.97 People who received mental health care provided by State and Territory public mental health services and who significantly improved, by service type and age group (per cent) (a), (b), (c)

	NSW	Vic (d)	Qld	WA	SA	Tas (e)	ACT (f)	NT (f)	Aust (d)
(e)	Industrial action in Tasmania has limited the available data quality and quantity of data for 2011-12.								
(f)	Some data for the ACT and the NT are np (not published) due to insufficient observations. The number of observations of consumer outcomes for some care types is too low to publish because conclusions based on such low numbers are known to have high levels of unreliability. For the purposes of this indicator, the threshold for the minimum number of observations to be reached was set at 200.								
(g)	Group A covers people who received a discrete episode of inpatient care within a state/territory designated psychiatric inpatient unit during the reference year. The defining characteristic of the group is that the episode of inpatient care commenced, and was completed, within the year. Outcome scores were calculated as the difference between the total score recorded at admission and discharge. The analysis excludes episodes where length of stay was three days or less because it is not meaningful to compare admission and discharge ratings for short duration episodes.								
(h)	Group B covers people who received relatively short term community care from a state/territory mental health service during the reference year. The defining characteristic of the group is that the episode of community care commenced, and was completed, within the year. Outcome scores were calculated as the difference between the total score recorded at admission to, and discharge from, community care. A subgroup of people whose episode of community care completed because they were admitted to hospital is not included in this analysis.								
(i)	Group C covers people receiving relatively long term community care from a state/territory mental health service. It includes people who were receiving care for the whole of the reference year, and those who commenced community care sometime after 1 July who continued under care for the rest of the year. The defining characteristic of the group is that all remained in ongoing care when the year ended (30 June). Outcome scores were calculated as the difference between the total score recorded on the first occasion rated and the last occasion rated in the year.								

na Not available. **np** Not published.

Source: AIHW (unpublished) from data provided by the Australian Mental Health Outcomes and Classification Network

Table 12A.98 Deflators used to calculate real State and Territory mental health expenditure (a)

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust</i>
2005-06	80.7	81.7	76.3	78.4	76.4	80.3	80.3	79.4	79.3
2006-07	84.3	84.3	79.8	81.4	79.2	83.7	83.4	82.3	82.6
2007-08	86.7	86.4	82.6	84.3	81.9	86.1	85.0	85.1	85.0
2008-09	88.8	89.2	85.8	87.6	85.1	88.7	88.3	88.2	87.8
2009-10	91.8	92.1	89.8	91.0	88.8	91.7	91.1	91.4	91.1
2010-11	92.7	93.5	91.8	91.8	89.6	92.7	92.0	92.3	92.3
2011-12	94.8	95.1	94.2	94.3	92.9	95.0	94.5	94.7	94.5
2012-13	97.6	97.4	97.1	96.9	96.5	97.5	97.0	98.2	97.3
2013-14	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

(a) The deflators used are the State and Territory implicit price deflators for general government final consumption expenditure on hospital and nursing home services.

Source: ABS (unpublished) *Australian National Accounts: National Income, Expenditure and Product*, Cat. no. 5204.0.

TABLE 12A.99

Table 12A.99 **Estimated resident populations used in mental health per head calculations (a)**

	<i>NSW</i>	<i>Vic</i>	<i>Qld</i>	<i>WA</i>	<i>SA</i>	<i>Tas</i>	<i>ACT</i>	<i>NT</i>	<i>Aust (b)</i>
2005-06	6 718 023	5 023 203	3 964 175	2 029 936	1 544 852	488 098	333 505	207 385	20 311 543
2006-07	6 786 160	5 103 965	4 055 845	2 076 867	1 561 300	491 515	338 381	211 029	20 627 547
2007-08	6 883 852	5 199 503	4 159 990	2 135 006	1 578 489	495 858	344 176	216 618	21 016 121
2008-09	7 001 782	5 313 285	4 275 551	2 208 928	1 597 880	501 774	351 101	222 526	21 475 625
2009-10	7 101 504	5 419 249	4 367 454	2 263 747	1 618 578	506 461	357 859	227 783	21 865 623
2010-11	7 179 891	5 495 711	4 436 882	2 319 063	1 632 482	510 219	364 833	230 299	22 172 469
2011-12	7 247 669	5 574 455	4 513 009	2 387 232	1 645 040	511 718	370 729	232 365	22 485 340
2012-13	7 348 899	5 679 633	4 610 932	2 472 717	1 662 169	512 422	379 554	236 869	22 906 352
2013-14	7 465 497	5 790 990	4 690 910	2 550 874	1 677 250	513 955	384 147	242 573	23 319 385

(a) The data represent the mid-point of the relevant financial year. For 2011-12 data, the mid-point is 31 December 2011.

(b) Includes other territories.

Source: ABS (various issues), *Australian Demographic Statistics, December (various years)*, Cat. no. 3101.0; table 2A.2.

Data quality information — Mental health management, chapter 12

Data Quality Information

Data quality information (DQI) provides information against the seven Australian Bureau of Statistics (ABS) data quality framework dimensions, for a selection of performance indicators in the Mental health management chapter. DQI for additional indicators will be progressively introduced in future reports.

Technical DQI has been supplied or agreed by relevant data providers. Additional Steering Committee commentary does not necessarily reflect the views of data providers.

DQI Contents

New client index	3
Mental health service use by special needs groups and total population	6
Primary mental health care for children and young people	12
Services reviewed against the National Standards	15
Services provided in an appropriate setting	18
Collection of outcomes information	21
Rate of seclusion — acute inpatient units	25
Consumer and carer involvement in decision making	30
Community follow-up after psychiatric admission/hospitalisation	32
Readmissions to hospital within 28 days of discharge	35
Cost of inpatient care — average recurrent cost per inpatient bed day	38
Cost of inpatient care — average length of inpatient stay	42
Cost of community-based residential care	45
Cost of ambulatory care	48
Rates of illicit and licit drug use	52
Prevalence of mental illness	56
Mortality due to suicide	58
Social and economic inclusion of people with a mental illness — participation in employment of working age population	61

Social and economic inclusion of people with a mental illness — participation in education and employment by young people	63
Mental health outcomes of consumers of specialised public mental health services	65

New client index

DQI for this indicator has been sourced from the Australian Institute of Health and Welfare (AIHW) and State and Territory health authorities with additional Steering Committee comments.

Indicator definition and description

Element	Equity — access
Indicator	New client index — State or Territory specialised public mental health services
Measure	<u>Description:</u>
(computation)	Proportion of total clients under the care of State or Territory specialised public mental health services who were new clients. A new client is a consumer who has not been seen by a specialised public mental health service in the five years preceding the initial contact with a service in the relevant reference period. <u>Numerator:</u> Number of new clients — clients who had not been seen by a public mental health service in the five years preceding the initial contact with a service in the relevant reference period. <u>Denominator:</u> Number of total clients under the care of State or Territory specialised public mental health services in the relevant reference period. <u>Computation:</u> Expressed as a proportion: (Numerator ÷ Denominator)*100.
Data source/s	The AIHW using data provided by State and Territory governments from the community mental health care (CMHC), residential mental health and admitted patients mental health collections.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW calculated the indicator based on data supplied by state and territory health authorities.</p> <p>The State and Territory health authorities provide these data according to specifications agreed under the <i>National Key Performance Indicators for Australian Public Mental Health Services</i>. State and Territory health authorities receive these data from specialised mental health organisations/units in psychiatric and acute hospitals, community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p>
Relevance	<p>Estimates are based on all 'in-scope' clients (new and total) who are in receipt of services from state and territory public psychiatric inpatient units, residential units and community mental health services. New clients are those who have not been seen by a public specialised mental health service in the five years preceding the initial contact with a service in the relevant reference period. A consumer is not considered to be 'new' client if they present with a new condition, but have previously received treatment for other conditions.</p> <p>Data for all years reflect full financial year activity — that is, all in scope clients from public specialised mental health services between the period 1 July and 30 June for each financial year.</p> <p>Only state and territory specialised public mental health services are included. New clients may have been treated in the preceding five years outside the state/territory specialised public mental health system in the primary mental health care or the specialist private mental health sector.</p> <p>States and territories vary in their capacity to accurately track clients across organisations, due to the lack of unique patient identifiers or data matching systems. SA</p>

indicated that the data submitted were not based on unique patient identifier or data matching approaches.

Remoteness and socioeconomic status have been allocated using the client's usual residence, not the location of the service provider. State/territory is reported for the state/territory of the service provider. NSW has used postcode based remoteness and SEIFA correspondence files, rather than SA2 concordance tables for these groups.

For NSW, residential clients are not included because their data is manually collected without a Statewide Unique Patient Identifier (SUPI) assigned, thus making the unique counts of the residential clients together with the inpatient and ambulatory clients not possible.

Data are not available for Victoria for 2011-12 or 2012-13. All Australian totals for 2011-12 and 2012-13 exclude Victoria.

All states except Victoria count triage and referral patients, that is those who are assessed and referred on.

Industrial action in Tasmania in 2011-12 and 2012-13 has limited the quality and quantity of community data.

Timeliness

State and Territory governments provide data to the AIHW for national collation, approximately twelve months after the reference period. The reference period for the latest data is 2013-14.

Accuracy

State and Territory governments are primarily responsible for the quality of the data they provide. The AIHW analyses the data, but cannot independently verify them.

Data are subject to ongoing historical validation. Due to this ongoing validation, 2009-10 and 2012-13 data might differ from previous reports.

States and territories differ in their capacity to accurately track clients across organisations or service types, due to the lack of unique patient identifiers or data matching systems. This has led to over/undercounting of clients in some jurisdictions.

- NSW has implemented a SUPI for mental health care. The identification of prior contacts for mental health clients is dependent upon the SUPI, both in coverage (all clients having a SUPI) and in the resolution of possible duplicates. There are differences in the completeness of coverage between the Local Health Districts/Networks and over time. The average SUPI coverage at a State level is 99.9 per cent for 2012-13. The numbers provided are a distinct count of individuals using the SUPI (majority) and a count of individuals at the facility level for a small percentage of clients without a SUPI in the reporting period (which may include some duplicates of those who attended multiple facilities).
- For NSW, residential clients are not included because their data is manually collected without SUPI assigned, thus making the unique counts of the residential clients together with the inpatient and ambulatory clients not possible. The client base of the NSW mental health residential is very small which will have minimal effect on the final result (for example, there were 237 total residential MH clients with 131 potential new clients in 2012-13).
- For Qld, unique client counts are identified through a statistical linkage process rather than a unique client record. Prior to 2009-10, person count was based on linkage of multiple databases for ambulatory mental health services, and the single inpatient database. From 2009-10 onwards, a single statewide system was introduced for ambulatory mental health services, and the unique identifier is used from this system. However, data are linked through a linkage process based on deterministic and probabilistic methods rather than use of a system level unique identifier.
- For WA, the matching of mental health community contacts to inpatient episodes from 2012-13 is done between two separate data systems and requires the use of record linkage to be able to identify the same person in both systems. There are delays associated in the use of record linkage and these delays can result in not getting a match between a community contact and a separation when there should be one. The number of unique consumers (both total and new) could be over-estimated as a result. Data before 2012-13 are based on data submitted for the National Minimum Data Set (NMDS) and have not been revised.
- For SA, the client counts are not unique: they are an aggregation of three

separate databases with no linkage between them. The impact on the result should be minimal due to populations being relatively stable within the three respective catchments.

- For Tasmania, information for years before 2012-13 were extracted from three different data sources and linked together with a Statistical Linkage Key (SLK) for each individual present in the extracts for the reporting period. While every attempt has been made to reduce any duplication of identified clients, using an SLK will lead to some duplication and can wrongly identify clients as new clients. Tasmania has been progressively implementing a state-wide patient identification system. Data for 2012-13 is considered to be the first collection period with this system fully implemented.

Coherence

Data are reported for each year from 2009-10 to 2013-14. There has been no major change to the methodology used to collect the data across years except as outlined below for WA.

The Australian totals for 2011-12 and 2012-13 exclude Victoria and are not comparable to other years.

Jurisdictions can differ in their approaches to counting clients under care. For example, people who are assessed for a mental health service but do not go on to be treated for a mental illness are included in the data by some jurisdictions but not others. Therefore, comparisons between jurisdictions should be made with caution.

States and territories differ in their capacity to accurately track clients across organisations or service types, this can affect the comparability of the results across jurisdictions (see the accuracy dimension).

For WA, data before 2012-13 are based on data submitted for the NMDS and have not been revised.

Accessibility

Data are also available for this indicator in the:

- National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data
- Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Interpretability

Information for understanding this indicator is available:

- the *Key Performance Indicators for Australian Public Mental Health Services, Second Edition* at [www.health.gov.au/internet/mhsc/publishing.nsf/Content/99A25CC5B3781660CA257A5D000235B3/\\$File/kpitech.pdf](http://www.health.gov.au/internet/mhsc/publishing.nsf/Content/99A25CC5B3781660CA257A5D000235B3/$File/kpitech.pdf)
- in the *Key Performance Indicators for Australian Public Mental Health Services, Third Edition*
- from the *Key Performance Indicators for Australian Public Mental Health Services* (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- States and territories vary in their capacity to accurately track clients across organisations, due to the lack of unique patient identifiers or data matching systems.
- Data are not available for Victoria for 2011-12 or 2012-13. All Australian totals for 2011-12 or 2012-13 exclude Victoria.
- Industrial action in Tasmania in 2011-12 and 2012-13 has limited the available data quality and quantity of community data.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Mental health service use by special needs groups and total population

DQI for this indicator has been sourced from the AIHW for the National Healthcare Agreement with additional Steering Committee comments.

Indicator definition and description

Elements	Equity — Access and Effectiveness — Access
Indicators	Mental health service use by special needs groups Mental health service use by total population
Measure (computation)	<p>The <i>numerator</i> is the number of people receiving mental health services, separately for three service types.</p> <p>The <i>denominator</i> is the Estimated Resident Population (ERP) as at 30 June.</p> <p><i>Calculation</i> is $100 \times (\text{Numerator} \div \text{Denominator})$, presented as a percentage and age-standardised to the Australian population as at 30 June 2001, using 5-year age groups to 84 years with ages over 84 years combined. Aboriginal and Torres Strait Islander population data are not available for all data sources for 5-year age groups beyond 64 years. Where data were not available, Aboriginal and Torres Strait Islander disaggregations were standardised to 64 years with ages over 64 years combined.</p> <p>These are calculated separately for public, private, Medicare Benefits Scheme- and Department of Veterans' Affairs (DVA)-funded services.</p>
Data source/s	<p><u>Numerators:</u></p> <p>For Public data: State/Territory data, including admitted specialised mental health hospital, residential mental health care and CMHC services.</p> <p>For Private data: Private Mental Health Alliance (PMHA) Centralised Data Management Service (CDMS) data.</p> <p>For MBS data: Australian Government Department of Health (Health) MBS Statistics.</p> <p>For DVA data: Australian Government DVA Statistical Services and Nominal Rolls using the Departmental Management Information System (DMIS). These data are known as Treatment Account System (TAS) data.</p> <p><u>Denominator:</u></p> <p>ABS ERP as at 30 June.</p> <p>ABS Aboriginal and Torres Strait Islander Estimates and Projections Series B.</p>

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW prepared the denominator and calculated the indicator based on numerators supplied by other data providers. The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>Numerators for this indicator were prepared by State and Territory health authorities, the PMHA, Health and DVA and quality-assessed by the AIHW.</p> <p>The AIHW drafted the initial data quality statement. The statement was finalised by AIHW following input from State and Territory health authorities, PMHA, Health and DVA. The AIHW does not hold the relevant mandated datasets required to independently verify the data tables for this indicator.</p> <p><u>Public data</u></p> <p>The State and Territory health authorities receive these data from public sector specialised mental health services. States and territories use these data for service</p>
----------------------------------	---

	<p>planning, monitoring and internal and public reporting.</p> <p><u>Private data</u></p> <p>The PMHA's CDMS provided data submitted by private hospitals with psychiatric beds. The data are used by hospitals for activities such as quality improvement.</p> <p><u>Health MBS and DVA TAS data</u></p> <p>The Department of Human Services (DHS) processes claims made under the <i>Medicare Australia Act 1973</i>. These data are then regularly provided to Health. DHS also processes claims for DVA Treatment Card holders made through the MBS under the <i>Veterans' Entitlements Act 1986</i>; <i>Military Rehabilitation and Compensation Act 2004</i> and <i>Medicare Australia Act 1973</i>. All claiming data is regularly provided to DVA as per the Memorandum of Understanding between DHS and DVA.</p>
Relevance	<p>Estimates are based on counts of individuals receiving care within the year, by each service type, where each individual is generally counted once regardless of the number of services received. Persons can receive services of more than one type within the year; a count of persons receiving services regardless of type is not available.</p> <p>Persons receiving mental health treatment who are not captured in these data sources include individuals receiving mental health services (other than as admitted patients in private hospitals) funded through other third party funders (e.g. transport accident insurers, workers compensation insurers) or out of pocket sources.</p> <p>There is likely to be considerable overlap between the various data sources since it is likely that patients accessing public services and private hospital services would also access MBS services.</p> <p>Remoteness and socioeconomic status have been allocated using the client's usual residence, not the location of the service provider. State/territory is reported for the state/territory of the service provider.</p> <p><u>Public data</u></p> <p>Person counts for State and Territory mental health services are counts of persons receiving one or more service contacts provided by public sector specialised mental health services, including admitted hospital, community and residential services.</p> <p><u>Private data</u></p> <p>Private hospital estimates are counts of individuals receiving admitted patient specialist psychiatric care in private hospitals.</p> <p><u>Health MBS and DVA TAS data</u></p> <p>Data are counts of individuals receiving mental health-specific MBS services for which Department of Human Services (DHS) has processed a claim.</p> <p>Analyses by state/territory, remoteness and socioeconomic status are based on postcode of residence of the client as recorded by DHS at the date of last service processed in the reference period. As clients may receive services in locations other than where they live, these data do not necessarily reflect the location in which services were received.</p> <p>DVA clients comprised of people receiving Australian Government (Medicare Benefits Scheme- and DVA-funded) clinical mental health services.</p>
Timeliness	<p>The reference periods for these data are 2007-08, 2008-09, 2009-10, 2010-11, 2011-12, 2012-13 and 2013-14.</p>
Accuracy	<p>Cells have been suppressed to protect confidentiality (where the presentation could identify a patient or a single service provider).</p> <p><u>Public data</u></p> <p>State and Territory jurisdictions differ in their capacity to provide accurate estimates of person receiving services. Additionally, jurisdictions differ in their approaches to counting clients under care. For example, people who are assessed for a mental health service but do not go on to be treated for a mental illness are included in the</p>

data by some jurisdictions but not others. Therefore, comparisons between jurisdictions should be made with caution.

The Indigenous status data should be interpreted with caution due to the varying and, in some instances, unknown quality of Indigenous identification across jurisdictions. Indigenous status was missing or not reported for around 7 per cent of all clients in 2013-14.

Private data

Coverage of private hospitals includes all private hospital with designated psychiatric beds and private psychiatric day hospitals.

The data provided are an estimate of overall activity. Actual counts are multiplied by a factor that accounts for the proportion of data missing from the CDMS collection. That adjustment is performed at the level of State and Territory and also financial year, since non-participation rates varied from state to state and financial year.

Patient counts are unique at the hospital level, therefore, duplication of persons in this data may be possible.

Indigenous status information is not collected for these data.

Health MBS and DVA TAS data

As with any administrative system a small degree of error may be present in the data captured.

Data used for statistical purposes are based on enrolment postcode of the patient. This postcode may not reflect the current postcode of the patient if an address change has not been notified to DHS.

The data provided are based on the date on which the claim was processed by DHS, not when the service was rendered. The use of data based on when the claim was processed, rather than when the service was rendered, produces little difference in the total number of persons included in the numerator for the reference period.

People who received more than one type of service are counted once only in the calculations for this indicator.

Health MBS data presented by Indigenous status have been adjusted for under-identification in the DHS Voluntary Indigenous Identifier (VII) database. Indigenous rates are therefore modelled and should be interpreted with caution. These statistics are not derived from the total Australian Indigenous population, but from those Aboriginal and Torres Strait Islander people who have voluntarily identified as Indigenous to DHS. The statistics have been adjusted to reflect demographic characteristics of the overall Indigenous population, but this adjustment may not address all the differences in the service use patterns of the enrolled population relative to the total Indigenous population. The level of VII enrolment (61 per cent nationally as at August 2012) varies across age-sex-remoteness-State/Territory sub-groups and over time which means that the extent of adjustment required varies across jurisdictions and over time. The methodology for this adjustment was developed and verified by the AIHW and Health for assessment of MBS and PBS service use and expenditure for Indigenous Australians. For an explanation of the methodology, see Expenditure on health for Aboriginal and Torres Strait Islander people 2006-07.

DVA TAS data are not available by Indigenous status.

Coherence

Following the 2011 Census of Population and Housing, the Australian Bureau of Statistics (ABS) has rebased the Australian population back to 1991. This rebasing had a significant impact on the population time series, therefore data were resupplied in previous reporting cycles for previous years using the rebased Estimated Resident Population (ERP) and rebased Indigenous population data.

In 2011, the ABS updated the standard geography used in Australia for most data collections from the Australian Standard Geographical Classification (ASGC) to the Australian Statistical Geography Standard (ASGS). Also updated at this time were remoteness areas and the Socio-Economic Indices for Areas (SEIFA), based on the 2011 ABS Census of Population and Housing. The new remoteness areas will be

referred to as RA 2011, and the previous remoteness areas as RA 2006. The new SEIFA will be referred to as SEIFA 2011, and the previous SEIFA as SEIFA 2006.

Data for 2007-08 through to 2010-11 reported by remoteness are reported for RA 2006. Data for 2011-12 and subsequent years are reported for RA 2011. The AIHW considers the change from RA 2006 to RA 2011 to be a series break when applied to data supplied for this indicator, therefore remoteness data for 2010-11 and previous years are not directly comparable to remoteness data for 2011-12 and subsequent years.

Data for 2007-08 through to 2010-11 reported for SEIFA deciles are reported using SEIFA 2006 at the Statistical Local Area (SLA) level. Data for 2011-12 are reported using SEIFA 2011 at the SLA level. Data for 2012-13 and subsequent years are reported using SEIFA 2011 at the Statistical Area (SA) 2 level. The AIHW considers the change from SEIFA 2006 to SEIFA 2011, and the change from SLA to SA2 to be series breaks when applied to data supplied for this indicator. Therefore SEIFA data for 2010-11 and previous years are not directly comparable with SEIFA data for 2011-12, and SEIFA data for 2011-12 and previous years are not directly comparable with SEIFA data for 2012-13 and subsequent years.

Public data

Public data for all collection periods were re-supplied in 2012-13 by jurisdictions due to an expanded scope that includes all specialised public mental health services. Historical disaggregated data were not re-supplied, therefore, comparisons with 2012-13 and later for any disaggregated data is not valid. As mentioned above, public historical Indigenous data were re-calculated with the revised Indigenous population data, however, as per this coherence issue, historical comparisons with 2012-13 and later data are not valid.

Queensland and Western Australia have provided updated data for 2012-13, thus the 2012-13 data have been updated and resupplied in this reporting cycle.

For public sector community mental health services, Victorian data is unavailable (for 2011-12 and 2012-13) due to service level collection gaps resulting from protected industrial action during this period. Industrial action during the 2011-12 and 2012-13 collection periods in Tasmania has limited the available data quality and quantity of data. Australian totals for 2011-12 and 2012-13 only include available data and should therefore be interpreted with caution. Australian totals for 2011-12 and 2012-13 should not be compared to previous or subsequent years.

Tasmania has been progressively implementing a state-wide patient identification system. Data for 2012-13 is considered to be the first collection period with this system fully implemented. Tasmanian data for 2007-08 and 2008-09 include people who received a Helpline services. From 2009-10 onwards these have been excluded consistent with the indicator definitions. Data for 2007-08 and 2008-09 are also limited to people who accessed Community mental health services only. Therefore, Tasmanian data is not comparable across years.

In past years there has been variation in the underlying concept used to allocate remoteness and socioeconomic status across jurisdictions (i.e. location of service provider, location of client or a combination of both). In addition, the underlying concordances used by jurisdictions to allocate remoteness may vary. Since 2009-10, remoteness and socioeconomic status have been allocated using the SLA of the client at last contact. For 2011-12 data all jurisdictions have used the same concordance and proportionally allocated records to remoteness and Socio-Economic Indexes for Areas (SEIFA) categories with the following exception:

- New South Wales and the Northern Territory used postcode concordance (rather than SLA concordance) to allocate records to remoteness and SEIFA.
- From 2009-10 onwards, disaggregation by SEIFA is based on a person's usual residence, not the location of the service provider, except for the Northern Territory data for which the majority of the data was based on the location of the service. Due to system-related issues impacting data quality, Tasmania was unable to provide data by SEIFA for 2008-09.

Comparisons over time for remoteness and socioeconomic status should therefore be interpreted with caution.

Private data

There has been no change to the methodology used to collect the data in 2013-14. Therefore, the data are comparable to previous reporting periods.

Health MBS and DVA TAS data

The same methodology to attribute demographic information to the data has been used in 2013-14 as in previous reporting periods.

For 2010-11 and previous years, remoteness and socioeconomic status for both Health MBS and DVA TAS data were allocated using a postcode concordance. For 2011-12 and subsequent years, DVA TAS data were allocated to remoteness using geocoding, and to socioeconomic status using an SLA/SA2 concordance.

MBS items 81325 and 81355 were added from 1 November 2008. These items relate to mental health or psychological services provided to a person who identified as being of Aboriginal or Torres Strait Islander descent.

On 1 January 2010, a new MBS item (2702) was introduced for patients of GPs who have not undertaken mental health skills training. Changes have been made to the existing MBS item 2710 to allow patients of GPs who have undertaken mental health skills training to access a higher rebate. Both of these items relate to the preparation of a GP mental health treatment plan.

On 1 November 2011, MBS items 2715 and 2717 were introduced to cover preparation of a GP mental health treatment plan by a GP who has undertaken mental health skills training. At the same time MBS items 2700 and 2701 were introduced to cover preparation of a GP mental health treatment plan by a GP who has not undertaken mental health skills training.

On 1 July 2011, MBS item 288 was introduced as a telehealth mental health related item.

MBS item 2719 existed from 1 November 2011 to 30 April 2012.

From 2011-12 MBS item 20104 is included to align with other national indicators.

Caution should be taken when interpreting Indigenous rates over time. All other data can be meaningfully compared across reference periods.

Other publications

The AIHW publication series Mental health services in Australia contains data that is comparable in coverage (using different MBS item splits) and includes a summary of MBS mental health-related items.

The data used in this indicator is also published in the COAG National Action Plan on Mental Health — final progress report covering implementation to 2010-11. There may be some differences between the data published in these two sources as:

rates may be calculated using different ERPs other than the June ERPs used for this indicator,

MBS numbers are extracted using a different methodology. The COAG National Action Plan on Mental Health — final progress report covering implementation to 2010-11 counts a patient in each state they resided in during the reference period but only once in the total whereas this indicator counts a patient in only one State/Territory.

The indicator specifications and analysis methodology used for this report are equivalent to the Healthcare 2011-12: comparing performance across Australia.

Accessibility

MBS statistics are available at:

- www.health.gov.au/internet/main/publishing.nsf/Content/Medicare+Statistics-1
- www.medicareaustralia.gov.au/statistics/mbs_item.shtml

Disaggregation of MBS data by SEIFA is not publicly available elsewhere.

Interpretability

Information is available for MBS data from:

- www.health.gov.au/internet/mbsonline/publishing.nsf/content/medicare-benefits-schedule-mbs-1

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following issues:

- This is a proxy measure of access to appropriate care.
- State and Territory jurisdictions differ in their approaches to counting clients under care, including different thresholds for registering a client. Additionally, they differ in their capacity to provide accurate estimates of individual persons receiving mental health services. Therefore comparisons between jurisdictions need to be made with caution.
- The Indigenous status data should be interpreted with caution:
 - public sector community mental health services (Public) data: There is varying and, in some instances, unknown quality of Aboriginal and Torres Strait Islander identification among jurisdictional data sources.
 - Medicare Benefits Schedule (MBS) data: have been adjusted for under-identification of Indigenous status in the Medicare Australia Voluntary Indigenous Identifier (VII) database
- Disaggregation of this indicator by Indigenous status for private patients and those recorded in DVA data is a priority.
- Remoteness data for 2010-11 and previous years are not directly comparable to remoteness data for 2011-12 and subsequent years.
- SEIFA data for 2010-11 and previous years are not directly comparable with SEIFA data for 2011-12, and SEIFA data for 2011-12 and previous years are not directly comparable with SEIFA data for 2012-13 and subsequent years.
- For public sector community mental health services, Victorian data is unavailable (for 2011-12 and 2012-13) due to service level collection gaps resulting from protected industrial action during this period. Industrial action during the 2011-12 and 2012-13 collection periods in Tasmania has limited the available data quality and quantity of data. Australian totals of public sector community mental health services for 2011-12 and 2012-13 only include available data and should therefore be interpreted with caution. Australian totals for 2011-12 and 2012-13 should not be compared to previous years.
- Historical disaggregated data up to 2011-12 does not match the scope of data from 2012-13, therefore, comparisons between these years for any disaggregated data is not valid.
- Data have been provided according to the State or Territory of service, but at the sub-state level (remoteness area) have been classified by the client's place of usual residence. For example, a person who usually resides in a very remote area of the NT and is treated by a service in a major city in Victoria would be classified at the sub-state level as a very remote area of Victoria (even though Victoria itself has no very remote areas under the ABS remoteness classification). Further work is required to determine whether geographic location for this indicator should be based on usual residence of the client (used for most indicators) or location of the service.
- Data linkage work is underway to obtain comprehensive and consistent data on people with mental illness across the full scope of service types.

Primary mental health care for children and young people

DQI for this indicator has been sourced from the Australian Government (Department of Health) with additional Steering Committee comments.

Indicator definition and description

Element	Effectiveness — access
Indicator	Primary mental health care for children and young people
Measure	<u>Description:</u>
(computation)	Proportion of young people aged under 25 years who received a primary mental health care services subsidised through the MBS. Data are also reported by four age cohorts: pre-school (0–<5 years), primary school (5–<12 years), secondary school (12–<18 years) and youth/young adult (18–<25 years). <u>Numerator:</u> Number of young people aged under 25 years who received a primary mental health care services subsidised through the MBS and by age cohort (pre-school (0–<5 years), primary school (5–<12 years), secondary school (12–<18 years) and youth/young adult (18–<25 years)). <u>Denominator:</u> ERP aged under 25 years and by age cohort (pre-school (0–<5 years), primary school (5–<12 years), secondary school (12–<18 years) and youth/young adult (18–<25 years)). <u>Computation:</u> Expressed as a proportion: (Numerator/s ÷ Denominator/s)*100. Calculated for all young people (aged under 25 years) and separately by age cohort, gender, Indigenous status, remoteness, SEIFA and service type.
Data source/s	<u>Numerator:</u> Department of Health MBS Statistics data. <u>Denominator:</u> ABS Australian Demographic Statistics.

Data Quality Framework Dimensions

Institutional environment	<p>MBS data are an administrative by-product of the DHS, Medicare fee-for-service payment systems. DHS, Medicare collects MBS data under the <i>Human Services (Medicare) Act 1973</i> and regularly provides the data to Department of Health.</p> <p>The ABS operates within a framework of the <i>Census and Statistics Act 1905</i> and the <i>Australian Bureau of Statistics Act 1975</i>. These ensure the independence and impartiality from political influence of the ABS, and the confidentiality of respondents.</p> <p>For more information on the institutional environment of the ABS, including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, please see ABS Institutional Environment at www.abs.gov.au.</p>
Relevance	<p>Includes primary mental health care covered by the MBS only. Other relevant forms of primary mental health care for young people are not incorporated due to a lack of available data, including community health centres, Aboriginal Community Controlled Health Services, school counsellors and health nurses, university and Technical and Further Education counselling services and a component of the mental health care provided by state/territory specialised public mental health services.</p> <p>MBS data are counts of young people receiving mental health-specific MBS services for which DHS has processed a claim, excluding those for psychiatrists. The relevant MBS items are as follows:</p> <ul style="list-style-type: none">• GP and other services include MBS items 170, 171, 172, 2574, 2575, 2577, 2578, 2700, 2701, 2702, 2704, 2705, 2707, 2708, 2710, 2712, 2713, 2715, 2717,

2719, 2721, 2723, 2725, 2727.

- Clinical psychologist services include MBS items 80000, 80005, 80010, 80015, 80020.
- Other allied health services include MBS items 10956, 10968, 80100, 80105, 80110, 80115, 80120, 80125, 80130, 80135, 80140, 80145, 80150, 80155, 80160, 80165, 80170, 81325, 81355, 82000, 82015.

Analyses by state/territory of MBS data is based on postcode of residence of the client as recorded by DHS at the date of last service processed in the reference period. As clients may receive services in locations other than where they live, these data do not necessarily reflect the location in which services were received. The allocation to the state/territory uses a concordance and splits a person where the postcode covers more than one state/territory, therefore the totals may not equal the sum of the individual cells due to rounding.

MBS data are based on the date the claim was processed. Age of the patient is based on their age at the date of the service.

The population data are for the 30 June of the proceeding financial year. For 2013-14 data, it is June 2013. All ERP data are based on the *2011 Census of Population and Housing*.

Timeliness MBS claims data are available within 14 days of the end of a month. The reference period for the latest data is 2014-15.

Accuracy DVA TAS data are not available.

As with any administrative system a small degree of error may be present in the data captured.

For SEIFA and remoteness reporting, data are excluded for patients whose postcodes do not map to a SEIFA or remoteness area. These patients are reported in a 'not stated' category.

Data used for statistical purposes are based on enrolment postcode of the patient. This postcode may not reflect the current postcode of the patient if an address change has not been notified to DHS.

Analyses by age are based on the client's as recorded by DHS, Medicare at the date the last service that was received in the reference period.

The data provided are based on the date on which the claim was processed by DHS, not when the service was rendered. The use of data based on when the claim was processed, rather than when the service was rendered, produces little difference in the total number of persons included in the numerator for the reference period.

People who received more than one type of service are counted once only in the calculations for this indicator.

Health MBS data presented by Indigenous status have been adjusted for under-identification in the DHS Voluntary Indigenous Identifier (VII) database. Aboriginal and Torres Strait Islander rates are therefore modelled and should be interpreted with caution. These statistics are not derived from the total Australian Aboriginal and Torres Strait Islander population, but from those Aboriginal and Torres Strait Islander people who have voluntarily identified as Aboriginal and Torres Strait Islander to DHS. The statistics have been adjusted to reflect demographic characteristics of the overall Aboriginal and Torres Strait Islander population, but this adjustment may not address all the differences in the service use patterns of the enrolled population relative to the total Aboriginal and Torres Strait Islander population. The level of VII enrolment (61 per cent nationally as at August 2012) varies across age-sex-remoteness-State/Territory sub-groups and over time which means that the extent of adjustment required varies across jurisdictions and over time. The methodology for this adjustment was developed and verified by the AIHW and Health for assessment of MBS and PBS service use and expenditure for Aboriginal and Torres Strait Islander Australians. For an explanation of the methodology, see Expenditure on health for Aboriginal and Torres Strait Islander people 2006-07.

Coherence Estimates are compiled the same way across jurisdictions and over time.

The MBS items included can change over time, for example 2700, 2701, 2715 and

	2719 were included for the later years of data.
Accessibility	<p>MBS statistics are available at:</p> <ul style="list-style-type: none"> • www.health.gov.au/internet/main/publishing.nsf/Content/Medicare+Statistics-1 • www.medicareaustralia.gov.au/statistics/mbs_item.shtml
Interpretability	<p>Information for understanding this indicator is available in the:</p> <ul style="list-style-type: none"> • Fourth national mental health plan: measurement strategy, www.health.gov.au/internet/mhsc/publishing.nsf/Content/pub-plan4-meas • National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data.
<u>Data Gaps/Issues Analysis</u>	
Key data gaps/issues	<p>The Steering Committee notes the following key data gaps/issues:</p> <ul style="list-style-type: none"> • Not all relevant forms of primary mental health care for young people are not incorporated due to a lack of available data. • Annual data are available. The most recent data available are for 2014-15. • The data are consistent and comparable over time.

Services reviewed against the National Standards

DQI for this indicator has been sourced from the AIHW and state and territory health authorities, with additional Steering Committee comments.

Indicator definition and description

Element	Effectiveness — appropriateness
Indicator	Services reviewed against the National Standards for Mental Health Services (<i>NSMHS</i>)
Measure	<u>Description:</u>
(computation)	Proportion of expenditure on specialised public mental health services that had completed a review by an external accreditation agency against the NSMHS. <u>Numerator/s:</u> Expenditure on service units, by assessed level (level 1, level 2, level 3, level 4). <u>Denominator:</u> Total expenditure on service units in scope for the NSMHS. <u>Computation:</u> Expressed as a proportion: (Numerator/s ÷ Denominator)*100. Calculated separately by assessed level.
Data source/s	AIHW from the Mental Health Establishments (MHE) NMDS.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW has provided the data for this indicator.</p> <p>The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.</p>
Relevance	<p>The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes). Specialised psychiatric care in non-specialised public mental health inpatient units is not in scope of the MHE NMDS.</p> <p>The NSMHS were first introduced in 1996 and were adopted by all public specialised mental health services and private psychiatric hospitals. Most non-government community mental health services found it difficult to apply many of the NSMHS to the context within which they operated¹. Revised standards were endorsed in September 2010 and these are designed to be applied across the broad range of mental health services (where mental health is the main focus of care), including non-government organisations and private office based services (such as GPs). Coverage of all publicly</p>

¹ DoHA 2010, *National Standards for Mental Health Services: Implementation guidelines for Non-government Community Services*, Australian Government, Canberra.

funded mental health services to which the revised NSMHS now apply would improve the relevance of these data to measurement of this indicator for future reports.

Services were assessed as level 1, level 2, level 3, or level 4 where these levels are defined as:

- *Services at level 1* — the number of specialised public mental health services that have been reviewed by an external accreditation agency and judged to have met all NSMHS.
- *Services at level 2* — the number of specialised public mental health services that have been reviewed by an external accreditation agency and judged to have met some but not all NSMHS.
- *Services at level 3* — the number of specialised public mental health services that are (i) in the process of being reviewed by an external accreditation agency but the outcomes are not known, or (ii) booked for review by an external accreditation agency.
- *Services at level 4* — the number of specialised public mental health services that do not meet criteria detailed under levels 1 to 3, except those for whom the NSMHS do not apply — code 8 in the MHE NMDS.

Assessments against the NSMHS are based on periodic reviews, usually conducted every three to five years. Services assigned a level 1 for the 2011-12 data may have been assessed at this level in a review that was conducted in 2005-06 and therefore this assessed level may not necessarily reflect the quality of the actual services delivered in the 2011-12 reference period, nor the extent to which the NSMHS are used for ongoing quality improvement.

The data element '*National standards for mental health service review status*' is collected at the statistical unit of service unit (admitted patient, ambulatory and residential). Specialised mental health service units relate to units in public psychiatric hospitals, designated psychiatric units in acute care hospitals, public community-based ambulatory and residential services and publicly funded private hospital and non-government residential service units. Non-government operated community residential service units are excluded from the analysis. Aged care community residential services in receipt of funding under the *Aged Care Act 1997* are subject to residential aged care reporting and service standard requirements and are therefore excluded from the NSMHS analysis. Ambulatory services managed by non-government organisations are not defined as statistical units for the MHE NMDS and therefore data on this element are not available for these service types.

Timeliness

State and territory health authorities provide the MHE NMDS data to the AIHW for national collation, on an annual basis approximately nine months after the reference period. The reference period for the most recent data is 2013-14.

Accuracy

Coverage of the MHE NMDS in-scope services for the '*National standards for mental health service review status*' data element is complete across jurisdictions and years.

States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.

The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year.

Due to ongoing validation, 2005-06 to 2012-13 data could differ from previous reports.

Coherence

Data are reported for each year from 2005-06 to 2013-14.

The data reported from 2005-06 to 2009-10 all relate to specialised mental health services assessed against the old NSMHS. Data from 2010-11 will progressively include larger proportions of services assessed against the revised NSMHS that were endorsed in September 2010.

External accreditation agencies can undertake accreditation of a parent health organisation (for example, a hospital) that can cover a number of specialised mental health service units. Accreditation of the parent organisation does not currently require an individual service unit (for example, a community-based ambulatory service managed by the hospital) to be assessed separately against the NSMHS. Assessment against the NSMHS for a service unit must be specifically requested and involves a separate review process. This leads to variation across states and territories in the method used to assign an assessment level (1, 2, 3 or 4) to service units. In some states and territories, if an organisation with multiple service units is assessed at a particular level all the organisation's units are 'counted' at that assessment level. In other jurisdictions, assessments are conducted at the service unit level and the level assigned may or may not be consistent with the other units within the organisation. The approach can also vary across organisations within a single jurisdiction.

The external accreditation agencies such as Australian Council on Healthcare Standards (ACHS) and Quality Improvement Council (QIC) can use differing review methods. In addition, external review is a process of negotiation between a mental health service organisation and the accrediting agency. There may be differences in the extent to which all or some of the NSMHS are considered to be applicable to individual service units.

Accessibility

The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:

- Mental Health Services in Australia — annual publication
- Australia's Health — a mental health chapter is included in this biennial publication
- National Mental Health Reports
- Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).

Unpublished MHE NMDS data are available from the AIHW on request, but clearance for use of these data for a specific purpose needs to be provided by states and territories and there may be costs incurred in gaining access. Cell sizes with small numbers may be suppressed.

Interpretability

Metadata information for the MHE NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.

Information for understanding this indicator is also available:

- in the Key Performance Indicators for Australian Public Mental Health Services, Third Edition
- from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis**Key data gaps/issues**

The Steering Committee notes the following key data gaps/issues:

- There is variation across and within states and territories in the method used to assign an assessment level (1, 2, 3 or 4) to service units. This may affect the comparability of the results across jurisdictions.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Services provided in an appropriate setting

DQI for this indicator has been sourced from the AIHW and state and territory health authorities, with additional Steering Committee comments.

Indicator definition and description

Element	Effectiveness — appropriateness
Indicator	Services provided in an appropriate setting
Measure	<u>Description:</u>
(computation)	Recurrent expenditure on community-based services as a proportion of total expenditure on mental health services. <u>Numerator:</u> Governments' recurrent expenditure on community-based specialised mental health services. Community-based recurrent expenditure for this indicator includes expenditure on ambulatory care, non-government organisations and adult residential services. Aged residential care expenditure is excluded. <u>Denominator:</u> Total government recurrent expenditure on specialised mental health services, excluding aged residential care expenditure and unapportioned indirect expenditure. <u>Computation:</u> Expressed as a proportion: (Numerator/Denominator)*100.
Data source/s	Numerator and Denominator: AIHW from the MHE NMDS.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW has provided the data for this indicator.</p> <p>The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.</p>
Relevance	<p>The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes). Specialised psychiatric care in non-specialised public mental health inpatient units is not in scope of the MHE NMDS.</p> <p>The data elements on direct and indirect recurrent expenditure and grants to non-government organisations are collected at levels in the hierarchy used to capture jurisdiction-wide information on mental health services (state/territory, region, organisation and service units). Non-government grants are collected at the regional and state and territory levels. Direct recurrent expenditure comprises salaries and wages and non-salary expenditure, and is collected at the individual service unit level. Indirect recurrent expenditure is additional expenditure associated with the provision of mental health services not incurred or reported at the individual service unit level. Some indirect expenditure reported at the organisational and regional level can be directly linked to the provision of services by service units and is apportioned to individual</p>

service units. The estimates do not include residual indirect expenditure incurred at the state and territory level or that unapportioned from the organisational or regional level.

Certain categories of expenditure collected under the MHE NMDS are excluded to derive this indicator and improve the relevance of these data to its measurement.

- Community aged residential care expenditure is excluded from community-based expenditure to improve comparability across states and territories. A significant share of jurisdictions do not have this service type.
- Indirect expenditure at the State and Territory level and indirect expenditure at the organisational or regional level that cannot be apportioned to individual services is also excluded. This indicator is seeking to measure the service mix by showing the proportion of expenditure that is community-based relative to the other categories of service expenditure (admitted patients) and not relative to total expenditure, which includes indirect expenditure at the State or Territory level on areas such as program administration and property leasing costs.

Government expenditure on mental health services that are out of scope of the MHE NMDS, such as Medicare-subsidises for community-based services provided by GPs or the personal helpers and mentors program is not included in the analysis.

Timeliness State and territory health authorities provide the MHE NMDS data to the AIHW for national collation, on an annual basis approximately nine months after the reference period. The reference period for the most recent data is 2013-14.

Accuracy Coverage of the MHE NMDS in-scope mental health services' recurrent expenditure is essentially complete across jurisdictions and years.

States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.

The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year.

Data are also subject to ongoing historical validation. Due to this ongoing validation, 2005-06 to 2012-13 data could differ from previous reports.

Coherence Data are reported for each year from 2005-06 to 2013-14. There has been no major change to the method used to collect the data or to derive the results across years for the majority of jurisdictions, therefore the data are largely comparable across most jurisdictions and years.

For NSW, Confused and Disturbed Elderly (CADE) residential mental health services were reclassified as admitted patient hospital services from 1 July 2007. All data relating to these services have been reclassified from 2007-08 onwards, including expenditure. Comparison of NSW data over time therefore should be approached with caution.

Accessibility The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:

- Mental Health Services in Australia — annual publication
- National Mental Health Reports.

Unpublished MHE NMDS data are available from the AIHW on request, but clearance for use of these data for a specific purpose needs to be provided by states and territories and there may be costs incurred in their provision. Cells may be suppressed for confidentiality reasons or where estimates are based on small numbers, resulting in low reliability.

Interpretability Metadata information for the MHE NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.

Data Gaps/Issues Analysis

Key data gaps/issues The Steering Committee notes the following key data gaps/issues:

- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Collection of outcomes information

DQI for this indicator has been sourced from the Australian Mental Health Outcomes and Classification Network (AMHOCN), Department of Health, AIHW and State and Territory governments with additional Steering Committee comments.

Indicator definition and description

Element	Effectiveness — appropriateness
Indicator	Collection of information on consumers' outcomes. This DQI should be considered in conjunction with the DQI for Mental health outcomes of consumers of specialised public mental health services.
Measure (computation)	<p><u>Description:</u></p> <p>Proportion of specialised public mental health service episodes with completed clinical mental health outcome measures data, by consumer type (people in ongoing community-based care, people discharged from community-based care and people discharged from hospital).</p> <p><u>Numerator:</u></p> <p>Number of specialised public mental health service episodes with completed clinical mental health outcome measures data, by consumer type.</p> <p><u>Denominator:</u></p> <p>Estimated number of specialised public mental health service episodes, by consumer type.</p> <p><u>Computation:</u></p> <p>Expressed as a proportion: $(\text{Numerator/s} \div \text{Denominator}) * 100$. Calculated separately by consumer type.</p>
Data source/s	<p><u>Numerator:</u></p> <p>State and territory health authorities' data reported to the National Outcomes and Casemix Collection (NOCC) and analysed by the AMHOCN.</p> <p><u>Denominator:</u></p> <p>State and territory health authorities' data as reported to CMHC NMDS and the Admitted Patient Mental Health Care (APMHC) NMDS and analysed by the Department of Health.</p>

Data Quality Framework Dimensions

Institutional environment	<p>Health Ministers adopted the routine measurement of consumer outcomes as a priority under the <i>National Mental Health Strategy (1992)</i> and in all subsequent National Mental Health Plans. It is also compatible with State and Territory governments' documented policy emphasis on high quality health services and increased consumer and carer participation.</p> <p>The AMHOCN prepared this indicator using the NOCC data on the Health of the Nation Outcome Scales (HoNOS) family of measures. The Australian Government contracts AMHOCN to support the implementation of the NOCC as part of routine clinical practice by undertaking three functions 1) data bureau — receives and processes information 2) analysis and reporting — analyses and reports on the submitted data and 3) training and service development — supports training in the measures and their use for clinical practice, service management and development purposes.</p> <p>The NOCC was endorsed by all State and Territory governments in 2003, and all jurisdictions have reported data since 2004-05. The NOCC protocol prescribes a set of standard measures to be collected at particular times (collection occasions) in the clinical process. Under the NOCC protocol, collection of outcomes data is mandatory at admission, review and discharge. Data collected outside of NOCC protocols are excluded from the analysis.</p>
Relevance	The scope of the NOCC is all specialised public mental health services managed by, or

in receipt of funds from, state or territory health authorities. Australian Government funded aged residential services are excluded.

The purpose of the NOCC is to measure consumer outcomes. This indicator relates only to the collection of data for the HoNOS family of measures (HoNOS; HoNOS for Older People (HoNOS 65+) and HoNOS for Children and Adolescents (HoNOSCA). Other consumer outcomes measures are also collected, including those completed by consumers. For adults and older persons these include: Kessler 10 (K10+), Behavior and Symptom Identification Scales (BASIS-32); for children and adolescents, the parent and youth versions of the Strengths and Difficulties Questionnaire (SDQ). The uptake of these measures is not captured by this indicator.

For an episode to be counted as one for which consumer outcome measures are collected, a minimum of two data collection occasions with 'valid' measures within the reference period are required. 'Valid' measures are those with a correctly completed specified number of items, for the:

- HoNOS/HoNOS 65+ — a minimum of 10 of the 12 items
- HoNOSCA — a minimum of 11 of the first 13 items.

Brief ambulatory and inpatient care episodes (defined as follows) are excluded.

- inpatient care — episodes 3 days or less.
- ambulatory — episodes where the consumers had a treatment period between 1 and 14 days inclusive.

Ambulatory episodes of mental health care that end because of admission to hospital or residential mental health care are also excluded.

The completion of outcomes data are calculated for three consumer groups. Further, the calculation varies depending on the setting and the duration of the episode of care:

- people discharged from hospital, episodes for people who were admitted and discharged from inpatient care during the reference period (an individual can have two episodes of care so the data represent episode-counts, rather than person-counts) — measures need to be 'valid' for both the admission and discharge occasions rated during the reference period
- people in ongoing community-based care, episodes for people who received community care for the whole of the reference period or who commenced community care sometime after 1 July (beginning of the period) and continued to receive care for the rest of the reference period — measures need to be 'valid' for both the first (either an admission or a review) and last (either an admission or a review) occasions rated during the year
- people discharged from community-based care, episodes for people who were discharged from community care (not including those discharged to hospital) who received an episode of community care that started and ended in the reference period — measures need to be 'valid' for both the admission and discharge occasions rated during the reference period.

Outcomes are measured for consumers discharged from residential mental health care also, but there were too few public mental health service episodes with completed clinical mental health outcome measures data to derive coverage estimates.

The number of 'in-scope' specialised public mental health service episodes, for which outcomes data should be collected (the denominator) is not provided directly to the NOCC, but is an estimate based on the CMHC or APMHC NMDs. For determining the denominators for consumers in ongoing ambulatory care and those discharged from ambulatory care the following distinguishing definitions are used:

- ongoing — the estimated unique count of consumers with CMHC treatment periods of greater than 91 days (that is, from their first service contact date to their last service contact date); LESS the estimated number of consumers whose episodes of care were left censored (that is, commenced in an earlier reporting period by finished within the current reporting periods)
- discharged — the estimated unique count of consumers with CMHC treatment periods of 91 days or less (that is, from their first service contact date to their last service contact date); LESS the estimated number of consumers whose episodes of care resulted in a discharge to an inpatient setting.

Data are not available for Victoria for 2011-12 or 2012-13. All Australian totals for

	2011-12 and 2012-13 exclude Victoria. Industrial action in Tasmania in 2011-12 and 2012-13 has affected the quality and quantity of data.
Timeliness	<p>State and territory health authorities provide the CMHC and APMHC NMDS data to the AIHW for national collation, on an annual basis approximately six months after the reference period.</p> <p>State and territory health authorities provide the NOCC data to AMHOCN for national collation, on an annual basis and all data are to be submitted approximately six months after the reference period.</p> <p>The reference period for the latest data is 2013-14.</p>
Accuracy	<p>States and territories are primarily responsible for the quality of the NOCC data they provide. However, AMHOCN undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage, concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage, primarily concerned with identifying inconsistent, anomalous, and exceptional issues in relation to the NOCC protocol as well as flagging invalid domain values and/or missing data.</p> <p>Change in consumers' clinical outcomes is not presented for states and territories with less than 200 unique consumer outcomes identifiable within the reference period. States and territories excluded from individual publication are used to calculate the change in consumers' clinical outcomes for Australia.</p>
Coherence	<p>Data are available for 2007-08 to 2013-14.</p> <p>The numerator and denominator are sourced from different data sets. Estimates of the total number of episodes requiring outcomes assessment is not provided directly to the NOCC, so it is indirectly estimated from the NMDSs (CMHC and APMHC).</p> <p>The Australian totals for 2011-12 and 2012-13 are not comparable to other years as they exclude data for Victoria.</p>
Accessibility	<p>The AIHW and Department of Health provide a variety of products that draw upon the CMHC and APMHC NMDS. Published products available on the AIHW or Department of Health websites include:</p> <ul style="list-style-type: none"> • Mental Health Services in Australia — annual publication mhsa.aihw.gov.au/home/ • Australia's Health — a mental health chapter is included in this biennial publication aihw.gov.au/publication-detail/?id=10737422172 • National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data • the Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/). <p>Unpublished NMDS data are available from the AIHW on request, but clearance for use of these data for a specific purpose needs to be provided by states and territories and there may be costs incurred in gaining access. Cell sizes with small numbers may be suppressed.</p> <p>NOCC data are available on the AMHOCN website amhocn.org/. The following on-line products are available:</p> <ul style="list-style-type: none"> • web decision support tool • NOCC Standard Reports • NOCC Volume and Percentage Clinical Ratings: Australia • NOCC data are also published in the National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data.
Interpretability	<p>Metadata information for the CMHC and APMHC NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.</p> <p>Metadata information for the NOCC are published on the AMHOCN website amhocn.org/.</p> <p>Information for understanding this indicator is also available:</p> <ul style="list-style-type: none"> • in the Key Performance Indicators for Australian Public Mental Health Services,

Third Edition

- from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- The numerator and denominator are sourced from different data sets. Estimates of the total number of episodes requiring outcomes assessment is not provided directly to the NOCC, so it is indirectly estimated from the NMDSs (CMHC and APMHC).
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Rate of seclusion — acute inpatient units

DQI for this indicator has been sourced from the AIHW and State and Territory health authorities with additional Steering Committee comments.

Indicator definition and description

Element	Effectiveness — quality — safety
Indicator	Rate of seclusion — acute inpatient units
Measure (computation)	<p><u>Definition:</u></p> <p>Number of seclusion events per 1000 patient bed days in specialised public mental health acute inpatient units</p> <p><u>Numerator:</u></p> <p>Number of seclusion events in specialised public mental health acute inpatient units.</p> <p><u>Denominator:</u></p> <p>Number of accrued mental health care days in specialised public mental health acute inpatient units.</p> <p><u>Computation:</u></p> <p>Expressed as a rate. Calculation is: (Numerator ÷ Denominator) x 1000.</p>
Data source/s	<p>AIHW 2015, <i>Mental Health Services in Australia Online</i>, mhsa.aihw.gov.au/home/.</p> <p>AIHW publishes data provided by State and Territory governments from their adhoc seclusion data collections.</p>

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW is a major national agency set up by the Australian Government under the <i>Australian Institute of Health and Welfare Act 1987</i> to provide reliable, regular and relevant information and statistics on Australia's health and welfare. It is an independent corporate Commonwealth entity established in 1987, governed by a management Board, and accountable to the Australian Parliament through the Health portfolio.</p> <p>The AIHW aims to improve the health and wellbeing of Australians through better health and welfare information and statistics. It collects and reports information on a wide range of topics and issues, ranging from health and welfare expenditure, hospitals, disease and injury, and mental health, to ageing, homelessness, disability and child protection.</p> <p>The Institute also plays a role in developing and maintaining national metadata standards. This work contributes to improving the quality and consistency of national health and welfare statistics. The Institute works closely with governments and non-government organisations to achieve greater adherence to these standards in administrative data collections to promote national consistency and comparability of data and reporting.</p> <p>One of the main functions of the AIHW is to work with the states and territories to improve the quality of administrative data and, where possible, to compile national datasets based on data from each jurisdiction, to analyse these datasets and disseminate information and statistics.</p> <p>The <i>Australian Institute of Health and Welfare Act 1987</i>, in conjunction with compliance to the <i>Privacy Act 1988 (Cth)</i> ensures that the data collections managed by the AIHW are kept securely, under the strictest conditions with respect to privacy and confidentiality.</p> <p>For further information see the AIHW website www.aihw.gov.au</p> <p>At present there is no formal, routine nationally agreed collection and reporting framework for seclusion events in specialised mental health public acute hospital services. Data are sourced from state and territory seclusion data collections for</p>
----------------------------------	--

specialised mental health public acute hospital services via Safety and Quality Partnership Standing Committee (SQPSC) a subcommittee, of the Mental Health, Drug and Alcohol Principal Committee (MHDAPC).

The Australian Health Ministers Advisory Council (AHMAC) mental health committees are in the process of formalising the current 'ad hoc' SQPSC seclusion data collection arrangements. The Mental Health Information Strategy Standing Committee (MHISSC) is working with AIHW to develop an aggregate seclusion and restraint Data Set Specification (DSS) to standardise the national collection of both seclusion and restraint data (and provide a more detailed data set) from the 2015-16 collection period.

Relevance

Seclusion is the confinement of the consumer at any time of the day or night alone in a room or area from which free exit is prevented. A seclusion event commences when a clinical decision is made to seclude a mental health consumer and ceases when there is a clinical decision to cease seclusion. If a consumer re-enters seclusion within a short period of time this is considered a new seclusion event. The term 'seclusion event' is utilised to differentiate it from the different definitions of 'seclusion episodes' used across jurisdictions.

Data on seclusion events relates to all specialised mental health public hospital acute services. Wards or units other than specialised mental health services, such as emergency departments, are out of scope for this data collection. Specialised mental health acute forensic hospital services are in scope, regardless of which department manages the service, for example a health department versus a correctional services department.

Timeliness

State and Territory governments provide the data to the AIHW via SQPSC for national collation, approximately three months after the reference period. Data are published within six months of the close of the reference period.

The reference period for the latest data is 2014-15.

Accuracy

Estimated acute bed coverage for 2014-15 seclusion data was over 95 per cent based on acute beds reported to the Mental Health Establishments National Minimum Data Set in 2013-14.

Occasionally, jurisdictions re-supply data for seclusion events or number of occupied bed days. Data re-submissions are highlighted in subsequent data supplies, with updated figures reported in the next annual publication. For 2014-15, historical data were re-supplied for two jurisdictions.

Integrity of the supplied seclusion data is tested by AIHW via a series of 'logical' validation checks. Any missing or unusual data is clarified with the supplying jurisdiction.

Some outliers (that is, a small number of clients who have an above average number of seclusion events) are apparent in the data and were not removed, this has the effect of skewed the rates of seclusion for some jurisdictions.

A new data element, average time in seclusion was captured for the 2013-14 collection period and subsequent collections. As the average time in seclusion is significantly higher for forensic units, these units were excluded from average time in seclusion calculations to provide a more realistic estimation of seclusion duration.

The absence of unit record data limits the ability to undertake analysis to provide context around the incidence of seclusion events. For example, the analysis of consumer attributes which may indicate risk factors or a vulnerability to experiencing seclusion and restraint events (that is, legal status, gender, date of birth, Indigenous status, country of birth).

In addition, identifying the timing of seclusion events within an episode of admitted patient mental health care may be informative in mitigating/pre-empting patterns in the use of restrictive practices. Some jurisdictions have the capacity to record and report this information but the lack of cross-jurisdictional consistency restricts the collection and reporting of unit record data at a national level.

Within the aggregate reporting framework, collection of service unit level data is currently not feasible. Although data are collected at target population, collection at the service unit level would improve consistency and comparability with other mental health collections such as the Mental Health Establishment NMDS.

The use of restrictive practices also includes restraint events. However, no national restraint data are currently reported representing a substantial data gap. The AIHW is currently working with the AHMAC mental health committees and jurisdictional representatives to develop national restraint data standards to facilitate the collection and reporting of national restraint event data.

Coherence

Variations in jurisdictional legislation may result in exceptions to the definition of a seclusion event. Data reported by jurisdictions may not be explicitly comparable, jurisdictional comparisons should therefore be made with caution.

Specific jurisdictional caveats are outlined below:

New South Wales

NSW does not have a centralised database for the collection of seclusion data. Services report seclusion rates regularly to the NSW Ministry of Health. Services are required to maintain local seclusion registers, which may be audited by NSW Official Visitors who function with legislative authority to raise issues in relation to patient safety, care or treatment. Seclusion rates are a Key Performance Indicator (KPI) in regular performance reporting to NSW Local Health Districts. Importantly, NSW seclusion rates include bed days for some forensic services managed by correctional facilities.

Note that in calculating seclusion rates at LHD and State level, all acute bed days are included in the denominator, as per national KPI specifications. This includes facilities where no seclusion occurs, since excluding these facilities would falsely increase the seclusion rate.

No seclusion episodes or bed days were provided for facilities which had not yet opened in the earlier part of the collection period.

The proportion of episodes with a seclusion event may be underestimated in some facilities containing multiple acute units, due to the duplicate counting of hospital stays at facility level. The method used in the seclusion collection for the admitted mental health separations will be reviewed.

Victoria

Victoria have comparably lower bed numbers than other jurisdictions, and as such, it may be useful to view the rate of seclusion events in a broader population context (rates per capita).

Seclusion events per 10 000 population in Victoria was 7.2 in 2011-12, 5.8 in 2012-13, 5.5 in 2013-14, and 4.5 in 2014-15.

Queensland

Queensland do not report any acute forensic services to the collection, however forensic patients can and do access acute care through general units.

Lady Cilento Children's Hospital commenced operation in December 2014 replacing the Mater Public Children's Hospital and Royal Children's Hospital. One hospital has activity that through the Patient Administration System is classified as having psychiatric care for children and adolescents. However, these beds are not classified as specialist mental health beds as reported by the hospital to the Mental Health Establishments (MHE) NMDS. Therefore there will be a mismatch of information between the establishment characteristics listed here and those listed in the MHE NMDS.

There are a number of extreme outliers in regards to duration that have significant impact on duration data. These are a combination of data entry error on legal documentation and actual long seclusions. Due to the timeframe required for submission, 2014-15 data is preliminary and includes imputed episodes of care where source data is not yet available.

Western Australia

It should be noted that WA does not have a centralised data base for the collection of seclusion data. Services provided seclusion data from their own data bases. The Chief Psychiatrist in WA has requested, from 1 July 2014, quarterly reporting of seclusion and restraint rates by all current reporting services.

WA has noted two methodologies for calculating 'Accrued mental health care days' to the Mental Health Establishment NMDS and to the Seclusion and Restraint 'ad hoc' collection denominator. Each has used their own data source and methodology to calculate 'Accrued mental health care days'. WA will investigate these differences further and work towards resolving the discrepancies for future data supplied.

In supplying 2014-15 Seclusion and Restraint data, the same methodology has been used as in 2013-14 for consistency, however as mentioned, there will be differences between the MHE NMDS. The 'N Accrued Care Days' for Seclusion and Restraint is supplied from the state data collection which counts the number of beds occupied every midnight. The following inclusions and exclusions that apply to the number of accrued care days for WA:

- Excludes same-day separations
- Excludes leave days where the bed is counted as not occupied at midnight, or where the bed is occupied by another patient at midnight during the leave period
- Includes all specialised mental health inpatient wards
- Includes hospital in the home wards
- Includes only those care days within each financial year, and includes patients admitted through the entire financial year.

South Australia

Recent data reporting improvements will impact on SA data. Importantly, bed days used to calculate SA's seclusion rates are estimated based on 100 per cent occupied bed numbers, which are fluctuating in relation to new infrastructure projects. During 2010-11, a substantial number of seclusion events in one particular hospital were for a small number of patients with over half of these being patient-requested events. This may have impacted on the overall seclusion rate reported for the state for 2010-11.

SA was unable to supply seclusion data for 2008-09. Information on seclusion duration is only available in 4 hour blocks, therefore averages cannot be calculated and seclusion duration figures for SA are not included in national totals.

Tasmania

The increase in the state-wide Tasmanian seclusion rate for 2012-13 and 2013-14 data is due to a small number of clients having an above average number of seclusion events.

Australian Capital Territory

When interpreting these data, the relative small size of the ACT should be noted, with a total of between 63 and 70 acute inpatient beds reported between 2008-09 and 2013-14.

Work is progressive and ongoing as part of a larger process of providing a place of improved safety and security, both for people experiencing an acute episode of mental ill health leading to an inpatient admission, visitors and for the staff who work in this challenging environment.

Northern Territory

The NT was unable to supply seclusion data for 2008-09.

The NT is unable to segregate Forensic Inpatient Episodes and Events from general events. Therefore all NT totals, wherever stated, are comprised of both General and Forensic Inpatient Episodes and Events. As this may artificially inflate NT data, caution should be used when comparing or interpreting this data.

Due to the low ratio of beds per person in the NT compared with other jurisdictions, the apparent rate of seclusion is inflated when reporting seclusion per patient day compared with reporting on a population basis.

Due to the low number of beds in NT, high rates of seclusion for a few individuals have a disproportional effect on the rate of seclusion reported.

NT seclusion data is therefore not directly comparable with other jurisdictions.

Accessibility

Seclusion data are available at AIHW's Mental Health Services in Australia — annual publication (<https://mhsa.aihw.gov.au/services/admitted-patient/restrictive-practices>).

Additional disaggregation of the seclusion data are in this AIHW publication. Additional disaggregation of the seclusion data are in this AIHW publication.

Interpretability Information is available for interpreting seclusion data from AIHW's Mental Health Services in Australia — annual publication (mhsa.aihw.gov.au/services/admitted-patient/restrictive-practices).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- The absence of unit record data limits the ability to undertake analysis to provide context around the incidence of seclusion events. For example, the analysis of consumer attributes which may indicate risk factors or a vulnerability to experiencing seclusion and restraint events (for example, legal status, gender, date of birth, Indigenous status, and country of birth).
- In addition, identifying the timing of seclusion events within an episode of admitted patient mental health care may be informative in mitigating/pre-empting patterns in the use of restrictive practices. Some jurisdictions have the capacity to record and report this information but the lack of cross-jurisdictional consistency restricts the collection and reporting of unit record data at a national level.
- Within the aggregate reporting framework, collection of service unit level data is currently not feasible. Collection at this level would improve consistency and comparability with other mental health collections such as the Mental Health Establishment NMDS.
- The use of restrictive practices also includes restraint events. However, no national restraint data are currently reported representing a substantial data gap. The AIHW is currently working with the AHMAC mental health committees and jurisdictional representatives to develop national restraint data standards to facilitate the collection and reporting of national restraint event data.

Consumer and carer involvement in decision making

DQI for this indicator has been sourced from the AIHW and state and territory health authorities, with additional Steering Committee comments.

Indicator definition and description

Element	Effectiveness — Quality — Responsiveness
Indicator	Consumer and carer involvement in decision making
Measure	<u>Description:</u>
(computation)	Number of paid full time equivalent (FTE) consumer OR carer staff per 1000 FTE direct care, carer and consumer staff
	<u>Numerator:</u>
	1) Number of paid FTE consumer staff.
	2) Number of paid FTE carer staff.
	<u>Denominator:</u>
	Number of paid FTE direct care, carer and consumer staff.
	<u>Computation:</u>
	Expressed as a proportion per 1000 FTE. Calculation is: (Numerator/Denominator*1000).
Data source/s	Numerator and Denominator: AIHW from the MHE NMDS.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW has provided the data for this indicator.</p> <p>The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.</p>
Relevance	<p>The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes). Specialised psychiatric care in non-specialised public mental health inpatient units is not in scope of the MHE NMDS.</p> <p>Direct care staff comprise consultant psychiatrists and psychiatrists, psychiatry registrars and trainees, other medical officers, registered nurses, enrolled nurses, occupational therapists, social workers, psychologists, other diagnostic and health professionals and other personal care staff. Other categories of staff who work in mental health services are collected under the MHE NMDS, such as administrative and clerical staff, but are not included.</p> <p>Mental health consumer and carer workers are individuals who are employed on a paid basis to represent the interests of consumers and carers, respectively, and advocate for their needs. The person must be employed for the expertise developed from their lived experience of mental illness. The person should also receive a salary or contract fee on a regular basis and it excludes individuals who only received reimbursement of</p>

	<p>expenses or occasional sitting fees for attendance at meetings.</p> <p>The MHE NMDS does not collect information on the staffing of, or consumer and carer participation in, specialised ambulatory mental health services managed by government-funded NGOs.</p>
Timeliness	State and territory health authorities provide the MHE NMDS data to the AIHW for national collation on an annual basis, approximately nine months after the reference period. The reference period for the most recent data is 2013-14.
Accuracy	<p>Coverage of the MHE NMDS in-scope mental health services for direct care staff and consumer and carer workers may not be complete across jurisdictions and years due to the transition from a count of consumer/carer consultants up to 2009-10 to a count of mental health consumer/carer workers from 2010-11.</p> <p>States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.</p> <p>Data are also subject to ongoing historical validation. Due to this ongoing validation, 2005-06 to 2012-13 data could differ from previous reports.</p> <p>The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year.</p> <p>WA have advised that data on FTE consumer or carer workers per 1000 direct care, consumer and carer staff do not accurately represent consumer and carer participation strategies used in WA.</p>
Coherence	Data are reported for each year from 2005-06 to 2013-14. Data up to 2009-10 were restricted to consumer/carer consultants. In 2010-11, the definitions were altered to include a broader range of roles in the contemporary mental health environment, transitioning to mental health consumer and carer workers. These improved definitions should promote greater consistency between jurisdictions. Comparisons between data up to 2009-10 and data from 2010-11 should not be made.
Accessibility	<p>The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:</p> <ul style="list-style-type: none"> • Mental Health Services in Australia — annual publication • Australia's Health — a mental health chapter is included in this biennial publication • National Mental Health Reports.
Interpretability	Metadata information for the MHE NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.

Data Gaps/Issues Analysis

Key data gaps/issues	<p>The Steering Committee notes the following key data gaps/issues:</p> <ul style="list-style-type: none"> • From 2010-11, the definitions of consumer/carer workers were altered to include a broader range of roles in the contemporary mental health environment, transitioning to mental health consumer and carer workers. These improved definitions should promote greater consistency between jurisdictions. Comparisons between data up to 2009-10 and data from 2010-11 should not be made. • Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.
-----------------------------	---

Community follow-up after psychiatric admission/hospitalisation

DQI for this indicator has been sourced from the AIHW for the National Healthcare Agreement with additional Steering Committee comments.

Indicator definition and description

Element	Quality – Continuity
Indicator	Community follow-up after psychiatric admission/hospitalisation — rate of community follow-up within first seven days of discharge from a psychiatric admission.
Measure (computation)	<p>Proportion of separations from the mental health service organisation’s acute psychiatric inpatient unit(s) for which a community ambulatory service contact, in which the consumer participated, was recorded in the seven days following that separation.</p> <p>The <i>numerator</i> is the number of in-scope separations from the mental health service organisation’s acute psychiatric inpatient unit(s) for which a community ambulatory service contact, in which the consumer participated, was recorded in the seven days following that separation.</p> <p>The <i>denominator</i> is the number of in-scope separations for the mental health service organisation’s acute psychiatric inpatient unit(s).</p> <p>Calculation is $100 \times (\text{Numerator} \div \text{Denominator})$</p>
Data source/s	State/territory admitted patient and CMHC data.

Data Quality Framework Dimensions

Institutional environment	<p>The tables for this indicator were prepared by the AIHW based on data supplied by state and territory health authorities. The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>AIHW drafted the initial data quality statement (including providing input about the methodology used to extract the data and any data anomalies) in consultation with State and Territory health authorities.</p> <p>The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities receive these data from public sector community mental health services and public hospitals. States and territories use these data for service planning, monitoring and internal and public reporting. The AIHW does not hold the relevant nationally mandated datasets required to independently verify the data tables for this indicator.</p> <p>Community mental health services and public hospitals may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data for publication in the RoGS 2016 and for the indicators section of the AIHW’s <i>Mental health services in Australia website</i>.</p>
Relevance	<p>Estimates are based on all ‘in scope’ separations from state and territory psychiatric acute inpatient units, where ‘in scope’ is defined as those separations for which it is meaningful to examine community follow-up rates. The following separations were excluded: same day separations; statistical and change of care type separations; separations that end by transfer to another acute psychiatric hospital; separations that end by death, left against medical advice/discharge at own risk; separations where the length of stay is one night and a procedure code for ECT is recorded and separations that end by transfer to community residential mental health services.</p> <p>Data for all years reflect full financial year activity – that is, all in scope separations from public sector acute psychiatric units between the period 1 July and 30 June for each financial year.</p> <p>Community mental health contacts counted for determining whether follow-up occurred are restricted to those in which the consumer participated. These may be face-to-face or ‘indirect’ (e.g., by telephone), but not contacts delivered ‘on behalf of the client’ in which they did not participate, with the exception of the NT which includes all contacts, but</p>

advised that the impact on the indicator is believed to be marginal. Contacts made on the day of discharge are also excluded for all jurisdictions.

Only community mental health contacts made by state and territory public mental health services are included. Where responsibility for clinical follow-up is managed outside the state/territory mental health system (e.g., by private psychiatrists, general practitioners), these contacts are not included.

States and territories vary in their capacity to accurately track post-discharge follow-up between hospital and community mental health care services, due to the lack of unique patient identifiers or data matching systems.

- Tasmania has been progressively implementing a state-wide patient identification system. Data for 2012-13 is considered to be the first collection period with this system fully implemented. The improved patient identification system has increased the percentage post-discharge community care reported by Tasmania in 2013-14. Therefore, Tasmanian data is not comparable across years.
- Western Australia indicated that submitted data was not based on a unique state-wide patient identifier system, but rather data linkage which uses probabilistic matching. Data is therefore subject to change as more information about the patient is collected in statewide data collections.

In 2011, the ABS updated the standard geography used in Australia for most data collections from the Australian Standard Geographical Classification (ASGC) to the Australian Statistical Geography Standard (ASGS). Also updated at this time were remoteness areas and the Socio-Economic Indices for Areas (SEIFA). The new remoteness areas are referred to as RA 2011. The new SEIFA are referred to as SEIFA 2011, and the previous SEIFA as SEIFA 2006.

Data for 2011-12 and subsequent years are reported for RA 2011.

Data for 2011-12 are reported using SEIFA 2011 at the Statistical Local Area level (an ASGC substate geographical unit). Data for 2012-13 and 2013-14 are reported using SEIFA 2011 at the Statistical Area (SA) 2 level (an ASGS substate geographical unit). The AIHW considers the change from SLA to SA2 to be a series break when applied to data supplied for this indicator. Therefore, SEIFA data for 2011-12 are not directly comparable with SEIFA data for 2012-13 and subsequent years.

Remoteness and socioeconomic status have been allocated using the SA2 or postcode concordance of the client at last contact. For 2012-13 and later years' data all jurisdictions have used the same concordance and proportionally allocated records to remoteness and Socio-Economic Indexes for Areas (SEIFA) categories with the following exception:

- New South Wales used postcode concordance (rather than SA2 concordance) to allocate records to remoteness and SEIFA.

Remoteness and socioeconomic status have been allocated using the client's usual residence, not the location of the service provider except for the Northern Territory data for which the majority of the data was based on the location of the service. State/territory is reported for the state/territory of the service provider.

Timeliness

The latest reference period for these data is 2013-14.

Accuracy

State and territory jurisdictions differ in their capacity to accurately track post-discharge follow-up between hospital and community service organisations (see Relevance section above for further information).

Due to ongoing validation, historical data could differ from previous reports. Queensland and WA have provided updated data for 2011-12 and 2012-13, thus these data have been updated and resupplied in this reporting cycle.

Coherence

Specifications for this indicator were revised for the National Healthcare Agreement to align with specifications for the nationally agreed key performance indicators for public mental health services. Specifically, the revised indicator focuses on follow-up care for people discharged from acute psychiatric units only, rather than discharges from all psychiatric units.

This indicator is currently reported in the RoGS and in the Indicators section of the AIHW's Mental health services in Australia website. It is also equivalent to the Key Performance Indicators for Australian Public Mental Health Services: MHS PI 12 —

Rates of post-discharge community care (which this new indicator is based on) and the Fourth National Mental Health Plan: NMHP PI 16 — Rates of post-discharge community care.

Clarification of the scope of the separations data was made to the 2012-13 data specification, however, jurisdictions advised that the impact on the overall data is likely to be minimal. Therefore, data are considered comparable across years in terms of the definitions.

Queensland and WA have provided updated data for 2011-12 and 2012-13, thus these data have been updated and resupplied in this reporting cycle.

For public sector community mental health services, Victorian data are unavailable (for 2011-12 and 2012-13) due to service level collection gaps resulting from protected industrial action during this period.

Industrial action during the 2011-12 and 2012-13 collection periods in Tasmania has limited the available data quality and quantity of community data. Australian totals for 2011-12 and 2012-13 should therefore be interpreted with caution.

For 2012-13, the ACT has refined its calculation methodology and as such, comparisons to previous years' results should be viewed with caution.

Source systems vary in terms of whether location data for the patients usual address is SA2 versus postcode.

All jurisdictions have used the same concordance and proportionally allocated records to remoteness and SEIFA categories.

Accessibility

These data are published in the:

- RoGS available at: www.pc.gov.au/rogs.
- Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).

Interpretability

Definitions for this indicator are published in the indicator specifications in METeOR.

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- States and territories vary in their capacity to accurately track post-discharge follow-up between hospital and community service organisations, due to the lack of unique patient identifiers or data matching systems.
- For public sector community mental health services, Victorian data are unavailable (for 2011-12 and 2012-13) due to service level collection gaps resulting from protected industrial action during this period. Industrial action during the 2011-12 and 2012-13 collection periods in Tasmania has limited the available data quality and quantity of community data. Australian totals for 2011-12 and 2012-13 should therefore be interpreted with caution.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.
- SEIFA data for 2011-12 are not directly comparable with SEIFA for 2012-13 or 2013-14.

Readmissions to hospital within 28 days of discharge

DQI for this indicator has been sourced from state and territory health authorities and AIHW with additional Steering Committee comments.

Indicator definition and description

Element	Quality— continuity
Indicator	Readmissions to hospital within 28 days of discharge
Measure (computation)	<p><u>Description:</u></p> <p>Proportion of 'in-scope' admitted patient overnight separations from public psychiatric acute inpatient services that were followed by readmission to public psychiatric acute inpatient services within 28 days of discharge.</p> <p><u>Numerator:</u></p> <p>Number of 'in-scope' admitted patient overnight separations from public psychiatric acute inpatient services that were followed by readmission to public psychiatric acute inpatient services within 28 days of discharge.</p> <p><u>Denominator:</u></p> <p>Number of 'in-scope' admitted patient overnight separations from public psychiatric acute inpatient services.</p> <p><u>Computation:</u></p> <p>Expressed as a proportion: (Numerator ÷ Denominator)*100.</p>
Data source/s	State and territory governments APMHC data set.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW calculated the indicator based on data supplied by state and territory health authorities. The state and territory health authorities receive these data from public hospitals. States and territories use these data for service planning, monitoring and internal and public reporting.</p> <p>Public hospitals may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p>
Relevance	<p>Estimates are based on all 'in scope' overnight separations from state and territory psychiatric acute inpatient units, where 'in scope' is defined as those separations for which it is meaningful to examine readmission after 28 days of discharge rates. The following separations were excluded: same day separations, including index separation and subsequent readmission; statistical and change of care type separations; separations that end by transfer to another acute or psychiatric hospital; separations that end by death, or instances where the person left against medical advice or discharged at own risk, separations where the length of stay is one night only and a procedure code for ECT is recorded.</p> <p>A readmission for any of the separations identified as 'in-scope' is an admission to any other public acute psychiatric unit within the jurisdiction. For this to occur a system of unique client identifiers needs to be in place that allows individuals to be 'tracked' across units. Such systems have been available in all states/territories for the full period (2005-06 to 2013-14), with the exception of Tasmania (which introduced such a system in 2007-08) and SA (which has not yet introduced such a system).</p> <p>Readmissions across state and territory boundaries or movements between public and private hospitals are not captured.</p> <p>No distinction is made between planned and unplanned readmissions because data collection systems in most Australian public mental health services do not include a reliable or consistent method to distinguish a planned from an unplanned admission to hospital.</p> <p>Remoteness and socioeconomic status have been allocated using the client's usual residence, not the location of the service provider. State/territory is reported for the</p>

	state/territory of the service provider.
Timeliness	<p>State and territory health authorities provide these data to Department of Health for national collation, on an annual basis approximately twelve months after the reference period.</p> <p>The latest year of data available is 2013-14.</p>
Accuracy	<p>Coverage of the 'in-scope' separations and readmissions is essentially complete across jurisdictions and years.</p> <p>States and territories are primarily responsible for the quality of these data. The AIHW analyses the data, but cannot independently verify them.</p> <p>SA was limited in their ability to accurately identify unique consumers due to lack of a unique patient identifier. Consequently the result may appear lower than it actually is, as missing or inaccurate linking data between the separate data sets means that some relevant re-admissions may not be identified.</p> <p>For SA, exclusion criteria based on separation end mode have been applied to the index separation only. Same-day separations and separations where length of stay is one night only with ECT procedure recorded are excluded from both index and readmission episode/separations..</p> <p>Data are also subject to ongoing historical validation. Due to this ongoing validation, 2005-06 to 2012-13 data could differ from previous reports.</p>
Coherence	<p>Data are available from 2005-06 to 2013-14. There has been no major change to the method used to collect the data or to derive the results across years for the majority of jurisdictions, therefore the data are largely comparable across most jurisdictions and years.</p> <p>States and territories differ in their capacity to accurately track clients across organisations or service types, this can affect the comparability of the results across jurisdictions (see the relevance and accuracy dimensions).</p> <p>For WA, changes in historical data are due to a change in methodology which has resulted in additional records being selected and used for calculating readmission; hence rates are slightly higher than previously reported.'</p> <p>An absence of a state-wide unique patient identifier in WA means there is a reliance on data linkage which uses probabilistic matching. Data is preliminary and is subject to change.</p> <p>For data before 2012-13, states and territories differed in the overnight separations that they count as 'in scope'. NSW and Queensland excluded separations where length of stay is one night only and the procedure code for ECT is recorded and the ACT excluded all overnight separations with the procedure code for ECT, whereas the others (Victoria, WA, SA, Tasmania and the NT) included all overnight separations for the procedure code for ECT.</p> <p>From 2012-13, the exclusion of overnight stays of one night with an ECT procedure code became a business rule for the calculation of data for this indicator. The change was considered likely to be minimal, therefore, historical data updates were not considered mandatory. The change is also unlikely to alter the interpretability of long term data trends.</p> <p>Queensland applies in-scope filtering to the subsequent readmission that matches the counterpart in-scope filter for index separations. For example, as outlined above, separations that end by transfer to another acute or psychiatric hospital separations are excluded from the denominator (index separation). By the same token, for the subsequent readmission, Queensland excludes separations that are commenced with code indicating they commenced with transfer from another acute or psychiatric hospital. However, these separations may be eligible for consideration as separate index separation if they did not end with a transfer to another facility.</p> <p>From 2012-13, the ACT refined its calculation methodology and as such, comparisons to previous years' results should be viewed with caution.</p>
Accessibility	<p>These data are also published in the:</p> <ul style="list-style-type: none"> • COAG national action plan on mental health progress reports available at

www.coag.gov.au

- National mental health reports available at www.health.gov.au/internet/main/publishing.nsf/Content/mental-data
- Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).

Interpretability

Further information to understand this indicator are available:

- in the COAG national action plan on mental health — progress report 2010-11
- in National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data
- in the Key Performance Indicators for Australian Public Mental Health Services, Third Edition
- from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- No distinction is made between planned and unplanned readmissions.
- States and territories differ in their capacity to accurately track clients across organisations or service types.
- States and territories differ in the overnight separations that they count as 'in scope'.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Cost of inpatient care — average recurrent cost per inpatient bed day

DQI for this indicator has been sourced from the AIHW and state and territory health authorities, with additional Steering Committee comments.

Indicator definition and description

Element	Efficiency
Indicator	Cost for inpatient care — average recurrent cost per inpatient bed day
Measure	<u>Description:</u>
(computation)	Average recurrent cost per inpatient bed day. <u>Numerator:</u> Expenditure on State and Territory funded specialised mental health admitted patient services, by hospital and program type and by target population and program type. <u>Denominator:</u> Number of inpatient bed days in State and Territory funded specialised mental health admitted patient services, by hospital and program type and by target population and program type. Disaggregations for numerator and denominator are: <u>By inpatient target population:</u> <ul style="list-style-type: none">• general, by acute and non-acute• child and adolescent, by acute and non-acute• older persons' psychiatry, by acute and non-acute• forensic psychiatry, by acute and non-acute <u>By hospital type:</u> <ul style="list-style-type: none">• psychiatric hospitals, by acute units and non-acute units• public acute hospital with a psychiatric unit or ward, by acute and non-acute units Computation: Expressed as \$ per bed day. Calculation is Numerator/Denominator. Real expenditure is reported across years. The general formula for applying the deflator (used in the attachment tables) to convert nominal dollars to real dollars is: $R_t = \frac{D_t}{N_t} \times 100$ Where: R_t is real dollars in year t D_t is nominal dollars in year t N_t is the new index based in year t. N_t is sourced from ABS unpublished, government final consumption expenditure on hospitals and nursing homes price deflator for 2013-14 dollars (2013-14=100).
Data source/s	Numerator and Denominator: AIHW from the MHE NMDS.

Data Quality Framework Dimensions

Institutional environment	The AIHW has provided the data for this indicator. The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website. The data were supplied to the AIHW by state and territory health authorities. The state
----------------------------------	--

	<p>and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.</p>
Relevance	<p>The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes). Specialised psychiatric care in non-specialised public mental health inpatient units is not in scope of the MHE NMDS.</p> <p>Bed days include those for same day admissions, which are counted as one day. Leave days are excluded. Same day admissions are a confounding issue that require the identification of intent of admission (that is, day care or overnight stay). Leave days also present complexities in the mental health area and further work is required to ensure that it does not distort this indicator.</p> <p>Expenditure data are for services provided in specialised mental health service units in public psychiatric hospitals, public acute hospitals and publicly funded private hospital units. Expenditure comprises direct and indirect expenditure incurred at the individual service unit level. Some indirect expenditure reported at the organisational and regional level can be directly linked to the provision of services by service units and is apportioned to individual service units. The residual indirect expenditure incurred at the state and territory level and that unapportioned from the organisational or regional level is not included in the estimates.</p> <p>Cost per inpatient bed day data are not adjusted for differences in the client mix. The client mix in inpatient settings can differ — for example, some jurisdictions treat a higher proportion of less complex patients in inpatient settings as distinct from treating them in the community. More relevant measures would be casemix adjusted, for which cost is adjusted to take into account the type and complexity of cases. Data for these measures are not yet available, as casemix funding has not been applied to specialised mental health services.</p>
Timeliness	<p>State and territory health authorities provide the MHE NMDS data to the AIHW for national collation on an annual basis, approximately nine months after the reference period. The reference period for the most recent data is 2013-14.</p>
Accuracy	<p>Coverage of the MHE NMDS in-scope mental health services for expenditure and bed days is essentially complete across jurisdictions and years.</p> <p>States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.</p> <p>Data are also subject to ongoing historical validation. Due to this ongoing validation, 2005-06 to 2011-12 data could differ from previous reports.</p> <p>The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year.</p>
Coherence	<p>Data are reported for each year from 2005-06 to 2013-14. Data should be reported</p>

consistently across most jurisdiction and across years within most jurisdictions.

Costs per inpatient bed day may not be comparable across jurisdictions. Classification of expenditure into target populations and program type is based on the classification of services as reported to the MHE NMDS rather than the characteristics of their patient populations. For a service to be classified as providing a child and adolescent, older persons' or forensic mental health service for example, it must be recognised by the relevant state or territory funding authority as having a corresponding specialised function and is specifically funded to provide such specialty services. It is likely that the cost per patient day for general mental health services in a jurisdiction that has separate child and adolescent and older persons services (for example, NSW and Victoria), may not be comparable to the average cost in a jurisdiction that has general services only (for example, NT).

For NSW, CADE residential mental health services were reclassified as admitted patient hospital services from 1 July 2007. All data relating to these services have been reclassified from 2007-08 onwards, including expenditure. Comparison of NSW data over time therefore should be approached with caution.

Caution is required when interpreting historical Queensland data, particularly as several services reported as forensic up to 2008-09 were reclassified as general services in 2009-10 to more accurately reflect the function of these services. For 2010-11 and 2011-12, a small number of Youth services have been included in the General category at the request of Queensland. Queensland public acute hospital data includes costs associated with extended treatment services (campus and non-campus based) reported as non-acute admitted patient services in public acute hospitals. Queensland does not provide any acute services in public psychiatric hospitals. Additionally, Queensland provides older persons' mental health inpatient services using a number of different service models, however the majority of older persons' acute care is reported through general adult units, which limits comparability with jurisdictions that report these services differently. Queensland does not report any acute forensic services, however forensic patients can and do access acute care through general units.

From 2010-11, a small number of Youth services have been included in the General category at the request of Queensland.

For WA data, a review of services resulted in the reclassification of beds between the acute and non-acute categories for the 2010-11 collection, to more accurately reflect the function of these services.

For SA, any increases in admitted patient expenditure in 2013-14 partly relate to genuine increases in mental health services. However, a significant proportion of the increases relate to improved identification and allocation of direct care and general overhead expenditure to mental health services.

ACT average costs for older person's mental health services during 2006-07 are based on a new 20 bed unit opened in March 2007, in which only 6-10 beds operated due to issues related to staffing resources. This has artificially inflated the average cost of older persons' mental health services relative to other jurisdictions and other years for the ACT.

Accessibility

The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:

- Mental Health Services in Australia — annual publication
- Australia's Health — a mental health chapter is included in this biennial publication
- National Mental Health Reports
- the Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).

Interpretability

Metadata information for the MHE NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.

Further information to understand this indicator are available:

- in the Key Performance Indicators for Australian Public Mental Health Services, Third Edition

-
- from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- The average recurrent cost per inpatient bed day measures are not adjusted for differences in the client mix and this reduces the relevance of these data to the measurement of efficiency.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Cost of inpatient care — average length of inpatient stay

DQI for this indicator has been sourced from the AIHW and state and territory health authorities, with additional Steering Committee comments.

Indicator definition and description

Element	Efficiency
Indicator	Cost of inpatient care — average length of inpatient stay
Measure	<u>Description:</u>
(computation)	Average length of inpatient stay in acute units, by target population. <u>Numerator:</u> Number of inpatient bed days in State and Territory funded specialised mental health admitted patient acute units, by target population. <u>Denominator:</u> Number of separations from State and Territory funded specialised mental health admitted patient acute units, by target population. <u>Disaggregations for numerator and denominator are:</u> <u>By inpatient target population:</u> <ul style="list-style-type: none">• general acute• child and adolescent acute• older persons' psychiatry acute• total acute (excluding forensic) <u>Computation:</u> Expressed as number of days per stay. Calculation is Numerator/Denominator.
Data source/s	Numerator and Denominator: AIHW from the MHE NMDS.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW has provided the data for this indicator.</p> <p>The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.</p>
Relevance	<p>The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes). Specialised psychiatric care in non-specialised public mental health inpatient units is not in scope of the MHE NMDS.</p> <p>Bed days include those for same day admissions, which are counted as one day. Leave days are excluded. Same day admissions are a confounding issue that require the identification of intent of admission (that is, day care or overnight stay). Leave days also present complexities in the mental health area and further work is required to ensure</p>

that it does not distort this indicator.

Average length of stay data are not adjusted for differences in the client mix. The client mix in inpatient settings can differ — for example, some jurisdictions treat a higher proportion of less complex patients in inpatient settings as distinct from treating them in the community. More relevant measures would be relative stay index, for which the length of stay index takes into account the type and complexity of cases. Data for these measures are not yet available, as casemix analysis has not been applied to specialised mental health services.

Patients days for clients who separated in the reference period that were during the previous period (for example, 2009-10), are excluded. Patient days for clients who remain in hospital (that is, are not included in the separations data) are included.

Average length of stay is not calculated for forensic services as the length of stay is determined by factors outside the control of the specialised mental health service. However, the child and adolescent and older persons' psychiatry target population services may include a forensic component.

Average length of stay is not calculated for non-acute inpatient units due to variability across jurisdictions in the models and mix of care (in particular, variability across jurisdiction in mix of non-acute inpatient and community-based residential care units) that would significantly affect the comparability of the average length of stay data.

Timeliness

State and territory health authorities provide the MHE NMDS data to the AIHW for national collation on an annual basis, approximately nine months after the reference period. The reference period for the most recent data are 2013-14.

Accuracy

Coverage of the MHE NMDS in-scope mental health services bed days and separations is essentially complete across jurisdictions.

States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.

The quality of the separations data used to derive this indicator is variable across jurisdictions. Until recently, these separations data were not subject to the level of in depth scrutiny that has applied to other data elements in the MHE NMDS. Therefore, data are only available from 2010-11. It is expected that the quality of these data will improve over time.

The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year.

Coherence

Average length of stay data may not be comparable across jurisdictions. Classification of inpatient days and separations into target populations and program type is based on the classification of services as reported to the MHE NMDS rather than the characteristics of their patient populations. For a service to be classified as providing a child and adolescent, older persons' or forensic mental health service for example, it must be recognised by the relevant state or territory funding authority as having a corresponding specialised function and is specifically funded to provide such specialty services. It is likely that the average length of stay for a general mental health services in a jurisdiction that has separate child and adolescent and older persons services (for example, NSW and Victoria) may not be comparable to the average length of stay that has general services only (for example, NT).

Queensland provides older persons' mental health inpatient services using a number of different service models, however the majority of older persons' acute care is reported through general adult units, which limits comparability with jurisdictions that report these

services differently.’

A small number of Youth services have been included in the General category at the request of Queensland.

Accessibility

The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:

- Mental Health Services in Australia — annual publication
- Australia’s Health — a mental health chapter is included in this biennial publication
- National Mental Health Reports
- the Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).

Interpretability

Metadata information for the MHE NMDS are published in the AIHW’s online metadata repository — METeOR and in the National health data dictionary.

Further information to understand this indicator are available:

- in the Key Performance Indicators for Australian Public Mental Health Services, Third Edition
- from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- The average length of stay measures are not adjusted for differences in the client mix and this reduces the relevance of these data to the measurement of efficiency.
- The quality of the separations data used to derive this indicator is variable across jurisdictions.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Cost of community-based residential care

DQI for this indicator has been sourced from the AIHW and state and territory health authorities with additional Steering Committee comments.

Indicator definition and description

Element	Efficiency
Indicator	Cost of community-based residential care
Measure	<u>Description:</u>
(computation)	Average recurrent cost per patient day for community-based residential care
	<u>Numerator:</u>
	Expenditure on community-based residential care, by target population and staffing provided
	<u>Denominator:</u>
	Number of patient days in community-based residential care, by target population and staffing provided.
	<u>Disaggregations</u> for the numerator and denominator are:
	General adult units
	<ul style="list-style-type: none">• 24 hour staffed• Non-24 hour staffed
	Older people's care units
	<ul style="list-style-type: none">• 24 hour staffed• Non-24 hour staffed
	<u>Computation:</u>
	Expressed as \$ per bed day. Calculation is Numerator/Denominator.
	Real expenditure is reported across years. The general formula for applying the deflator (used in the attachment tables) to convert nominal dollars to real dollars is:
	$R_t = \frac{D_t}{N_t} \times 100$
	Where:
	R_t is real dollars in year t
	D_t is nominal dollars in year t
	N_t is the new index based in year t. N_t is sourced from ABS unpublished, government final consumption expenditure on hospitals and nursing homes price deflator for 2013-14 dollars (2013-14=100).
Data source/s	Numerator and Denominator: AIHW from the MHE NMDS.

Data Quality Framework Dimensions

Institutional environment	The AIHW has provided the data for this indicator.
	The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.
	The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory

and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.

States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.

Relevance

The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes).

Patient days and expenditure relating to community residential services includes that for publicly funded residential services operated by non-government organisations.

Expenditure data are for services provided in community residential units. Expenditure comprises direct and indirect expenditure incurred at the individual service unit level. Some indirect expenditure reported at the organisational and regional level can be directly linked to the provision of services by service units and is apportioned to individual service units. The residual indirect expenditure incurred at the state and territory level and that unapportioned from the organisational or regional level is not included in the estimates.

Cost per patient day data are not adjusted for differences in the client mix. The client mix in community residential settings can differ — for example, some jurisdictions treat a higher proportion of more complex patients in community residential services. More relevant measures would be casemix adjusted to take into account the type and complexity of cases. Data for these measures are not yet available, as casemix funding has not been applied to specialised mental health services.

Data for child and adolescent community-based residential units are included in the data for general acute units for NSW and the ACT. Other jurisdictions do not have these types of units.

For 2011-12, a small number of Youth services have been included in the General category at the request of WA. From 2012-13, a small number of Youth services have been included in the General category at the request of Victoria, WA and the ACT.

Queensland does not report any in-scope government-operated residential mental health services to the MHE NMDS. However, it funds a number of extended treatment services (campus and non-campus based) with full clinical staffing for 24 hours a day, 7 days a week that are reported as non-acute admitted patient services.

From 2013-14, a small number of residential services reported by SA and the NT as Forensic were included in the General category at the request of those jurisdictions.

Timeliness

State and territory health authorities provide the MHE NMDS data to the AIHW for national collation on an annual basis, approximately nine months after the reference period. The reference period for the most recent data are 2013-14.

Accuracy

Coverage of the MHE NMDS in-scope mental health services community residential expenditure and bed days is complete across jurisdictions and years.

States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.

Data are also subject to ongoing historical validation. Due to this ongoing validation, 2005-06 to 2012-13 data could differ from previous reports.

The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year. Delays caused by this change in completing the NSW 2010-11 MHE NMDS has also meant that the figures provided for the RoGS have not completed full validation and may be different to the finalised data that will be provided for the *National Mental Health Report*.

Coherence

Data are reported for each year from 2005-06 to 2013-14. Data should be reported consistently across years within most jurisdictions.

Classification of expenditure and inpatient days into target populations is based on the classification of services as reported to the MHE NMDS rather than the characteristics of their patient populations. For a service to be classified as providing a general or older persons' mental health service, it must be recognised by the relevant state or territory funding authority as having a corresponding specialised function and is specifically funded to provide such specialty services. For NSW and the ACT, some child and adolescent services are reclassified to general adult to protect agency confidentiality.

For NSW, CADE residential mental health services were reclassified as admitted patient hospital services from 1 July 2007. All data relating to these services have been reclassified from 2007-08 onwards, including patient days. Comparison of NSW data over time therefore should be approached with caution.

Several WA residential services reported as 24-hour staffed services in 2009-10 transitioned to a non-24-hour staffed model of care as of 1 July 2010.

Accessibility

The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:

- Mental Health Services in Australia — annual publication
- Australia's Health — a mental health chapter is included in this biennial publication
- National Mental Health Reports.

Interpretability

Metadata information for the MHE NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- The cost of community-based residential care measures are not adjusted for differences in the client mix and this reduces the relevance of these data to the measurement of efficiency.
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.

Cost of ambulatory care

DQI for this indicator has been sourced from the AIHW, state and territory health authorities and Department of Health with additional Steering Committee comments.

Indicator definition and description

Element	Efficiency
Indicator	Cost of ambulatory care
Measure	<u>Description:</u>
(computation)	Average treatment days per episode of ambulatory care. Average cost per treatment day of ambulatory care <u>Numerator:</u> (1) Number of treatment days in ambulatory care. (2) Adjusted recurrent expenditure on ambulatory care. <u>Denominator:</u> (1) Number of statistical episodes of ambulatory care. (2) Number of treatment days in ambulatory care. <u>Computation:</u> Expressed as treatment days per episode OR cost per episode. Calculation is Numerator (1 OR 2)/Denominator (1 OR 2).
Data source/s	Numerator (1): AIHW from the CMHC NMDS. Numerator (2): AIHW from the MHE NMDS Denominator/s: AIHW from the CMHC NMDS.

Data Quality Framework Dimensions

Institutional environment	<p>The AIHW has provided the data for this indicator.</p> <p>The AIHW is an independent corporate Commonwealth entity within the Health portfolio, which is accountable to the Parliament of Australia through the Minister for Health. For further information see the AIHW website.</p> <p>The data were supplied to the AIHW by state and territory health authorities. The state and territory health authorities received these data from specialised mental health organisations/units in psychiatric and acute hospitals, and community-based ambulatory and residential settings. States and territories use these data for service planning, monitoring and internal and public reporting. Organisations may be required to provide data to states and territories through a variety of administrative arrangements, contractual requirements or legislation.</p> <p>States and territories supplied these data under the auspices of the National Healthcare Agreement and the terms of the National Health Information Agreement.</p>
Relevance	<p>The scope of the MHE NMDS is all specialised mental health services managed by, or in receipt of funds from, state or territory health authorities. The purpose of the MHE NMDS is to collect information on the characteristics of specialised mental health services (for example, program type and target populations) and summary information on their expenditure, staffing and activity (for example, patient days, available beds, separations, contact and episodes). Ambulatory services managed by non-government organisations are not defined as statistical units for the MHE NMDS and therefore excluded.</p> <p>The scope of the CMHC NMDS is government-operated community (also termed ambulatory) mental health services. Data collected includes information relating to each individual service contact provided by an in-scope mental health service. Examples of data elements are demographic characteristics of patients, such as age and sex, clinical information, such as principal diagnosis and mental health legal status, and service</p>

provision information, such as contact duration and session type. Ambulatory services managed by non-government organisations are not considered in-scope for the CMHC NMDS and are therefore excluded.

All activity (treatment days and statistical episodes) and expenditure associated with non-uniquely identified consumers is excluded.

Expenditure data are for services provided in public specialised mental health ambulatory services. Expenditure comprises direct and indirect expenditure incurred at the individual service unit level. Some indirect expenditure reported at the organisational and regional level can be directly linked to the provision of services by service units and is apportioned to individual service units. The residual indirect expenditure incurred at the state and territory level and that unapportioned from the organisational or regional level is not included in the estimates.

Treatment days per episode or expenditure per treatment day are not adjusted for differences in the client mix. The client mix in ambulatory settings can differ — for example, some jurisdictions treat a higher proportion of more complex patients in ambulatory settings as distinct from treating them in hospitals. More relevant measures would be casemix adjusted to take into account the type and complexity of cases. Data for these measures are not yet available, as casemix funding/analysis has not been applied to specialised mental health services.

Treatment day refers to any day on which one or more community contacts (direct or indirect) are recorded for a registered client during an ambulatory care episode. 'One treatment day' episodes are included. These episodes are a confounding issue and a method for accounting for 'one treatment day' ambulatory episodes might provide more relevant measures.

An episode of ambulatory care is a three month period of ambulatory care for an individual registered patient where the patient was under 'active care' (one or more treatment days in the period). Community-based periods relate to the following four fixed three monthly periods: January to March, April to June, July to September, and October to December. The three month period used in this indicator to define a treatment episode is arbitrary. Further development of episode-based funding models may enable more meaningful/relevant measures in future.

Data are not available for Victoria for 2012-13. All Australian totals for 2012-13 exclude Victoria.

Industrial action in Tasmania in 2011-12 and 2012-13 affected the quality and quantity of Tasmania's CMHC data.

Timeliness

State and territory health authorities provide the MHE NMDS data to the AIHW for national collation on an annual basis, approximately nine months after the reference period.

State and territory health authorities provide the CMHC NMDS data to the AIHW for national collation on an annual basis, approximately six months after the reference period.

The reference period for the most recent data are 2013-14.

Accuracy

Coverage of the MHE NMDS in-scope expenditure is essentially complete across years. Coverage of the CMHC NMDS in-scope mental health services contacts is variable among the jurisdictions, with coverage issues for both the services in-scope for collection and the reporting of service contacts between clinicians and clients. Work is ongoing to clarify coverage for jurisdictions.

States and territories are primarily responsible for the quality of the MHE NMDS data they provide. However, the Department of Health and the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is overseen by the Department of Health and managed by the AIHW and is concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is managed by the AIHW and is primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues, including invalid values, missing data and historical inconsistency. Potential validation errors are queried with jurisdictions, and corrections and resubmissions are made in response to these edit queries. The AIHW does not

adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.

States and territories are primarily responsible for the quality of the CMHC NMDS data they provide. However, the AIHW undertake extensive validation. Validation is conducted in two stages: (1) The compliance stage is concerned with ensuring that the data file supplied is structurally compliant and correctly formatted. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage is series of edit checks to ensure that the data supplied are consistent, logical and with valid values. Potential validation errors are queried with jurisdictions, and where the priority for correction is considered high, resubmissions are requested in response to these edit queries. A series of additional edit checks are conducted by the AIHW including coverage checks, historical validation and state/territory comparisons. The AIHW does not adjust data to account for possible data errors or missing or incorrect values. Jurisdictions are responsible for adjusting any data that is identified as problematic and re-submitting improved data files.

Data are also subject to ongoing historical validation. Due to this ongoing validation, 2005-06 to 2012-13 data could differ from previous reports.

The quality of the NSW 2010-11 MHE NMDS has been affected by the reconfiguration of 10 Area Health Services into 18 Local Health Districts mid the 2010-11 financial year.

Coherence

Data are reported for each year from 2005-06 to 2013-14.

'Non-uniquely identifiable consumers' are defined as those with service contacts for which a unique person identifier was not recorded. The proportion of contacts attributed to these consumers varies across jurisdictions (for example, from zero to 15 per cent) and can vary in one jurisdiction across time (for example, from 76 to 99 per cent). As all activity (treatment days and statistical episodes) and expenditure associated with non-uniquely identified consumers are excluded using these proportions, the coherence and comparability of the results across jurisdictions and across time may be affected.

The Australian totals for 2011-12 and 2012-13 are not comparable to other years as they exclude data for Victoria.

Accessibility

The AIHW and Department of Health provide a variety of products that draw upon the MHE NMDS. Published products available on the AIHW or Department of Health websites include:

- Mental Health Services in Australia — annual publication
- Australia's Health — a mental health chapter is included in this biennial publication
- National Mental Health Reports
- the Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).

Unpublished MHE NMDS data are available from the AIHW on request, but clearance for use of these data for a specific purpose needs to be provided by states and territories and there may be costs incur in their provision. Cells may be suppressed for confidentiality reasons or where estimates are based on small numbers, resulting in low reliability.

Interpretability

Metadata information for the MHE NMDS are published in the AIHW's online metadata repository — METeOR and in the National health data dictionary.

Further information to understand this indicator are available:

- in the Key Performance Indicators for Australian Public Mental Health Services, Third Edition
- from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- The cost of ambulatory care measures are not adjusted for differences in the client mix and this reduces the relevance of these data to the measurement of efficiency.

-
- Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data.
 - The exclusion of activity (treatment days and statistical episodes) and expenditure associated with non-uniquely identified consumers means that the coherence and comparability of the results across jurisdictions and across time may be affected.

Rates of illicit and licit drug use

DQI for this indicator has been sourced from the AIHW with additional Steering Committee comments.

Indicator definition and description

Element	Outcome
Indicator	Rates of illicit and licit drug use
Measure (computation)	<p><u>Description:</u></p> <p>Proportion of people aged 14 years or over who use specific licit and illicit drugs in the preceding 12 months — by drug type: alcohol, cannabis, ecstasy, cocaine, meth/amphetamine, hallucinogens, Gamma-hydroxybutyrate (GHB), inhalants, and heroin.</p> <p><u>Numerator:</u></p> <p>Number of people aged 14 years or over who use specific licit and illicit drugs in the preceding 12 months — by drug type.</p> <p><u>Denominator:</u></p> <p>Total population aged 14 years or over.</p> <p><u>Computation:</u></p> <p>$(\text{Numerator} \div \text{Denominator}) * 100$</p> <p>Calculated separately, by drug type.</p>
Data source/s	<p>AIHW 2014, <i>National Drug Strategy Household Survey (NDSHS) detailed report 2013</i>, Drug statistics series no. 28, Cat. no. PHE 183, Canberra.</p> <p>AIHW 2011, <i>2010 NDSHS Report</i>, Drug statistics series no. 25, Cat. no. PHE 145.</p>

Data Quality Framework Dimensions

Institutional environment	<p>The NDSHS data were managed, analysed and published by the AIHW. The AIHW is a major national agency set up by the Australian Government under the <i>Australian Institute of Health and Welfare Act 1987</i> to provide reliable, regular and relevant information and statistics on Australia's health and welfare. The AIHW is an independent corporate Commonwealth entity within the Health portfolio, governed by a management Board, and accountable to the Australian Parliament through the Health and Ageing portfolio.</p> <p>The NDSHS is one of the key data collections that support the <i>National Drug Strategy</i>. The last survey in this program was conducted in 2013, with previous surveys in 1985, 1988, 1991, 1993, 1995, 2001, 2004, 2007 and 2010. The data collected from these surveys have contributed to the development of policies for Australia's response to drug-related issues.</p>
Relevance	<p><u>Scope and coverage</u></p> <p>The NDSHS collects self-reported information on tobacco, alcohol and illicit drug use and attitudes from persons aged 12 years and over.</p> <p>Excluded from sampling were non-private dwellings (hotels, motels, boarding houses, etc.) and institutional settings (hospitals, nursing homes, other clinical settings such as drug and alcohol rehabilitation centres, prisons, military establishments and university halls of residence). Homeless persons were also excluded as well as the territories of Jervis Bay, Christmas Island and Cocos Island.</p> <p>The exclusion of people from non-private dwellings and institutional settings, and the difficulty in reaching marginalised people are likely to have affected estimates.</p> <p>The 2013 NDSHS was designed to provide reliable estimates at the national level. The survey was not specifically designed to obtain reliable national estimates for Aboriginal and Torres Strait Islander people, as there was no target sample size for Aboriginal and Torres Strait Islander Australians. In 2013, the sample size for Aboriginal and Torres</p>

Strait Islander Australians was smaller than anticipated based on population estimates, and so estimates based on this population group should be interpreted with caution.

Reference period

The fieldwork was conducted from 31 July to 1 December 2013. Respondents to the survey were asked questions relating to their beliefs and experiences covering differing time periods, predominantly over the previous 12 months.

Geographic detail

In 2013, data were coded to the census collector's district level. Data are generally published at the national level with a selection of data published at the State/Territory and Remoteness Area levels.

Statistical standards

Data on alcohol consumption was collected in accordance with World Health Organization standards and alcohol risk data were reported in accordance with the current 2009 National Health and Medical Research Council's 'Australian Guidelines to Reduce Health Risks from Drinking Alcohol'.

Timeliness

The NDSHS is conducted approximately every three years over a three-four month period. 2013 data were collected between late-July and early December 2013.

A preliminary data set was received by the AIHW in late-January 2014 and initial data checks were completed in early February 2014.

Key findings from the 2013 NDSHS were released on 17 July 2014.

Accuracy

Perceptions of behaviour

It is known from past studies of alcohol consumption that respondents tend to underestimate actual consumption levels (Stockwell et al. 2004). There are no equivalent data on the tendencies for under- or over-reporting of actual illicit drug use.

However, illicit drug users, by definition, have committed illegal acts. They are, in part, marginalised and difficult to reach. Accordingly, estimates of illicit drug use and related behaviours are likely to be underestimates of actual practice

Sample design

The 2013 sample was stratified by region (15 strata in total – capital city and rest of state for each state and territory, with the exception of the ACT, which operated as one stratum). To produce reliable estimates for the smaller states and territories, sample sizes were boosted in Tasmania, the ACT and the NT.

The over-sampling of lesser populated states and territories produced a sample that was not proportional to the state/territory distribution of the Australian population aged 12 years or older. Weighting was applied to adjust for imbalances arising from execution of the sampling and differential response rates, and to ensure that the results relate to the Australian population.

Sampling error

The measure used to indicate reliability of individual estimates reported in 2013 was the relative standard error (RSE). Only estimates with RSEs of less than 25 per cent are considered sufficiently reliable for most purposes. Results subject to RSEs of between 25 per cent and 50 per cent should be considered with caution and those with RSEs greater than 50 per cent should be considered as unreliable for most practical purposes.

Non-sampling error

In addition to sampling errors, the estimates are subject to non-sampling errors. These can arise from errors in reporting of responses (for example, failure of respondents' memories, incorrect completion of the survey form), the unwillingness of respondents to reveal their true responses and the higher levels of non-response from certain subgroups of the population.

Reported findings are based on self-reported data and not empirically verified by blood tests or other screening measures.

Response rates and contact rates

Overall for the 2013 Survey, contact was made with 48 579 in-scope households, of which 23 855 questionnaires were categorised as being complete and useable, representing a response rate for the 2010 survey of 49.1 per cent, slightly lower than the drop and collect component of the 2010 survey (50.6 per cent).

Some survey respondents did not answer all questions, either because they were unable or unwilling to provide a response. The survey responses for these people were retained in the sample, and the missing values were recorded as not answered. No attempt was made to deduce or impute these missing values.

A low response rate does not necessarily mean that the results are biased. As long as the non-respondents are not systematically different in terms of how they would have answered the questions, there is no bias. Given the nature of the topics in this survey, some non-response bias is expected. If non-response bias in the NDSHS is to be eliminated as far as possible, there would need to be additional work conducted to investigate the demographic profile of the non-respondents and the answers they may have given had they chosen to respond.

Aboriginal and Torres Strait Islander Data

The survey was not specifically designed to obtain reliable national estimates for Aboriginal and Torres Strait Islander people, as there was no target sample size for Indigenous Australians. In the 2013 NDSHS, 1.9 per cent of the sample (or approximately 461 respondents) identified as being of Aboriginal or Torres Strait Islander origin. The sample size for Indigenous Australians was smaller than anticipated based on population estimates, and so estimates based on this population group should be interpreted with caution.

The total population of Aboriginal and Torres Strait Islander people forms a very small part of the total Australian population. At the August 2011 census, the Aboriginal and Torres Strait Islander population was officially calculated at 670,000 people, or 2.1 per cent of the total Australian population. At that time, about one-third (35 per cent) of the Aboriginal and Torres Strait Islander population lived in Major cities, 22 per cent in Inner regional areas, 22 per cent in Outer regional areas, 8% in Remote areas and 14 per cent in Very remote areas.

The Aboriginal and Torres Strait Islander population living in Very remote areas shows other differences to populations living in Major cities including in household structure, size and age distribution. The NDSHS sample design is stratified by region and not by remoteness. Due to this sampling design, the NDSHS sample of Indigenous Australians living in Very remote areas comprised of 9 per cent of the population in those regions compared with 14 per cent of Indigenous Australians living in Very remote areas based on the 2011 Census. Therefore, Aboriginal and Torres Strait Islander people in Very remote areas are underrepresented, and it becomes difficult to generalise results from Major cities and regional areas to the whole Indigenous population.

The sampling method employed for the NDSHS invited one participant aged over 12 years to take part in the survey. The sample strategy did not take into account the size of the household selected. This is an issue for respondent selection for Indigenous Australians, as often they live in larger households compared with non-Indigenous Australians. This selection process means that Aboriginal and Torres Strait Islander people are proportionately less likely to be selected.

The NDSHS uses a self-completion questionnaire, and requires good comprehension of the English language (as it is not translated into other languages) and the ability to follow instructions. Practicality of the survey design meant that some Aboriginal communities and those with low levels of English literacy may have been excluded. Response rates are reported in the relevant NDSHS reports.

Coherence

Surveys in this series commenced in 1985. Over time, modifications have been made to the survey's methodology and questionnaire design. The 2013 survey differs from previous versions of the survey in some of the questions asked and also used three follow-up attempts by interviewers instead of the two used in 2010.

Methodology

The 2013 survey was the second to exclusively use the drop and collect method, the first being 2010. In 2007 and 2004, a combination of computer-assisted telephone interviews (CATI) and drop and collect methods were used, and in earlier waves,

personal interviews were also conducted.

The change in methodology in 2010 does have some impact on time series data, and users should exercise some degree of caution when comparing data over time.

Fieldwork was conducted between July and December 2013, slightly later than in previous wave. The collection period also coincided with the 2013 federal election, although no questionnaires were placed on that day.

Sample

To produce reliable estimates for the smaller states and territories, sample sizes were boosted in Tasmania, the ACT and the NT.

In 2013 and 2010, to improve the geographic coverage of the survey, interviewers were flown to Very remote areas selected in the sample. In previous surveys, some Very remote areas that were initially selected in the sample would have been deemed inaccessible and not included in the final sample.

Questionnaire

The 2013 questionnaire was modelled on the 2010 version, to maintain maximum comparability. However, some refinements were made to ensure the questions remained relevant and useful. For more information on questionnaire changes in 2013 see chapter 1 of the 2013 NDSHS report.

Accessibility

Results from the 2013 NDSHS are available on the AIHW website. Key findings can be found in the web compendium: Highlights from the 2013 survey and full published results can be found in the 2013 NDSHS report.

Users can request data not available online or in form the AIHW. Requests that take longer than half an hour to compile are charged for on a cost-recovery basis.

A confidentialised unit record file is available for third party analysis through the Australian Data Archive. Access to the master unit record file may be requested through the AIHW Ethics Committee.

Data for this indicator are also reported in the National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data

Interpretability

Information to aid in interpretation of 2013 NDSHS results may be found in chapter 1 of the 2013 NDSHS report titled 'Introduction'. In addition, the 2013 Technical Report, code book and other supporting documentation are available through the Australian Data Archive website or may be requested from AIHW

Further information to understand this indicator are available in:

- the COAG national action plan on mental health — progress report 2010-11
- National mental health reports www.health.gov.au/internet/main/publishing.nsf/Content/mental-data.

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following key data gaps/issues:

- Respondents tend to underestimate actual alcohol consumption levels.
- Estimates of illicit drug use are also likely to be underestimates of actual practice.
- Reported findings are based on self-reported data and are not independently verified.
- The response rate for the 2013 survey was 49.1 per cent. Some non-response bias is expected — this bias has not been measured

Prevalence of mental illness

DQI for this indicator has been sourced from the ABS with additional Steering Committee comments.

Indicator definition and description

Element	Outcome
Indicator	Prevalence of mental illness
Measure (computation)	<p><i>Numerator:</i></p> <p>Number of people aged 16–85 years who had a mental health disorder diagnosed by the World Mental Health Composite Interviewing Diagnostic Instrument (CIDI), with symptoms in last 12 months.</p> <p><i>Denominator:</i></p> <p>Total population aged 16–85 years.</p> <p><i>Computation:</i></p> <p>$(\text{Numerator} \div \text{Denominator}) * 100$</p> <p>Disaggregated by disorder type and age or sex (national only), State and Territory, by disorder type.</p>
Data source/s	ABS unpublished, <i>2007 National Survey of Mental Health and Wellbeing</i> (Cat. no. 4326.0).

Data Quality Framework Dimensions

Institutional environment	For information on the institutional environment of the ABS, including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, see ABS Institutional Environment (available www.abs.gov.au).
Relevance	<p>The 2007 National Survey of Mental Health and Wellbeing (SMHWP) provides information about the prevalence of selected <i>high prevalence</i> mental disorders in the Australian population aged 16–85 years, the level of impairment associated with these disorders, physical conditions, and the use of health services, such as consultations with health practitioners or visits to hospital. The survey also provides information on the strength of social networks, caring responsibilities and a range of socio-economic and demographic characteristics.</p> <p>The SMHWP was designed to provide prevalence estimates for the mental disorders that are considered to have the highest incidence rates in the population — anxiety disorders (such as social phobia), affective disorders (such as depression) and substance use disorders (such as harmful alcohol use). The SMHWP was not designed to measure the prevalence of all mental health conditions, therefore some severe mental disorders, such as schizophrenia, were not collected.</p> <p>The SMHWP is based on an international survey instrument, the CIDI, developed by the World Health Organization (WHO) for use by participants in the World Mental Health Survey Initiative.</p> <p>The 2007 survey was designed to provide data that were internationally comparable, rather than to provide comparisons with the 1997 survey. The survey was also designed to provide estimates of the prevalence of mental disorders at a national rather than a state/territory level.</p>
Timeliness	<p>The SMHWP was conducted in 1997 and 2007.</p> <p>Results from the 2007 survey were released ten months after the completion of enumeration, in the publication <i>National Survey of Mental Health and Wellbeing: Summary of Results</i> (cat. no. 4326.0).</p>
Accuracy	Estimates from the 2007 SMHWP are subject to sampling and non-sampling errors. The RSE is a measure of the size of the sampling error affecting an estimate; that

is, the error introduced by basing estimates on a sample of the population rather than the full population. Estimates should be considered with reference to their RSEs. Estimates with an RSE between 25 per cent and 50 per cent should be used with caution, and those with an RSE greater than 50 per cent are considered too unreliable for general use. Non-sampling errors are inaccuracies that occur because of imperfections in reporting by respondents and interviewers, as well as errors made in coding and processing the data.

The SMHWP was designed primarily to provide estimates at the national level. Due to the higher than expected non-response rate, RSEs were somewhat larger than originally designed. While broad estimates are available for the larger states, users should exercise caution when using estimates at this level due to relatively high sampling errors.

Coherence The 2007 SMHWP was the second survey of this type conducted by the ABS, with the previous survey conducted in 1997. Care should be exercised when comparing data between surveys as there have been a number of changes to the scope, design, collection, methodology and content.

Supporting documentation released with the survey data can assist in understanding the relationships between data variables within the dataset and in comparisons with data from other sources.

Accessibility The main products available from this survey are:

- National Survey of Mental Health and Wellbeing: Summary of Results, 2007 (Cat. no. 4326.0)
- National Survey of Mental Health and Wellbeing: Users' Guide, 2007 (Cat. no. 4327.0)
- Microdata: National Survey of Mental Health and Wellbeing, Basic and Expanded Confidentialised Unit Record Files, 2007 (Cat. no. 4326.0.30.001)
- Technical Manual: National Survey of Mental Health and Wellbeing, Confidentialised Unit Record Files (Cat. no. 4329.0).

Further information may be available on request. The ABS observes strict confidentiality protocols as required by the *Census and Statistics Act (1905)*. This may restrict access to data at a very detailed level.

Interpretability The *National Survey of Mental Health and Wellbeing: Summary of Results* (Cat. no. 4326.0) includes explanatory material to aid the interpretation of the survey results. More detailed information is available in the *National Survey of Mental Health and Wellbeing: Users' Guide* (Cat. no. 4327.0).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following issues:

- The SMHWP was designed to provide estimates at the national level. Broad estimates are available for the larger states, but users should exercise caution when using estimates at this level due to relatively high sampling errors.
- The SMHWP was designed to provide prevalence estimates for the mental disorders that are considered to have the highest incidence rates in the population — anxiety disorders (such as social phobia), affective disorders (such as depression) and substance use disorders (such as harmful alcohol use). It does not measure the prevalence of some severe mental disorders, such as schizophrenia (which are the mental illnesses most frequently treated by specialised public mental health services).

Mortality due to suicide

DQI for this indicator has been sourced from the ABS with additional Steering Committee comments.

Indicator definition and description

Element	Outcome
Indicator	Mortality due to suicide
Measure (computation)	<p><i>Numerator:</i></p> <p>Number of people who have died by suicide over the relevant reference period:</p> <ul style="list-style-type: none">• five year period (2008–2012)• single reference year (2012) <p><i>Denominator:</i></p> <p>ERP.</p> <p><i>Computation:</i></p> <p>$(\text{Numerator} \div \text{Denominator}) * 100\,000$</p> <p>Expressed as crude, age-specific or age standardised rates.</p> <p>Disaggregated by age and sex (national only), State and territory for all persons, young people (15–24 years), by geographical region and Indigenous status.</p>
Data source/s	<p><i>Numerator:</i> ABS <i>Causes of Death</i> collection (Cat. no. 3303.0)</p> <p><i>Denominator:</i> ABS ERP (Cat. no. 3101.0); Experimental Estimates and Projections, Aboriginal and Torres Strait Islander Australians, 1991 to 2021 (Cat. no. 3238.0); ASGC (Cat. no. 1216.0).</p>

Data Quality Framework Dimensions

Institutional environment	<p>Statistics presented in <i>Causes of Death, Australia, 2012</i> (Cat. no. 3303.0) are sourced from deaths registrations administered by the various state and territory Registrars of Births, Deaths and Marriages. It is a legal requirement of each state and territory that all deaths are registered. Information about the deceased is supplied by a relative or other person acquainted with the deceased, or by an official of the institution where the death occurred on a <i>Death Registration Form</i>. As part of the registration process, information on the cause of death is either supplied by the medical practitioner certifying the death on a <i>Medical Certificate of Cause of Death</i>, or supplied as a result of a coronial investigation.</p> <p>Death records are provided electronically to the ABS by individual Registrars on a monthly basis. Each death record contains both demographic data and medical information from the <i>Medical Certificate of Cause of Death</i> where available. Information from coronial investigations are provided to the ABS through the National Coroners Information System (NCIS).</p> <p>For information on the institutional environment of the ABS, including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, see ABS Institutional Environment (available www.abs.gov.au).</p>
Relevance	<p>The ABS Causes of Death collection includes all deaths that occurred and were registered in Australia, including deaths of persons whose usual residence is overseas. Deaths of Australian residents that occurred outside Australia may be registered by individual Registrars, but are not included in ABS deaths or causes of death statistics.</p> <p>From the 2007 reference year, the scope of the collection is:</p> <ul style="list-style-type: none">• all deaths registered in Australia for the reference year and are received by the ABS by the end of the March quarter of the subsequent year; and• deaths registered prior to the reference year but not previously received from

the Registrar nor included in any statistics reported for an earlier period.

For example, records received by the ABS during the March quarter of 2011 which were initially registered in 2010 or prior (but not forwarded to the ABS until 2011) are assigned to the 2010 reference year. Any registrations relating to 2010 which are received by the ABS after the end of the March 2011 quarter are assigned to the 2011 reference year.

Data in the Causes of Death collection include demographic items, as well as causes of death information, which is coded according to the International Classification of Diseases (ICD). ICD is the international standard classification for epidemiological purposes and is designed to promote international comparability in the collection, processing, classification, and presentation of causes of death statistics. The classification is used to classify diseases and causes of disease or injury as recorded on many types of medical records as well as death records. The ICD has been revised periodically to incorporate changes in the medical field. The 10th revision of ICD (ICD-10) has been used since 1997.

Non-Indigenous data from the Causes of Death collection do not include death registrations with a 'not stated' Indigenous status.

Timeliness

Causes of death data are published on an annual basis.

There is a focus on fitness for purpose when causes of death statistics are released. To meet user requirements for accurate causes of death data it is necessary to obtain information from other administrative sources before all information for the reference period is available (for example, information from finalisation of coronial proceedings to code an accurate cause of death). A balance therefore needs to be maintained between accuracy (completeness) of data and timeliness. ABS provides the data in a timely manner, ensuring that all coding possible can be undertaken with accuracy prior to publication.

In addition, to address the issues which arise through the publication of causes of death data for open coroners cases, these data are subject to a revisions process. This process enables the use of additional information relating to coroner certified deaths either 12 or 24 months after initial processing.

Accuracy

Information on causes of death is obtained from a complete enumeration of deaths registered during a specified period and are not subject to sampling error. However, deaths data sources are subject to non-sampling error which can arise from inaccuracies in collecting, recording and processing the data. Every effort is made to minimise non-sample error by working closely with data providers, running quality checks throughout the data processing cycle, training of processing staff, and efficient data processing procedures.

Causes of death data for 2006, 2007, 2008, 2009, 2010, 2011 and 2012 have been subject to revision. All coroner certified deaths registered after 1 January 2006 are subject to a revision process. This is a change from previous years where all ABS processing of causes of death data for a particular reference period was finalised approximately 13 months after the end of the reference period. Where insufficient information was available to code a cause of death (for example, a coroner certified death was yet to be finalised by the Coroner), less specific ICD codes were assigned as required by the ICD coding rules. The revision process enables the use of additional information relating to coroner certified deaths as it becomes available over time. This results in increased specificity of the assigned ICD-10 codes.

For this year's report, causes of death data for 2011, 2012 were updated as more information became available. Final data for 2006, 2007, 2008, 2009, 2010 and 2011, revised data for 2012 and preliminary data for 2013 have been published in the *2013 Causes of Death* publication, released in March 2015. 2012 and 2013 causes of death will be revised in the 2014 Causes of Death publication due for release in 2016. Revisions will only affect coroner certified deaths, as further information becomes available to the ABS about the causes of these deaths. See *Causes of Death, Australia* (Cat. no. 3303.0).

Some rates are unreliable due to small numbers of deaths over the reference period. Resultant rates could be misleading for example where the non-Indigenous mortality rate is higher than the Aboriginal and Torres Strait Islander mortality rate. All rates for this indicator must be used with caution.

Non-Indigenous population estimates are available for census years only. In the intervening years, Aboriginal and Torres Strait Islander population figures are derived from assumptions about past and future levels of fertility, mortality and migration. In the absence of non-Indigenous population figures for these years, it is possible to derive denominators for calculating non-Indigenous rates by subtracting the Aboriginal and Torres Strait Islander population from the total population. Such figures have a degree of uncertainty and should be used with caution, particularly as the time from the base year of the projection series increases.

Coherence

The methods used to construct the indicator are consistent and comparable with other collections and with international practice.

The completeness or quality of older (unrevised) versus newer data (subject to a revisions process) can affect comparisons across time. The accuracy dimension contains information pertinent to coroner certified deaths affected by the revision process.

The ERPs used to derived rates differ across years and tables. Some are derived using ERPs based on the 2001 Census, 2006 Census or 2011 Census. See particular tables for details. Rates derived using ERPs based on different Censuses are not comparable.

Accessibility

Causes of death data are available in a variety of formats on the ABS website under the 3303.0 product family. ERP data are available in a variety of formats on the ABS website under the 3101.0 and 3201.0 product families. Further information on deaths and mortality may be available on request. The ABS observes strict confidentiality protocols as required by the *Census and Statistics Act (1905)*. This may restrict access to data at a very detailed level.

Interpretability

Information on how to interpret and use cause of death data are available from Explanatory Notes in *Causes of Death, Australia* (Cat. no. 3303.0).

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following issue:

- Causes of death data are subject to a revisions process. Final data for 2006, 2007, 2008, 2009, 2010 and 2011, revised data for 2012 and preliminary data for 2013 have been published in the 2013 Causes of Death publication. Data for 2012 and 2013 causes of death will be revised in 2016.

Social and economic inclusion of people with a mental illness — participation in employment of working age population

DQI for this indicator has been sourced from the ABS with additional Steering Committee comments.

Indicator definition and description

Element	Outcome
Indicator	Social and economic inclusion of people with a mental illness — participation in employment of working age population.
Measure (computation)	<i>Numerator:</i> Number of people aged 16-64 years who are employed (by mental health status) <i>Denominator:</i> Number of people aged 16-64 years in the population (by mental health status) <i>Computation:</i> (Numerator ÷ Denominator)*100 Note: People with a mental health condition are defined as having a self-reported mental or behavioural problem that has lasted for six months, or which the respondent expects to last for six months or more.
Data source/s	ABS unpublished, Australian Health Survey (AHS) 2011–13 (2011-12 National Health Survey component).

Data Quality Framework Dimensions

Institutional environment	<p>The AHS was collected, processed, and published by the ABS. The ABS operates within a framework of the Census and Statistics Act 1905 and the Australian Bureau of Statistics Act 1975. These ensure the independence and impartiality from political influence of the ABS, and the confidentiality of respondents.</p> <p>For more information on the institutional environment of the ABS, including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, please see ABS Institutional Environment at www.abs.gov.au.</p>
Relevance	<p>Long-term health conditions described in this publication are classified to a classification developed for use in the NHS (or variants of that classification), based on the ICD. The 2011-12 AHS collected data on self-reported mental and behavioural problems that have lasted for six months, or which the respondent expects to last for six months or more. Estimates for people with 'mental illness' will differ to those that are derived under the SMHWB using the CIDI.</p> <p>The definitions of employment, unemployment and the labour force are consistent with those used in ABS labour force surveys.</p>
Timeliness	<p>The AHS is conducted every three years over a 12 month period. Results from the 2011-12 NHS component of the AHS were released in October 2012.</p>
Accuracy	<p>The AHS is conducted in all States and Territories, excluding very remote areas and discrete Aboriginal and Torres Strait Islander communities. Non-private dwellings such as hotels, motels, hospitals, nursing homes and short-stay caravan parks were also not included in the survey. The exclusion of persons usually residing in very remote areas and discrete Aboriginal and Torres Strait Islander communities has a small impact on estimates, except for the NT where such persons comprise around 25 per cent of the estimated resident population. The response rate for the 2011-12 NHS component was 85 per cent. Results are weighted to account for non-response.</p> <p>As it is drawn from a sample survey, the indicator is subject to sampling error. Sampling error occurs because only a small proportion of the population is used</p>

to produce estimates that represent the whole population. Sampling error can be reliably estimated as it is calculated based on the scientific methods used to design surveys. Rates should be considered with reference to their RSE. Estimates with RSEs between 25 per cent and 50 per cent should be used with caution. Estimates with RSEs greater than 50 per cent are generally considered too unreliable for general use. The attachment tables identify those estimates with RSEs between 25 per cent and 50 per cent.

For information on AHS survey design, see the Australian Health Survey: Users' Guide on the ABS website.

Coherence The methods used to construct the indicator are consistent and comparable with other collections and with international practise.

Accessibility See *Australian Health Survey: First Results (cat. no. 4364.0.55.001)* for an overview of results from the NHS component of the AHS. Other information from this survey is also available on request.

Further information may be available on request. The ABS observes strict confidentiality protocols as required by the *Census and Statistics Act (1905)*. This may restrict access to data at a very detailed level.

Interpretability Information to aid interpretation of the data are available from the Australian Health Survey: Users' Guide on the ABS website.

Many health-related issues are closely associated with age; therefore data for this indicator have been age-standardised to the 2001 total Australian population to account for differences in the age structures of the States and Territories. Age standardised rates should be used to assess the relative differences between groups, not to infer the rates that actually exist in the population.

Data Gaps/Issues Analysis

Key data gaps/issues

The Steering Committee notes the following issues:

- The AHS collects data on self-reported mental and behavioural problems that have lasted for six months, or which the respondent expects to last for six months or more. The data may not be as reliable as or comparable with the data collected under the SMHWB that uses a diagnostic tool to identify mental illnesses.

Social and economic inclusion of people with a mental illness — participation in education and employment by young people

DQI for this indicator has been sourced from the ABS with additional Steering Committee comments.

Indicator definition and description

Element	Outcome
Indicator	Social and economic inclusion of people with a mental illness — participation in education and employment by young people.
Measure (computation)	<p><i>Numerator:</i></p> <p>Number of people aged 16–30 years who are employed and/or are enrolled for study in a formal secondary or tertiary qualification (studying full or part-time) (by mental health status).</p> <p><i>Denominator:</i></p> <p>Number of people in aged 16–30 years in the population (by mental health status).</p> <p><i>Computation:</i></p> <p>$(\text{Numerator} \div \text{Denominator}) * 100$</p> <p>Note: People with a mental health condition are defined as having a self-reported mental and behavioural problems that have lasted for six months, or which the respondent expects to last for six months or more.</p>
Data source/s	ABS unpublished, <i>AHS 2011–13</i> (2011-12 National Health Survey component).

Data Quality Framework Dimensions

Institutional environment	<p>The AHS was collected, processed, and published by the ABS. The ABS operates within a framework of the Census and Statistics Act 1905 and the Australian Bureau of Statistics Act 1975. These ensure the independence and impartiality from political influence of the ABS, and the confidentiality of respondents.</p> <p>For more information on the institutional environment of the ABS, including the legislative obligations of the ABS, financing and governance arrangements, and mechanisms for scrutiny of ABS operations, please see ABS Institutional Environment at www.abs.gov.au.</p>
Relevance	<p>Long-term health conditions described in this publication are classified to a classification developed for use in the NHS (or variants of that classification), based on the ICD. The 2011-12 AHS collected data on self-reported mental and behavioural problems that have lasted for six months, or which the respondent expects to last for six months or more. Estimates for people with 'mental illness' will differ to those that are derived under the SMHWB using the CIDI.</p> <p>The definitions of employment are consistent with those used in ABS labour force surveys.</p>
Timeliness	The AHS is conducted every three years over a 12 month period. Results from the 2011-12 NHS component of the AHS were released in October 2012.
Accuracy	<p>The AHS is conducted in all states and territories, excluding very remote areas. Non-private dwellings such as hotels, motels, hospitals, nursing homes and short-stay caravan parks were also not included in the survey. The exclusion of persons usually residing in very remote areas and discrete Aboriginal and Torres Strait Islander communities has a small impact on estimates, except for the NT where such persons comprise around 25 per cent of the estimated resident population. The response rate for the 2011-12 NHS component was 85 per cent. Results are weighted to account for non-response.</p> <p>As it is drawn from a sample survey, the indicator is subject to sampling error.</p>

Sampling error occurs because only a small proportion of the population is used to produce estimates that represent the whole population. Sampling error can be reliably estimated as it is calculated based on the scientific methods used to design surveys. Rates should be considered with reference to their RSE. Estimates with RSEs between 25 per cent and 50 per cent should be used with caution. Estimates with RSEs greater than 50 per cent are generally considered too unreliable for general use. The attachment tables identify those estimates with RSEs between 25 per cent and 50 per cent.

For information on AHS survey design, see the Australian Health Survey: Users' Guide on the ABS website.

Coherence The methods used to construct the indicator are consistent and comparable with other collections and with international practise.

Accessibility See *Australian Health Survey: First Results (cat. no. 4364.0.55.001)* for an overview of results from the NHS component of the AHS. Other information from this survey is also available on request.

Further information may be available on request. The ABS observes strict confidentiality protocols as required by the *Census and Statistics Act (1905)*. This may restrict access to data at a very detailed level.

Interpretability Information to aid interpretation of the data are available from the Australian Health Survey: Users' Guide on the ABS website.

Many health-related issues are closely associated with age; therefore data for this indicator have been age-standardised to the 2001 total Australian population to account for differences in the age structures of the States and Territories. Age standardised rates should be used to assess the relative differences between groups, not to infer the rates that actually exist in the population.

Data Gaps/Issues Analysis

Key data The Steering Committee notes the following issues:

gaps/issues

- The AHS collects data on self-reported mental and behavioural problems that have lasted for six months, or which the respondent expects to last for six months or more. The data may not be as reliable as, or comparable with, the data collected under the National Survey of Mental Health and Wellbeing that uses a diagnostic tool to identify mental illnesses.

Mental health outcomes of consumers of specialised public mental health services

DQI for this indicator has been sourced from the AMHOCN and Australian, State and Territory governments with additional Steering Committee comments.

Indicator definition and description

Element	Outcome
Indicator	Mental health outcomes of consumers of specialised public mental health services. This DQI should be considered in conjunction with DQI for Collection of information on consumers' outcomes.
Measure (computation)	<p><u>Description:</u></p> <p>Proportion of people receiving care in specialised public mental health services who had a significant improvement in their clinical mental health outcomes. Data are also reported on the proportion who experienced no significant change or a significant deterioration in their mental health outcomes. Data are reported by consumer type: people in ongoing community-based care, people discharged from community-based care and people discharged from a hospital psychiatric inpatient unit.</p> <p><u>Numerator/s:</u></p> <p>Number of people receiving care in specialised public mental health services who had a <u>significant improvement</u> in their clinical mental health outcomes, by consumer type.</p> <p>Number of people receiving care in specialised public mental health services who had <u>no significant change</u> in their clinical mental health outcomes, by consumer type.</p> <p>Number of people receiving care in specialised public mental health services who had a <u>significant deterioration</u> in their clinical mental health outcomes, by consumer type.</p> <p><u>Denominator:</u></p> <p>Number of specialised public mental health service episodes with completed clinical mental health outcome measures data, by consumer type.</p> <p><u>Computation:</u></p> <p>Expressed as a proportion: (Numerator ÷ Denominator)*100. Calculated separately by consumer type.</p>
Data source/s	State and Territory data reported to NOCC and analysed by AMHOCN.

Data Quality Framework Dimensions

Institutional environment	<p>Health Ministers adopted the routine measurement of consumer outcomes as a priority under the <i>National Mental Health Strategy (1992)</i> and in all subsequent National Mental Health Plans. It is also compatible with State and Territory governments' documented policy emphasis on high quality health services and increased consumer and carer participation.</p> <p>The AMHOCN prepared this indicator using the NOCC data on HoNOS family of measures. The Australian Government (Department of Health) contracts AMHOCN to support the implementation of the NOCC as part of routine clinical practice by undertaking three functions 1) data bureau — receives and processes information 2) analysis and reporting — analyses and reports on the submitted data and 3) training and service development — supports training in the measures and their use for clinical practice, service management and development purposes.</p> <p>The NOCC 1.50 was endorsed by all State and Territory governments in 2003, and all jurisdictions have reported data since 2004-05. The NOCC Technical Specification was revised to 1.60 in 2009. All jurisdictions have supplied, or resupplied NOCC data according to 1.60 from 2007-08. The NOCC protocol prescribes a set of standard measures to be collected at particular times (collection occasions) in the clinical process. Under the NOCC protocol, collection of outcomes data is mandatory at admission, review and discharge. Data collected outside of NOCC protocols are</p>
----------------------------------	---

Relevance

excluded from the analysis.

The scope of the NOCC is all specialised public mental health services managed by, or in receipt of funds from, state or territory health authorities Australian Government funded aged residential services are excluded.

The purpose of the NOCC is to measure consumer outcomes. This indicator relates only to consumer outcomes data collected through the HoNOS family of measures (HoNOS; HoNOS for Older People (HoNOS 65+) and HoNOS for Children and Adolescents (HoNOSCA). Other consumer outcome measures are also collected. For adults and older persons these include: Kessler 10 (K10+), Behavior and Symptom Identification Scales (BASIS-32), or Mental Health Inventory (MHI-38); for children and adolescents, the parent and youth versions of the SDQ. The uptake of these measures is not captured by this indicator.

Only episodes that have valid measures for two specified data collection occasions are included. 'Valid' measures are those with a correctly completed specified number of items, for the:

- HoNOS/HoNOS 65+ — a minimum of 10 of the 12 items
- HoNOSCA — a minimum of 11 of the first 13 items.

Brief ambulatory care episodes are excluded from this indicator.

The denominator for the 'completed inpatient' group excludes those episodes that were partially completed within the year and had a length of less than 3 days. The denominator for the 'completed ambulatory' group is made up of those episodes that started and finished within the year. The denominator for the 'ongoing ambulatory' group is made up of those ambulatory episodes that started within the reference year and were still open at 30 June or were open at the start and end of the reference year.

Outcome scores are classified based on effect size — a statistic used to assess the magnitude of a treatment effect. The effect size is the ratio of the difference between the pre- and post- scores to the standard deviation of the pre-score. Individual episodes are classified as 'significant improvement' if the effect size index is greater than or equal to positive 0.5; 'no change' if the index is between -0.5 and 0.5; and 'significant deterioration' if the effect size index is less than or equal to -0.5.

Outcomes are calculated for each of the following three consumer groups and the calculation varies depending on the setting and the duration of the episode of care:

- people discharged from hospital, episodes for people who were admitted and discharged from inpatient care during the reference period (an individual can have two episodes of care so the data represent episode-counts, rather than person-counts) — the admission and discharge occasions rated during the reference period are used
- people in ongoing community-based care, episodes for people who received community care for the whole of the reference period or who commenced community care sometime after 1 July (beginning of the period) and continued to receive care for the rest of the reference period — the first and last occasions rated during the reference period are used
- people discharged from community-based care, episodes for people who were discharged from community care (not including those discharged to hospital) and who received an episode of community care that started and ended in the reference period — the admission and discharge occasions rated during the reference period are used.

Outcomes are measured for consumers discharged from residential mental health care, but there were too few episodes with completed clinical mental health to derive outcome results.

A single 'average score' by consumer type does not reflect the complex service system in which services are delivered across multiple settings (inpatient, community and residential) and provided as both discrete, short term episodes of care and prolonged care over indefinite periods. The approach separates a consumer's care into segments (hospital versus the community) rather than tracking the person's overall outcomes across treatment settings. In addition, consumers' outcomes are measured from the clinician's perspective and not as the 'lived experience' from the consumer's viewpoint.

Data are not available for Victoria for 2011-12 and 2012-13. All Australian totals for 2011-12 and 2012-13 exclude Victoria.

	Tasmanian data for 2009-10 are considered unreliable and not reported. As a result, 2008-09 data are used in the calculation of the 2009-10 Australian coverage estimates.
Timeliness	<p>State and territory health authorities provide the NOCC data to AMHOCN for national collation on a quarterly/annual basis and all data are to be submitted approximately six months after the reference period.</p> <p>The latest reference period for this data set is 2013-14.</p>
Accuracy	<p>States and territories are primarily responsible for the quality of the NOCC data they provide. However, AMHOCN undertakes extensive validation. Validation is conducted in two stages: (1) The compliance stage, concerned with ensuring that the data file will load and is structurally compliant. A non-compliant file is rejected and a new file needs to be submitted. (2) The data validation stage, primarily concerned with identifying and explaining or fixing inconsistent, anomalous, and exceptional issues in relation to the NOCC protocol as well as flagging , including invalid domain values and/or, missing data.</p> <p>The proportion of episodes for which 'valid' outcomes data are collected is less than 50 per cent of expected coverage. It is not known if the results for those for whom data are collected are representative of the consumer population.</p>
Coherence	<p>Data are available for 2007-08 to 2012-13. The comparability of the outcomes data across jurisdictions and years may be affected by the relatively low proportion of episodes for which 'valid' outcomes data are collected and the degree to which this proportion varies across jurisdictions and years.</p> <p>The Australian totals for 2011-12 and 2012-13 are not comparable to other years as they exclude data for Victoria.</p>
Accessibility	<p>Data for this indicator are published in the National mental health reports: www.health.gov.au/internet/main/publishing.nsf/Content/mental-data and in the Key Performance Indicators for Australian Public Mental Health Services set (available at mhsa.aihw.gov.au/indicators/nkpi/).</p> <p>NOCC data are available on the AMHOCN website amhocn.org/. The following on-line products are available:</p> <ul style="list-style-type: none"> • web decision support tool • NOCC Standard Reports • NOCC Volume and Percentage Clinical Ratings: Australia
Interpretability	<p>Metadata information for the NOCC are published on the AMHOCN website amhocn.org/.</p> <p>Further information to understand this indicator are available:</p> <ul style="list-style-type: none"> • in the Key Performance Indicators for Australian Public Mental Health Services, Third Edition • from the Key Performance Indicators for Australian Public Mental Health Services (available at mhsa.aihw.gov.au/indicators/nkpi/).

Data Gaps/Issues Analysis

Key data gaps/issues	<p>The Steering Committee notes the following key data gaps/issues:</p> <ul style="list-style-type: none"> • Data are not provided for the latest reference period (2014-15). Further work is required to ensure availability of more timely data. • There are differences in the relative proportions of 'matched pair' HoNOS/CA/65+ ratings. <ul style="list-style-type: none"> – NOCC completion rates for people discharged from hospital and people in ongoing community based care are approximately 85 per cent. – NOCC completion rates for people discharged from community based care, are lower, at approximately 65 per cent. This pattern has been stable over time and generally consistent for all consumer age groups and jurisdictions, with the exception of ACT where technical issues have not enabled linkage of admission and discharge ratings for this consumer group. It is likely that the overall lower completion rate for this consumer group arises when consumers are administratively discharged from care following a period of no active care in the preceding period.
-----------------------------	---