

5A Early child development and growth — attachment

The tables in this file accompany the Report, *Overcoming Indigenous Disadvantage: Key Indicators 2007*, prepared by the Steering Committee for the Review of Government Service Provision. Background and definitions are available in the Report, which is available on the Review website (www.pc.gov.au/gsp).

This file is available in both Microsoft Excel and Adobe PDF formats on the Review website (www.pc.gov.au/gsp). Users without Internet access can contact the Secretariat to obtain these tables (details inside the front cover of the Report).

Contents

5A.1 Injury and preventable diseases

Table 5A.1.1 Number and rate of hospitalisations for injury and potentially preventable diseases for children aged less than 4 years, Qld, WA, SA, and public hospitals in the NT, 2004-05

Table 5A.1.2 Deaths from external causes and preventable diseases for children aged less than 5 years, 2001-2005

5A.2 Infant mortality

Table 5A.2.1 Infant mortality rates (per 1000 live births)

5A.3 Birthweight

Table 5A.3.1 Baby's birthweight, by live births and fetal deaths, 2002-2004

Table 5A.3.2 Baby's birthweight, by live births and fetal deaths, 2001-2003

Table 5A.3.3 Baby's birthweight, by live births and fetal deaths, 2000-2002

Table 5A.3.4 Baby's birthweight, by live births and fetal deaths, 1999-2001

Table 5A.3.5 Baby's birthweight, by live births and fetal deaths, 1998-2000

5A.4 Hearing impediments

Table 5A.4.1 Persons with a long-term hearing problem: type of ear/hearing problem, by age and Indigenous status, 2004-05 (per cent of population)

Table 5A.4.2 Persons with a long-term hearing problem: type of ear/hearing problem, by age and Indigenous status, 2001 (per cent of population)

Table 5A.4.3 Persons with a long-term hearing problem: type of ear/hearing problem, by remoteness areas and Indigenous status, 2004-05 (per cent of population)

Table 5A.4.4 Persons with a long-term hearing problem: type of ear/hearing problem, by remoteness areas and Indigenous status, 2001 (per cent of population)

Table 5A.4.5 Persons with otitis media, by age and Indigenous status, 2004-05 (per cent of population)

Table 5A.4.6 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA, and public hospitals in NT, 2004-05

Table 5A.4.7 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA, and public hospitals in NT, 2003-04

Table 5A.4.8 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA and public hospitals in NT, 2002-03

Table 5A.4.9 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA and public hospitals in NT, 2001-02

5A.5 Children with tooth decay

Table 5A.5.1 Decayed and decay-free infant teeth in 4- to 10-year-old children enrolled in school dental services in NSW, SA and the NT, by Indigenous status and location

Table 5A.5.2 Decayed and decay-free permanent teeth in 6- to 14-year-old children enrolled in school dental services in NSW, SA and the NT, by Indigenous status and location

Table 5A.5.3 Mean number of teeth with decay for children enrolled in the NT School Dental Service, by age and Indigenous status, 2002-03

Table 5A.5.4 Mean tooth decay (dmft) for 4- to 10-year-old children enrolled in the NT School Dental Service, by SEIFA index of relative socio-economic disadvantage, age and Indigenous status, 2002-03

Table 5A.5.5 Mean tooth decay (DMFT) for 6- to 13-year-old children enrolled in the NT School Dental Service, by SEIFA index of relative socio-economic disadvantage, age and Indigenous status, 2002-03

Contents

-
- Table 5A.5.6** SA, infant and permanent tooth decay, mean dmft and DMFT by Indigenous status and age, 2002 Child Dental Health Survey
- Table 5A.5.7** SA, proportion of children with calculus and gingival bleeding by Indigenous status and age, children attending school dental service, 2002 Child Dental Health Survey
- Table 5A.5.8** Dental hospitalisations for children aged 2 to 14 years, per 100 000 children, by sociodemographic characteristics and Indigenous status, 2002-03
- Table 5A.5.9** Hospital dental procedure rates for children aged 2 to 14 years, per 100 000 children, by sex and Indigenous status, 2002-03
- Table 5A.5.10** Hospital dental procedure rates for children aged 2 to 14 years, per 100 000 children, by age group and Indigenous status, 2002-03
- Table 5A.5.11** Hospital dental procedure rates for children aged 2 to 14 years, per 100 000 children, by remoteness area and Indigenous status, 2002-03

Table 5A.1.1

Number and rate of hospitalisations for injury and potentially preventable diseases for children aged less than 4 years, Qld, WA, SA, and public hospitals in the NT, 2004-05 (a), (b)

ICD-10-AM code and description	Separations (no.)			Age specific rate (c)			Rate ratio (d)
	Indigenous	non- Indigenous (d)	Total	Indigenous	non- Indigenous (d)		
Qld							
Certain infectious and parasitic diseases (A00–B99)	429	3 853	4 282	31.1	20.7	1.5	
Nutritional anaemias (D50–D53) and malnutrition (E40–E46)	np	np	30	np	np	np	
Diseases of the ear and mastoid process (H60–H95)	np	np	2 409	np	np	0.8	
Diseases of the respiratory system (J00–J99)	1 089	9 575	10 664	79.0	51.4	1.5	
Diseases of oral cavity, salivary glands and jaws (K00–K14)	158	1 090	1 248	11.5	5.9	2.0	
Diseases of the skin and subcutaneous tissue (L00–L99)	207	618	825	15.0	3.3	4.5	
Injury, poisoning and certain other consequences of external causes (S00–T98)	397	4 051	4 448	28.8	21.7	1.3	
Transport accidents (V01–V99)	29	176	205	2.1	0.9	2.2	
Other external causes of accidental injury (W00–X59)	292	3 475	3 767	21.2	18.7	1.1	
Assault (X85–Y09)	22	42	64	1.6	0.2	7.1	
Complications of medical and surgical care (Y40–Y84)	45	347	392	3.3	1.9	1.8	
Other	9	11	20	0.7	0.1	11.1	
Total potentially preventable hospitalisations	2 419	21 487	23 906	175.4	115.3	1.5	
WA							
Certain infectious and parasitic diseases (A00–B99)	499	1 993	2 492	71.7	21.8	3.3	
Nutritional anaemias (D50–D53) and malnutrition (E40–E46)	23	7	30	3.3	0.1	43.1	
Diseases of the ear and mastoid process (H60–H95)	96	1 456	1 552	13.8	15.9	0.9	
Diseases of the respiratory system (J00–J99)	1 137	4 709	5 846	163.4	51.5	3.2	
Diseases of oral cavity, salivary glands and jaws (K00–K14)	67	628	695	9.6	6.9	1.4	
Diseases of the skin and subcutaneous tissue (L00–L99)	139	355	494	20.0	3.9	5.1	
Injury, poisoning and certain other consequences of external causes (S00–T98)	203	1 689	1 892	29.2	18.5	1.6	
Transport accidents (V01–V99)	17	70	87	2.4	0.8	3.2	
Other external causes of accidental injury (W00–X59)	158	1 485	1 643	22.7	16.3	1.4	
Assault (X85–Y09)	21	22	43	3.0	0.2	12.5	
Complications of medical and surgical care (Y40–Y84)	np	105	np	np	1.1	np	
Other	np	7	np	np	0.1	np	
Total potentially preventable hospitalisations	2 164	10 837	13 063	311.0	118.6	2.6	

Table 5A.1.1

Number and rate of hospitalisations for injury and potentially preventable diseases for children aged less than 4 years, Qld, WA, SA, and public hospitals in the NT, 2004-05 (a), (b)

ICD-10-AM code and description	Separations (no.)			Age specific rate (c)			Rate ratio
	Indigenous	non-Indigenous (d)	Total	Indigenous	non-Indigenous (d)		
SA							
Certain infectious and parasitic diseases (A00–B99)	164	1 766	1 930	63.0	26.2	2.4	
Nutritional anaemias (D50–D53) and malnutrition (E40–E46)	11	7	18	4.2	0.1	40.7	
Diseases of the ear and mastoid process (H60–H95)	39	2 057	2 096	15.0	30.5	0.5	
Diseases of the respiratory system (J00–J99)	309	4 747	5 056	118.8	70.4	1.7	
Diseases of oral cavity, salivary glands and jaws (K00–K14)	27	409	436	10.4	6.1	1.7	
Diseases of the skin and subcutaneous tissue (L00–L99)	25	251	276	9.6	3.7	2.6	
Injury, poisoning and certain other consequences of external causes (S00–T98)	85	1 127	1 212	32.7	16.7	2.0	
Transport accidents (V01–V99)	6	33	39	2.3	0.5	4.7	
Other external causes of accidental injury (W00–X59)	58	991	1 049	22.3	14.7	1.5	
Assault (X85–Y09)	np	np	np	np	np	np	
Complications of medical and surgical care (Y40–Y84)	6	86	92	2.3	1.3	1.8	
Other	np	np	np	np	np	np	
Total potentially preventable hospitalisations	660	10 364	13 063	253.7	153.7	1.7	
Public hospitals in the NT							
Certain infectious and parasitic diseases (A00–B99)	639	180	819	114.6	21.0	5.4	
Nutritional anaemias (D50–D53) and malnutrition (E40–E46)	np	np	69	np	np	np	
Diseases of the ear and mastoid process (H60–H95)	np	np	59	np	np	1.9	
Diseases of the respiratory system (J00–J99)	881	254	1 135	158.0	29.7	5.3	
Diseases of oral cavity, salivary glands and jaws (K00–K14)	45	19	64	8.1	2.2	3.6	
Diseases of the skin and subcutaneous tissue (L00–L99)	223	49	272	40.0	5.7	7.0	
Injury, poisoning and certain other consequences of external causes (S00–T98)	130	136	266	23.3	15.9	1.5	
Transport accidents (V01–V99)	10	7	17	1.8	0.8	2.2	
Other external causes of accidental injury (W00–X59)	94	116	210	16.9	13.6	1.2	
Assault (X85–Y09)	np	np	np	np	np	np	
Complications of medical and surgical care (Y40–Y84)	np	np	np	np	np	np	
Other	np	np	np	np	np	np	
Total potentially preventable hospitalisations	2 017	667	13 063	361.8	78.0	4.6	

Table 5A.1.1

Table 5A.1.1 Number and rate of hospitalisations for injury and potentially preventable diseases for children aged less than 4 years, Qld, WA, SA, and public hospitals in the NT, 2004-05 (a), (b)

ICD-10-AM code and description	Separations (no.)			Age specific rate (c)		
	Indigenous	non-Indigenous (d)	Total	Indigenous	non-Indigenous (d)	Rate ratio
Qld, WA, SA, and public hospitals in the NT						
Certain infectious and parasitic diseases (A00–B99)	1 731	7 792	9 523	59.8	22.0	2.7
Nutritional anaemias (D50–D53) and malnutrition (E40–E46)	104	43	147	3.6	0.1	29.6
Diseases of the ear and mastoid process (H60–H95)	303	5 813	6 116	10.5	16.4	0.6
Diseases of the respiratory system (J00–J99)	3 416	19 285	22 701	118.1	54.5	2.2
Diseases of oral cavity, salivary glands and jaws (K00–K14)	297	2 146	2 443	10.3	6.1	1.7
Diseases of the skin and subcutaneous tissue (L00–L99)	594	1 273	1 867	20.5	3.6	5.7
Injury, poisoning and certain other consequences of external causes (S00–T98)	815	7 003	7 818	28.2	19.8	1.4
Transport accidents (V01–V99)	62	286	348	2.1	0.8	2.7
Other external causes of accidental injury (W00–X59)	602	6 067	6 669	20.8	17.2	1.2
Assault (X85–Y09)	73	79	152	2.5	0.2	11.3
Complications of medical and surgical care (Y40–Y84)	66	551	617	2.3	1.6	1.5
Other	12	20	32	0.4	0.1	7.3
Total potentially preventable hospitalisations	7 260	43 355	13 063	251.0	122.6	2.0

(a) Data are based on principal diagnosis as classified by the ICD-10-AM code and description.

(b) Data are based on state of usual residence.

(c) Age specific rates are as per 1000 people in that age group (based on ABS 2002 ERP).

(d) Separations where Indigenous status was not stated are included in the non-Indigenous numbers and rates.
np Not published.

Source: AIHW National Hospital Morbidity Database (unpublished).

Table 5A.1.2

Deaths from external causes and preventable diseases for children aged less than 5 years, 2001–2005 (a)

	Indigenous					Non-Indigenous				
	NSW	Qld	WA	SA	NT	NSW	Qld	WA	SA	NT
External causes and preventable diseases	no.	25	48	37	12	38	283	211	100	59
All causes	no.	156	232	143	34	155	2218	1331	524	413
Diseases as a percentage of all causes	%	16.0	20.7	25.9	35.3	24.5	12.8	15.9	19.1	14.3
Death rate per 10 000 children aged less than 5 years		np	5.7	9.3	7.6	10.8	np	1.8	1.7	1.4
2003 Population aged less than 5 years	no.	np	16 966	7 928	3 163	7 056	np	np	np	np

(a) Calculations of rates for the Indigenous population are based on ABS Experimental Projections, Aboriginal and Torres Strait Islander Australians (low series, 2001 base). There are no comparable population data for the non-Indigenous population. Calculations of rates for the non-Indigenous population are based on data derived by subtracting Indigenous population projections from total population estimates and should be used with care.

np Not published.

Source: ABS *Causes of Death* 2005, Cat. no. 3303.0 (unpublished).

Table 5A.2.1

Table 5A.2.1 Infant mortality rates (per 1000 live births) (a), (b), (c)

	NSW	Qld	WA	SA	NT
Indigenous					
1997-99	10.4	12.6	15.8	6.9	23.5
1998-00	11.8	12.5	16.9	7.8	21.5
1999-01	10.9	11.7	16.6	8.0	19.2
2000-02	9.5	11.5	16.5	10.4	18.1
2001-03	8.6	11.2	15.9	9.1	14.8
2002-04	8.5	10.9	14.1	9.4	15.4
2003-05	8.4	10.9	12.8	7.7	15.6
Total persons					
1997-99	5.1	6.0	5.0	4.4	12.2
1998-00	5.1	6.1	4.7	4.3	11.9
1999-01	5.4	5.9	4.7	4.5	11.4
2000-02	5.0	6.0	4.6	4.8	11.2
2001-03	4.8	5.5	4.5	4.5	10.1
2002-04	4.6	5.3	4.1	4.0	10.1
2003-05	4.7	5.0	4.2	4.0	9.5
Indigenous, by sex, 2003-05					
Males	8.8	14.6	13.9	7.1	21.2
Females	7.9	6.9	11.6	8.3	9.5

- (a) Victoria, Tasmania and the ACT are excluded due to poor coverage rates or small numbers.
- (b) Deaths for whom Indigenous origin was not stated have not been prorated between Indigenous and non-Indigenous deaths. As a result, Indigenous and non-Indigenous infant mortality rates may be underestimated.
- (c) Contribution of Indigenous deaths to total deaths is much larger in the NT than in other states.

Source: ABS 2006, *Deaths Australia, 2005* (and previous issues), Cat. no. 3302.0, Canberra.

Table 5A.3.1

Baby's birthweight, by live births and fetal deaths, 2002–2004 (a), (b)

Australia	Aboriginal and Torres Strait Islander (c)		Non-Indigenous (c)		Unknown Indigenous status (c)		All births (d)	
	Live births		Fetal deaths (e)		Unknown		Unknown	
	no.	%	no.	%	no.	%	no.	%
Australia								
Low birthweight (<2500g)	3 459	13.0	241	78.2	3 700	13.8	43 974	6.1
Very low birthweight (<1500g)	630	2.4	206	66.9	836	3.1	7 275	1.0
Extremely low birthweight (<1000g)	286	1.1	184	59.7	480	1.8	3 182	0.4
All births	26 583	100.0	308	100.0	28 891	100.0	729 309	100.0
Mean birthweight (g)	3 161	..	1 300	..	3 140	..	3 380	..
NSW								
Low birthweight (<2500g)	805	12.1	45	72.6	850	12.7	14 367	5.8
Very low birthweight (<1500g)	120	1.8	36	58.1	156	2.3	2 303	0.9
Extremely low birthweight (<1000g)	45	0.7	30	48.4	75	1.1	982	0.4
All births	6 643	100.0	62	100.0	6 705	100.0	249 682	100.0
Mean birthweight (g)	3 196	..	1 398	..	3 180	..	3 386	..
Victoria								
Low birthweight (<2500g)	174	14.3	17	89.5	191	15.5	11 948	6.4
Very low birthweight (<1500g)	32	2.6	16	84.2	48	3.9	2 072	1.1
Extremely low birthweight (<1000g)	16	1.3	15	78.9	31	2.5	969	0.5
All births	1 215	100.0	19	100.0	1 234	100.0	187 564	100.0
Mean birthweight (g)	3 143	..	772	..	3 108	..	3 370	..
Queensland								
Low birthweight (<2500g)	974	11.7	71	81.6	1 045	12.4	8 949	6.3
Very low birthweight (<1500g)	193	2.3	62	71.3	255	3.0	1 525	1.1
Extremely low birthweight (<1000g)	87	1.0	57	65.5	144	1.7	659	0.5
All births	8 354	100.0	87	100.0	8 441	100.0	141 112	100.0
Mean birthweight (g)	3 196	..	1 243	..	3 176	..	3 390	..

Table 5A.3.1

Table 5A.3.1

Table 5A.3.1

Baby's birthweight, by live births and fetal deaths, 2002-2004 (a), (b)

	Aboriginal and Torres Strait Islander (c)					Non-Indigenous (c)					Unknown Indigenous status (c)					All births (d)											
	Live births		Fetal deaths (e)		Total births	Live births		Fetal deaths (e)		Total births	Live births		Fetal deaths (e)		Total births	Live births		Fetal deaths (e)		Unknown							
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.						
WA																											
Low birthweight (<2500g)	652	14.1	53	77.9	705	15.0	4 112	5.9	384	80.2	-	4 496	6.4	-	-	-	4 764	6.4	437	79.9	-	5 201	6.9				
Very low birthweight (<1500g)	113	2.4	47	69.1	160	3.4	614	0.9	333	69.5	-	947	1.3	-	-	-	-	727	1.0	380	69.5	-	1 107	1.5			
Extremely low birthweight (<1000g)	57	1.2	41	60.3	98	2.1	247	0.4	298	62.2	-	545	0.8	-	-	-	-	304	0.4	339	62.0	-	643	0.9			
All births	4 633	100.0	68	100.0	4 701	100.0	69 813	100.0	479	100.0	-	70 292	100.0	-	-	-	-	74 446	100.0	547	100.0	-	74 993	100.0			
Mean birthweight (g)	3 132	..	1 318	..	3 106	..	3 368	..	1 214	..	-	3 354	..	-	-	-	-	3 354	..	1 227	..	-	3 338	..			
SAs																											
Low birthweight (<2500g)	250	17.9	15	78.9	265	18.7	3 208	6.2	262	74.9	-	3 470	6.7	-	-	-	-	-	3 458	6.6	277	75.1	-	3 735	7.0		
Very low birthweight (<1500g)	63	4.5	13	68.4	76	5.4	532	1.0	222	63.4	-	754	1.5	-	-	-	-	-	595	1.1	235	63.7	-	830	1.6		
Extremely low birthweight (<1000g)	29	2.1	13	68.4	42	3.0	222	0.4	195	55.7	-	417	0.8	-	-	-	-	-	251	0.5	208	56.4	-	459	0.9		
All births	1 395	100.0	19	100.0	1 414	100.0	51 346	100.0	350	100.0	-	51 696	100.0	-	-	-	-	-	52 741	100.0	369	100.0	-	53 110	100.0		
Mean birthweight (g)	3 055	..	1 243	..	3 031	..	3 370	..	1 396	..	-	3 357	..	-	-	-	-	-	3 362	..	1 379	..	-	3 348	..		
Tas																											
Low birthweight (<2500g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na					
Very low birthweight (<1500g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na					
Extremely low birthweight (<1000g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na					
All births	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na					
Mean birthweight (g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na					
ACT (f)																											
Low birthweight (<2500g)	np	np	<5	np	42	18.4	np	np	np	np	-	-	1 087	7.6	4	6.5	-	0.0	-	4	6.3	1 032	7.1	101	82.1		
Very low birthweight (<1500g)	np	np	<5	np	16	7.0	np	np	np	np	-	-	254	1.8	-	0.0	-	-	-	184	1.3	86	69.9	-	-	270	1.8
Extremely low birthweight (<1000g)	np	np	<5	np	10	4.4	np	np	np	np	-	-	157	1.1	-	0.0	-	-	0.0	90	0.6	77	62.6	-	-	167	1.1
All births	np	np	<5	np	228	100.0	np	np	np	np	-	-	14 315	100.0	62	100.0	1	100.0	-	63	100.0	14 483	100.0	-	-	14 606	100.0
Mean birthweight (g)	3 116	..	489	..	3 070	..	3 397	..	1 211	..	-	3 379	..	-	-	-	-	-	3 392	..	2 590	..	-	-	3 374	..	

Table 5A.3.1

Baby's birthweight, by live births and fetal deaths, 2002-2004 (a), (b)

NT	Aboriginal and Torres Strait Islander (c)										Non-Indigenous (c)										Unknown Indigenous status (c)										All births (d)									
	Live births					Fetal deaths (e)					Total births					Live births					Fetal deaths (e)					Unknown					Total births									
	no.	%	no.	%	no.	no.	%	no.	%	no.	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%		
Low birthweight (<2500g)	566	13.7	36	73.5	602	14.4	400	6.1	34	77.3	-	-	434	6.5	10	20.8	-	-	-	-	10	20.8	976	9.1	70	75.3	-	-	-	-	1046	9.6								
Very low birthweight (<1500g)	97	2.4	28	57.1	125	3.0	57	0.9	28	63.6	-	-	85	1.3	3	6.3	-	-	-	-	3	6.3	157	1.5	56	60.2	-	-	-	-	213	2.0								
Extremely low birthweight (<1000g)	56	1.4	24	49.0	80	1.9	19	0.3	25	56.8	-	-	44	0.7	3	6.3	-	-	-	-	3	6.3	78	0.7	49	52.7	-	-	-	-	127	1.2								
All births	4 119	100.0	49	100.0	4 168	100.0	6 595	100.0	44	100.0	-	-	6 639	100.0	48	100.0	-	-	-	-	48	100.0	10 762	100.0	93	100.0	-	-	-	-	10 855	100.0								
Mean birthweight (g)	3 109	..	1548	..	3 091	..	3 363	..	1 337	3 350	..	3 094	3 094	..	3 265	..	1 447	3 249	..								

(a) Data are presented in a three year grouping due to small numbers from year to year. Tasmania not available because the not stated category for Indigenous status was not able to be distinguished from the non-Indigenous category.

(b) Data on Indigenous births relate to babies born to Indigenous mothers only, and excludes babies born to non-Indigenous mothers and Indigenous fathers. Thus, the information is not based on the total count of Indigenous babies.

(c) Totals from the Australian sections of births broken-down by Indigenous status do not include data for Tasmania.

(d) The Australian all births section totals include data for Tasmania.

(e) The denominator for the fetal death percentages is fetal deaths rather than births.

(f) ACT data includes non-ACT residents who gave birth in the ACT.

(g) n/a Not available, .. Not applicable, - Nil or rounded to zero, np Not published.

Source: National Perinatal Data Collection, AIHW National Perinatal Statistics Unit (unpublished).

Table 5A.3.2

Table 5A.3.2 Baby's birthweight, by live births and fetal deaths, 2001–2003 (a), (b)

	Non-Indigenous (c)												Unknown Indigenous status (c)												All births (d)											
	Aboriginal and Torres Strait Islander (c)						Live births Fetal deaths (e)						Unknown						Live births Fetal deaths (e)						Unknown											
	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%						
Australia																																				
Low birthweight (<2500g)	3 403	12.9	244	75.1	3 647	13.7	43 475	6.1	3 671	76.2	—	—	47 146	6.5	19	7.0	2	66.7	—	—	21	7.6	48 068	6.3	4 029	76.2	—	—	52 097	6.8						
Very low birthweight (<1500g)	614	2.3	197	60.6	811	3.0	7 267	1.0	3 083	64.0	—	—	10 350	1.4	4	1.5	1	33.3	—	—	5	1.8	8 072	1.1	3 370	63.7	—	—	11 442	1.5						
Extremely low birthweight (<1000g)	284	1.1	176	54.2	460	1.7	3 176	0.4	2 732	56.7	—	—	5 908	0.8	3	1.1	1	33.3	—	—	4	1.4	3 535	0.5	2 985	56.5	—	—	6 520	0.9						
All births	26 553	100.0	325	100.0	26 678	100.0	717 635	100.0	4 815	100.0	—	—	722 450	100.0	273	10.0	3	100.0	—	—	276	100.0	761 059	100.0	5 287	100.0	—	—	766 346	100.0						
Mean birthweight (g)	3 136	..	1 450	..	3 143	..	3 380	..	1 326	..	—	..	3 367	..	3 380	..	1 705	..	—	..	3 361	..	3 373	..	1 394	..	—	..	3 359	..						
NSW																																				
Low birthweight (<2500g)	784	12.2	52	732	836	12.8	14 451	5.8	1 108	73.7	—	—	15 559	6.2	5	3.7	1	100.0	—	—	6	4.4	15 240	5.9	1 161	73.7	—	—	16 401	6.4						
Very low birthweight (<1500g)	120	1.9	39	54.9	159	2.4	2 323	0.9	896	59.6	—	—	3 219	1.3	—	0.0	—	0.0	—	—	—	0.0	2 443	1.0	935	59.3	—	—	3 378	1.3						
Extremely low birthweight (<1000g)	46	0.7	33	46.5	79	1.2	1 021	0.4	786	52.3	—	—	1 807	0.7	—	0.0	—	0.0	—	—	—	0.0	1 067	0.4	819	52.0	—	—	1 886	0.7						
All births	6 439	100.0	71	100.0	6 510	100.0	250 128	100.0	1 504	100.0	—	—	251 632	100.0	134	100.0	1	100.0	—	—	135	100.0	256 701	100.0	1 576	100.0	—	—	258 277	100.0						
Mean birthweight (g)	3 192	..	1 435	..	3 174	..	3 387	..	1 409	..	—	..	3 375	..	3 499	..	2 074	..	—	..	3 488	..	3 382	..	1 411	..	—	..	3 370	..						
Victoria																																				
Low birthweight (<2500g)	152	12.7	18	818	170	14.0	11 814	6.3	1 108	77.9	—	—	12 922	6.9	—	1	100.0	—	—	1	100.0	—	—	1 100	11 966	6.4	1 127	78.0	—	—	13 093	6.9				
Very low birthweight (<1500g)	25	2.1	16	727	41	3.4	2 064	1.1	960	67.5	—	—	3 024	1.6	—	1	100.0	—	—	1	100.0	—	—	1	2 089	1.1	977	67.6	—	—	3 066	1.6				
Extremely low birthweight (<1000g)	12	1.0	15	682	27	2.2	953	0.5	865	60.8	—	—	1 818	1.0	—	—	1	100.0	—	—	1	100.0	—	—	1	965	0.5	881	61.0	—	—	1 846	1.0			
All births	1 194	100.0	22	100.0	1 216	100.0	186 195	100.0	1 422	100.0	—	—	187 617	100.0	—	—	1	100.0	—	—	1	100.0	—	—	187 389	100.0	1 445	100.0	—	—	188 834	100.0				
Mean birthweight (g)	3 176	..	1 210	..	3 141	..	3 368	..	1 228	..	—	..	3 352	..	—	..	450	..	—	..	450	..	—	..	3 366	..	1 227	..	—	..	3 350	..				

Table 5A.3.2

Table 5A.3.2 Baby's birthweight, by live births and fetal deaths, 2001–2003 (a), (b)

Queensland	Non-Indigenous (c)												Unknown Indigenous status (c)												All births (d)											
	Aboriginal and Torres Strait Islander (c)						Live births Fetal deaths (e)						Unknown						Live births Fetal deaths (e)						Unknown											
	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%	Live births no.	%	Fetal deaths no.	%	Total births no.	%						
Low birthweight (<2500g)	956	11.5	71	78.9	1 027	12.3	8 671	6.2	687	75.6	—	—	9 358	6.6	3	10.7	—	—	3	10.7	9 630	6.5	758	75.9	—	—	10 388	7.0								
Very low birthweight (<1500g)	188	2.3	55	61.1	243	2.9	1 500	1.1	579	63.7	—	—	2 079	1.5	1	3.6	—	—	1	3.6	1 689	1.1	634	63.5	—	—	2 323	1.6								
Extremely low birthweight (<1000g)	91	1.1	50	55.6	141	1.7	609	0.4	504	55.4	—	—	1 113	0.8	—	0.0	—	—	—	—	—	700	0.5	554	55.5	—	—	1 254	0.8							
All births	8 278	100.0	90	100.0	8 368	100.0	139 947	100.0	909	100.0	—	—	140 856	100.0	28	100.0	—	—	28	100.0	148 253	100.0	998	100.0	—	—	149 252	100.0								
Mean birthweight (g)	3 204	..	1 474	..	3 186	..	3 394	..	1 371	..	—	—	3 381	..	3 255	..	—	—	3 255	..	3 383	..	1 381	..	—	—	3 370	..								
WA																																				
Low birthweight (<2500g)	675	14.5	48	72.7	723	15.3	4 042	5.8	370	80.6	—	—	4 412	6.3	—	—	—	—	—	—	—	—	—	4 717	6.4	418	79.6	—	—	5 135	6.9					
Very low birthweight (<1500g)	124	2.7	42	63.6	166	3.5	590	0.9	321	69.9	—	—	911	1.3	—	—	—	—	—	—	—	—	—	714	1.0	363	69.1	—	—	1 077	1.4					
Extremely low birthweight (<1000g)	61	1.3	37	56.1	98	2.1	233	0.3	284	61.9	—	—	517	0.7	—	—	—	—	—	—	—	—	—	294	0.4	321	61.1	—	—	616	0.8					
All births	4 661	100.0	66	100.0	4 727	100.0	69 218	100.0	459	100.0	—	—	69 677	100.0	—	—	—	—	—	—	—	—	—	—	73 879	100.0	525	100.0	—	—	74 404	100.0				
Mean birthweight (g)	3 123	..	1 480	..	3 100	..	3 368	..	1 213	..	—	—	3 354	..	—	—	—	—	—	—	—	—	—	—	3 352	..	1 246	..	—	—	3 338	..				
SA																																				
Low birthweight (<2500g)	229	17.6	15	682	244	18.4	3 193	6.2	268	75.7	—	—	3 461	6.7	—	—	—	—	—	—	—	—	—	—	3 422	6.5	283	75.3	—	—	3 705	7.0				
Very low birthweight (<1500g)	56	4.3	13	59.1	69	5.2	557	1.1	224	63.3	—	—	781	1.5	—	—	—	—	—	—	—	—	—	—	613	1.2	237	63.0	—	—	850	1.6				
Extremely low birthweight (<1000g)	24	1.8	13	59.1	37	2.8	256	0.5	199	56.2	—	—	455	0.9	—	—	—	—	—	—	—	—	—	—	280	0.5	212	56.4	—	—	492	0.9				
All births	1 301	100.0	22	100.0	1 323	100.0	51 616	100.0	354	100.0	—	—	51 970	100.0	—	—	—	—	—	—	—	—	—	—	52 917	100.0	376	100.0	—	—	53 293	100.0				
Mean birthweight (g)	3 060	..	1 496	..	3 034	..	3 372	..	1 381	..	—	—	3 358	..	—	—	—	—	—	—	—	—	—	—	3 364	..	1 388	..	—	—	3 350	..				

Table 5A.3.2

Baby's birthweight, by live births and fetal deaths, 2001–2003 (a), (b)

Table 5A.3.2

Tas	Aboriginal and Torres Strait Islander (c)						Non-Indigenous (c)						Unknown Indigenous status (c)						All births (d)								
	Live births			Fetal deaths (e)			Total births			Live births			Fetal deaths (e)			Total births			Live births			Fetal deaths (e)			Total births		
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	
Low birthweight (<2500g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na		
Very low birthweight (<1500g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na		
Extremely low birthweight (<1000g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na		
All births	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na		
Mean birthweight (g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na		
ACT (f)																											
Low birthweight (<2500g)	np	np	<5	np	43	20.7	np	np	np	np	np	np	np	np	np	np	np	np	3	5.4	0.0	—	—	3	5.3		
Very low birthweight (<1500g)	np	np	<5	np	16	7.7	np	np	np	np	np	np	np	np	np	np	np	np	—	—	—	—	189	1.3	77		
Extremely low birthweight (<1000g)	np	np	<5	np	10	4.8	np	np	np	np	np	np	np	np	np	np	np	np	—	0.0	0.0	—	—	91	0.6	70	
All births	np	np	<5	np	208	100.0	np	np	np	np	np	np	np	np	np	np	np	np	56	100.0	1	100.0	57	100.0	125	100.0	
Mean birthweight (g)	3 085	..	3 036	..	3 397	..	1 398	..	3 380	..	3 373	..	2 590	..	2 580	..	2 570	..	3 359	..	3 392	..	1 379	..	3 374	..	
NT																											
Low birthweight (<2500g)	568	13.3	36	72.0	604	14.0	402	6.0	37	78.7	—	—	439	6.5	8	14.5	—	—	—	8	14.5	97.8	8.8	73	75.3		
Very low birthweight (<1500g)	89	2.1	28	56.0	117	2.7	56	0.8	30	63.8	—	—	86	1.3	3	5.5	—	—	—	3	5.5	148	1.3	58	59.8		
Extremely low birthweight (<1000g)	44	1.0	24	48.0	68	1.6	19	0.3	28	59.6	—	—	47	0.7	3	5.5	—	—	—	3	5.5	66	0.6	52	53.6		
All births	4 276	100.0	50	100.0	4 326	100.0	6 723	100.0	47	100.0	—	—	6 770	100.0	55	100.0	—	—	—	56	100.0	11 054	100.0	97	100.0		
Mean birthweight (g)	3 118	..	1 562	..	3 100	..	3 362	..	1 291	..	—	—	—	—	—	—	—	—	3 348	..	3 170	..	3 267	..	1 424	..	
(a)	Data are presented in a three year grouping due to small numbers from year to year. Tasmania not available because the not stated category for Indigenous status was not able to be distinguished from the non-Indigenous category.																										
(b)	Data on Indigenous births relate to babies born to Indigenous mothers only, and excludes babies born to non-Indigenous mothers and Indigenous fathers. Thus, the information is not based on the total count of Indigenous babies.																										
(c)	Totals from the Australian sections of births broken-down by Indigenous status do not include data for Tasmania.																										
(d)	The Australian all births section totals include data for Tasmania.																										
(e)	The denominator for the fetal death percentages is fetal deaths rather than births.																										
(f)	ACT data includes non-ACT residents who gave birth in the ACT.																										
na	Not available. .. Not applicable – Nil or rounded to zero. np Not published.																										
Source:	National Perinatal Data Collection, AIHW National Perinatal Statistics Unit (unpublished).																										

Table 5A.3.3

Baby's birthweight, by live births and fetal deaths, 2000–2002 (a), (b)

	All births (d)											
	Aboriginal and Torres Strait Islander (c)						Non-Indigenous (c)					
	Live births			Fetal deaths (e)			Unknown			Total births		
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%
Australia												
Low birthweight (<2500g)	3 356	12.9	251	74.3	3 607	13.6	43 606	6.1	3 660	76.6	3	42.9
Very low birthweight (<1500g)	614	2.4	202	59.8	816	3.1	7 350	1.0	3 041	63.7	1	14.3
Extremely low birthweight (<1000g)	285	1.1	176	52.1	461	1.7	3 154	0.4	2 674	56.0	1	14.3
All births	26 106	100.0	338	100.0	26 444	100.0	717 858	100.0	4 776	100.0	7	100.0
Mean birthweight (g)	3 169	..	1 485	..	3 148	..	3 382	..	1 341	..	2 450	..
NSW												
Low birthweight (<2500g)	765	12.0	51	71.8	816	12.7	14 686	5.8	1 163	73.8	3	75.0
Very low birthweight (<1500g)	124	1.9	37	52.1	161	2.5	2 366	0.9	934	59.3	1	25.0
Extremely low birthweight (<1000g)	47	0.7	33	46.5	80	1.2	1 028	0.4	812	51.5	1	25.0
All births	6 371	100.0	71	100.0	6 442	100.0	251 617	100.0	1 576	100.0	4	100.0
Mean birthweight (g)	3 203	..	1 462	..	3 185	..	3 389	..	1 435	..	2 128	..
Victoria												
Low birthweight (<2500g)	157	13.1	15	78.9	172	14.1	11 607	6.3	1 026	78.4	—	—
Very low birthweight (<1500g)	26	2.2	12	63.2	38	3.1	2 011	1.1	871	66.6	—	—
Extremely low birthweight (<1000g)	13	1.1	10	52.6	23	1.9	888	0.5	772	59.0	—	—
All births	1 203	100.0	19	100.0	1 222	100.0	185 316	100.0	1 308	100.0	—	—
Mean birthweight (g)	3 176	..	1 386	..	3 148	..	3 368	..	1 256	..	3 354	..
All indigenous status (c)												
Live births	10 392	1.4	6	2.2	8 161	1.1	3 321	63.0	1	14.3	11 483	1.5
Fetal deaths (e)	22	8.1	2	100.0	—	—	24	8.8	48 100	6.3	4 013	76.2
Total births	100.0	—	—	—	—	—	6	2.2	8 161	1.1	3 321	63.0
Unknown	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%
Fatal deaths (e)	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%
Total births	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%

Table 5A.3.3

Baby's birthweight, by live births and fetal deaths, 2000–2002 (a), (b)

queensland	Aboriginal and Torres Strait Islander (c)						Non-Indigenous (c)						Unknown Indigenous status (c)						All births (d)												
	Live births			Fetal deaths (e)			Total births			Live births			Fetal deaths (e)			Total births			Live births			Fetal deaths (e)			Total births						
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%					
Low birthweight (<2500g)	950	11.6	88	79.3	1 038	12.5	8 648	6.2	721	76.9	—	—	9 369	6.7	5	17.2	1	100.0	—	—	6	20.0	9 603	6.5	810	77.1	—	—	10 413	7.0	
Very low birthweight (<1500g)	185	2.3	70	63.1	255	3.1	1 484	1.1	613	65.4	—	—	2 097	1.5	1	3.4	—	0.0	—	—	1	3.3	1 670	1.1	683	65.0	—	—	2 353	1.6	
Extremely low birthweight (<1000g)	95	1.2	60	54.1	155	1.9	618	0.4	537	57.2	—	—	1 155	0.8	—	0.0	—	0.0	—	—	0.0	—	713	0.5	597	56.9	—	—	1 310	0.9	
All births	8 211	100.0	111	100.0	8 322	100.0	138 912	100.0	938	100.0	—	—	139 850	100.0	29	100.0	1	100.0	—	—	30	100.0	147 152	100.0	1 050	100.0	—	—	148 202	100.0	
Mean birthweight (g)	3 210	..	1 472	..	3 187	..	3 396	..	1 235	..	—	—	3 382	..	3 161	..	1 775	..	—	—	..	3 114	..	3 385	..	1 341	..	—	..	3 371	..
WA																															
Low birthweight (<2500g)	669	14.5	49	76.6	718	15.3	4 073	5.8	386	79.9	—	—	4 459	6.3	—	—	—	—	—	—	—	—	—	4 742	6.4	436	79.5	—	—	5 177	6.9
Very low birthweight (<1500g)	127	2.7	43	67.2	170	3.6	631	0.9	332	68.7	—	—	963	1.4	—	—	—	—	—	—	—	—	—	758	1.0	375	68.6	—	—	1 133	1.5
Extremely low birthweight (<1000g)	57	1.2	38	59.4	95	2.0	252	0.4	296	61.3	—	—	548	0.8	—	—	—	—	—	—	—	—	—	309	0.4	334	61.1	—	—	643	0.9
All births	4 629	100.0	64	100.0	4 693	100.0	69 776	100.0	483	100.0	—	—	70 259	100.0	—	—	—	—	—	—	—	—	—	74 405	100.0	547	100.0	—	—	74 952	100.0
Mean birthweight (g)	3 118	..	1 390	..	3 094	..	3 368	..	1 236	..	—	—	3 354	..	—	—	—	—	—	—	—	—	—	3 353	..	1 254	..	—	..	3 338	..
SA																															
Low birthweight (<2500g)	226	17.7	14	60.9	240	18.4	3 258	6.3	253	77.8	—	—	3 511	6.7	—	—	—	—	—	—	—	—	—	3 484	6.6	267	76.7	—	—	3 751	7.0
Very low birthweight (<1500g)	54	4.2	13	56.5	67	5.1	611	1.2	207	63.7	—	—	818	1.6	—	—	—	—	—	—	—	—	—	665	1.3	220	63.2	—	—	885	1.7
Extremely low birthweight (<1000g)	26	2.0	12	52.2	38	2.9	271	0.5	181	55.7	—	—	452	0.9	—	—	—	—	—	—	—	—	—	297	0.6	193	55.5	—	—	490	0.9
All births	1 278	100.0	23	100.0	1 301	100.0	51 695	100.0	325	100.0	—	—	52 020	100.0	—	—	—	—	—	—	—	—	—	52 973	100.0	348	100.0	—	—	53 321	100.0
Mean birthweight (g)	3 069	..	1 745	..	3 046	..	3 371	..	1 367	..	—	—	3 359	..	—	—	—	—	—	—	—	—	—	3 364	..	1 392	..	—	..	3 351	..

Table 5A.3.3

Baby's birthweight, by live births and fetal deaths, 2000–2002 (a), (b)

			All births (d)																																																
			Unknown Indigenous status (c)				Non-Indigenous (c)				Unknown Indigenous status (c)				Non-Indigenous (c)				Unknown Indigenous status (c)				Non-Indigenous (c)				Unknown Indigenous status (c)				Non-Indigenous (c)				Unknown Indigenous status (c)				Non-Indigenous (c)												
	Aboriginal and Torres Strait Islander (c)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	Live births				Fetal deaths (e)	Total births	
Tas	Live births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%	Live births no.	%	Total births no.	%											
Low birthweight (<2500g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na											
Very low birthweight (<1500g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na											
Extremely low birthweight (<1000g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na											
All births	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na											
Mean birthweight (g)	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na											
ACT (f)																																																			
Low birthweight (<2500g)	np	np	<5	np	42	23.1	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np											
Very low birthweight (<1500g)	np	np	<5	np	18	9.9	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np											
Extremely low birthweight (<1000g)	np	np	<5	np	9	4.9	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np											
All births	np	np	<5	np	182	100.0	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np	np												
Mean birthweight (g)	3 037	..	545	..	2 996	..	3 396	..	1 562	..	3 383	..	3 467
NT																																																			
Low birthweight (<2500g)	550	13.0	31	66.0	581	13.6	406	6.0	32	78.0	—	0.0	438	6.4	7	12.3	—	—	—	—	7	12.3	—	—	—	—	963	8.7	63	71.6	—	0.0	1 026	9.2																	
Very low birthweight (<1500g)	83	2.0	24	51.1	107	2.5	62	0.9	26	63.4	—	0.0	88	1.3	3	5.3	—	—	—	—	3	5.3	148	1.3	50	56.8	—	0.0	198	1.8																					
Extremely low birthweight (<1000g)	41	1.0	20	42.6	61	1.4	19	0.3	24	58.5	—	0.0	43	0.6	3	5.3	—	—	—	—	3	5.3	63	0.6	44	50.0	—	0.0	107	1.0																					
All births	4 235	100.0	47	100.0	4 282	100.0	6 779	100.0	41	100.0	3	100.0	6 823	100.0	57	100.0	—	—	—	—	57	100.0	11 071	100.0	88	100.0	3	100.0	11 162	100.0																					
Mean birthweight (g)	3 126	..	1 656	..	3 110	..	3 364	..	1 309	..	3 095	..	3 351	..	3 216	..	3 216	..	3 272	..	3 272	..	3 493	..	3 272	..	3 272	..	3 258	..	3 258	..	3 258	..	3 258	..	3 258	..	3 258	..											

(a) Data are presented in a three year grouping due to small numbers from year to year. Tasmania's not available because the not stated category for Indigenous status was not able to be distinguished from the non-Indigenous category.
(b) Data on Indigenous births relate to babies born to Indigenous mothers only, and excludes babies born to non-Indigenous mothers and Indigenous fathers. Thus, the information is not based on the total count of Indigenous babies.

(c) Totals from the Australian sections of births broken-down by Indigenous status do not include data for Tasmania.
(d) The Australian all births section totals include data for Tasmania.
(e) The denominator for the fetal death percentages is fetal deaths rather than births.

(f) ACT data includes non-ACT residents who gave birth in the ACT.
(g) NA Not available... Not applicable. — Nil or rounded to zero. np Not published.

Source: National Perinatal Data Collection, AIHW, National Perinatal Statistics Unit (unpublished).

Table 5A.3.4 Baby's birthweight, by live births and fetal deaths, 1999–2001 (a), (b)

Table 5A.3.4

	Non-Indigenous												Unknown Indigenous status												All births (c)											
	Aboriginal and Torres Strait Islander						Total births						Fetal deaths						Live births						Fetal deaths						Unknown					
	Live births no.	Live births %	Fetal deaths no.	Fetal deaths %	Total births no.	Total births %	Live births no.	Live births %	Fetal deaths no.	Fetal deaths %	Total births no.	Total births %	Live births no.	Live births %	Fetal deaths no.	Fetal deaths %	Total births no.	Total births %	Live births no.	Live births %	Fetal deaths no.	Fetal deaths %	Total births no.	Total births %	Live births no.	Live births %	Fetal deaths no.	Fetal deaths %	Total births no.	Total births %						
Australia (excluding Tas and ACT)																																				
Low birthweight (<2500g)	3 241	13	255	75	3 496	13 –	42 418	6	3 609	76	3	13	46 030	6	15	10	2	100	–	–	17	11	47 740	6	4 038	75	3	13	51 781	7						
Very low birthweight (<1500g)	594	2	207	61	801	3 –	7 192	1	2 949	62	1	4	10 142	1	3	2	–	–	–	–	3	2	8 200	1	3 290	62	1	4	11 491	1						
Extremely low birthweight (<1000g)	276	1	181	53	457	2 –	3 126	0	2 583	55	1	4	5 710	1	2	1	–	–	–	2	1	3 569	0	2 878	54	1	4	6 448	1							
All births	25 807	100	342	100	26 149	100 –	70 6230	100	4 733	100	23	100	710 986	100	152	100	2	100	1	100	155	100	763 635	100	5 349	100	24	100	769 008	100						
Mean birthweight (g)	3 171	..	1 432	..	3 149	..	3 382	..	1 373	..	–	..	3 369	..	–	..	–	..	–	..	3 375	..	1 393	..	–	..	3 362	..								
NSW																																				
Low birthweight (<2500g)	743	12	55	74	798	13	14 738	6	1 163	73	3	15	15 904	6	10	10	1	100	–	–	11	10	15 491	6	1 219	73	3	14	16 713	6						
Very low birthweight (<1500g)	121	2	35	47	156	2	2 429	1	908	57	1	5	3 338	1	2	2	–	–	–	–	2	2	2 552	1	943	57	1	5	3 496	1						
Extremely low birthweight (<1000g)	48	1	31	42	79	1	1 050	0	783	49	1	5	1 834	1	2	2	–	–	–	2	2	1 100	–	814	49	1	5	1 915	1							
All births	6 264	100	74	100	6 338	100	25 3013	100	1 591	100	20	100	264 624	100	104	100	1	100	1	100	106	100	259 381	100	1 666	100	21	100	261 068	100						
Mean birthweight (g)	3 199	..	1 552	..	3 181	..	3 390	..	1 482	..	–	..	3 378	..	–	..	–	..	–	3 385	..	1 486	..	–	..	3 373	..							
Victoria																																				
Low birthweight (<2500g)	167	14	14	74	181	14	11 500	6	1 020	77	–	–	12 520	7	–	–	–	–	–	–	–	–	–	11 667	6	1 034	77	–	–	12 701	7					
Very low birthweight (<1500g)	24	2	12	63	36	3	1 933	1	844	64	–	–	2 777	1	–	–	–	–	–	–	–	–	–	1 957	1	856	64	–	–	2 813	2					
Extremely low birthweight (<1000g)	17	1	8	42	25	2	856	0	736	55	–	–	1 592	1	–	–	–	–	–	–	–	–	–	873	0	744	55	–	–	1 617	1					
All births	1 236	100	19	100	1 255	100	184 813	100	1 328	100	–	–	186 141	100	6	100	–	–	–	–	6	100	186 055	100	1 347	100	–	–	187 402	100						
Mean birthweight (g)	3 164	..	1 538	..	3 139	..	3 387	..	1 338	..	–	..	3 353	..	–	..	–	..	–	3 366	..	1 340	..	–	..	3 352	..							

Table 5A.3.4 Baby's birthweight, by live births and fetal deaths, 1999–2001 (a), (b)

Table 5A.3.4

Queensland	Aboriginal and Torres Strait Islander				Non-Indigenous				Unknown Indigenous status				All births (c)			
	Live births		Fetal deaths		Total births		Live births		Fetal deaths		Unknown		Total births		Live births	
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%
Low birthweight (<2500g)	923	11	86	80	1 009	12	8 368	6	754	79	—	—	9 122	7	3	17
Very low birthweight (<1500g)	182	2	73	68	255	3	1 501	1	634	66	—	—	2 135	2	1	6
Extremely low birthweight (<1000g)	87	1	64	60	151	2	648	0	556	58	—	—	1 204	1	—	—
All births	8 346	100	107	100	8 453	100	138 319	100	956	100	—	—	139 275	100	18	100
Mean birthweight (g)	3 220	..	1 321	..	3 196	..	3 399	..	1 298	..	—	—	3 385	..	—	..
WA																
Low birthweight (<2500g)	666	15	47	78	713	15	4 127	6	387	79	—	—	4 514	6	—	—
Very low birthweight (<1500g)	139	3	42	70	181	4	644	1	331	67	—	—	975	1	—	—
Extremely low birthweight (<1000g)	61	1	38	63	99	2	282	0	295	60	—	—	557	1	—	—
All births	4 574	100	60	100	4 634	100	70 814	100	491	100	—	—	71 305	100	—	—
Mean birthweight (g)	3 105	..	1 246	..	3 081	..	3 369	..	1 288	..	—	—	3 355	..	—	..
SA																
Low birthweight (<2500g)	211	17	15	63	226	17	3 238	6	248	78	—	—	3 486	7	—	—
Very low birthweight (<1500g)	48	4	11	46	59	5	610	1	204	64	—	—	814	2	—	—
Extremely low birthweight (<1000g)	24	2	9	38	33	3	284	1	187	59	—	—	471	1	—	—
All births	1 275	100	24	100	1 299	100	52 479	100	317	100	—	—	52 796	100	—	—
Mean birthweight (g)	3 105	..	1 856	..	3 082	..	3 373	..	1 331	..	—	—	3 361	..	—	..
Overcoming Indigenous Disadvantage: Key Indicators 2007																
ATTACHMENT TABLES																

Table 5A.3.4

Baby's birthweight, by live births and fetal deaths, 1999–2001 (a), (b)

Table 5A.3.4

Tas	Aboriginal and Torres Strait Islander		Non-Indigenous		Unknown Indigenous status		All births (c)		
	Live births		Fetal deaths	Total births	Unknown		Total births	Unknown	
	no.	%	no.	%	no.	%	no.	%	no.
Low birthweight (<2500g)	na	na	na	na	na	na	na	na	na
Very low birthweight (<1500g)	na	na	na	na	na	na	na	na	na
Extremely low birthweight (<1000g)	na	na	na	na	na	na	na	na	na
All births	na	na	na	na	na	na	na	na	na
Mean birthweight (g)	na	na	na	na	na	na	na	na	na
ACT									
Low birthweight (<2500g)	np	np	np	np	np	np	np	np	np
Very low birthweight (<1500g)	np	np	np	np	np	np	np	np	np
Extremely low birthweight (<1000g)	np	np	np	np	np	np	np	np	np
All births	np	np	np	np	np	np	np	np	np
Mean birthweight (g)	np	np	np	np	np	np	np	np	np
NT									
Low birthweight (<2500g)	531	13	38	66	569	14	447	7	37
Very low birthweight (<1500g)	80	2	34	59	114	3	75	1	28
Extremely low birthweight (<1000g)	39	1	31	53	70	2	26	0	52
All births	4 112	100	58	100	4 170	100	6 792	100	3 100
Mean birthweight (g)	3 123	..	1 472	..	3 101	..	3 349	..	3 336

(a) Data are presented in a three year grouping due to small numbers from year to year. ACT not published due to small numbers.

(b) Data on Indigenous births relate to babies born to Indigenous mothers only, and excludes babies born to non-Indigenous mothers and Indigenous fathers. Thus, the information is not based on the total count of Indigenous babies.

(c) The Australian all births section totals includes data for Tas and ACT.

na Not available. – Not applicable. – Nil or rounded to zero. np Not published.

Source: National Perinatal Data Collection, AIHW National Perinatal Statistics Unit (unpublished).

Total births

no.

%

%

All births (c)

no.

%

%

na

Table 5A.3.5

Baby's birthweight, by live births and fetal deaths, 1998–2000 (a), (b), (c)

	Indigenous						Non-Indigenous						All births					
	Live births		Fetal deaths		Total births		Live births		Fetal deaths		Total births		Live births		Fetal deaths		Total births	
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%
Australia																		
Number of low birthweight (<2500g)	3 101	11.9	258	76.6	3 359	12.8	44 503	6.0	3 715	74.7	48 218	6.5	47 604	6.2	3 973	74.8	51 577	6.7
Number of very low birthweight (<1500g)	569	2.2	212	62.9	781	3.0	7 519	1.0	2 971	59.7	10 490	1.4	8 088	1.1	3 183	59.9	11 271	1.5
Number of extremely low birthweight (<1000g)	276	1.1	177	52.5	453	1.7	3 261	0.4	2 588	52.0	5 849	0.8	3 537	0.5	2 765	52.1	6 302	0.8
All births	25 970	100.0	337	100.0	26 307	100.0	738 721	100.0	4 974	100.0	743 695	100.0	764 691	100.0	5 311	100.0	770 002	100.0
Mean birthweight (g)	3 179	..	1 371	..	3 156	..	3 382	..	1 432	..	3 369	..	3 375	..	1 428	..	3 362	..
NSW																		
Number of low birthweight (<2500g)	684	11.0	45	68.2	729	11.6	14 526	5.7	1 180	71.2	15 706	6.2	15 210	5.9	1 225	71.1	16 435	6.3
Number of very low birthweight (<1500g)	109	1.8	30	45.5	139	2.2	2 346	0.9	895	54.0	3 241	1.3	2 455	0.9	925	53.7	3 380	1.3
Number of extremely low birthweight (<1000g)	50	0.8	26	39.4	76	1.2	993	0.4	775	46.8	1 768	0.7	1 043	0.4	801	46.5	1 844	0.7
All births	6 203	100.0	66	100.0	6 269	100.0	253 590	100.0	1 657	100	255 247	100	259 793	100	1 723	100	261 516	100
Mean birthweight (g)	3 207	..	1 658	..	3 191	..	3 391	..	1 546	..	3 378	..	3 386	..	1 550	..	3 374	..
Victoria																		
Number of low birthweight (<2500g)	171	13.4	12	85.7	183	14.2	11 542	6.2	983	76.8	12 525	6.7	11 713	6.3	995	76.9	12 708	6.8
Number of very low birthweight (<1500g)	27	2.1	9	64.3	36	2.8	1 963	1.1	793	62.0	2 756	1.5	1 990	1.1	802	62.0	2 792	1.5
Number of extremely low birthweight (<1000g)	20	1.6	5	35.7	25	1.9	860	0.5	677	52.9	1 537	0.8	880	0.5	682	52.7	1 562	0.8
All births	1 273	100.0	14	100.0	1 287	100.0	184 777	100.0	1 280	100.0	186 057	100.0	186 050	100.0	1 294	100.0	187 344	100.0
Mean birthweight (g)	3 157	..	1 353	..	3 137	..	3 367	..	1 386	..	3 354	..	3 366	..	1 386	..	3 352	..

Table 5A.3.5

Baby's birthweight, by live births and fetal deaths, 1998–2000 (a), (b), (c)

Table 5A.3.5

	Indigenous						Non-Indigenous						All births					
	Live births			Fetal deaths			Total births			Live births			Fetal deaths			Total births		
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%
Queensland																		
Number of low birthweight (<2500g)	907	10.8	88	82.2	995	11.7	8 322	6.1	738	78.9	9 060	6.6	9 229	6.4	826	79.3	10 055	6.9
Number of very low birthweight (<1500g)	171	2.0	77	72.0	248	2.9	1 463	1.1	611	65.3	2 074	1.5	1 634	1.1	688	66.0	2 322	1.6
Number of extremely low birthweight (<1000g)	86	1.0	66	61.7	152	1.8	657	0.5	541	57.9	1 198	0.9	743	0.5	607	58.3	1 350	0.9
All births	8 374	100.0	107	100.0	8 481	100.0	136 801	100.0	935	100.0	137 736	100.0	145 175	100.0	1 042	100.0	146 217	100.0
Mean birthweight (g)	3 223	..	1 234	..	3 198	..	3 396	..	1 310	..	3 382	..	3 386	..	1 302	..	3 372	..
WA																		
Number of low birthweight (<2500g)	606	13.3	46	78.0	652	14.2	4 160	5.8	388	79.2	4 548	6.3	4 766	6.2	505	434	79.053	5 200 6.782
Number of very low birthweight (<1500g)	127	2.8	41	69.5	168	3.7	642	0.9	328	66.9	970	1.3	769	1.0	101	369	67.213	1 138 1.484
Number of extremely low birthweight (<1000g)	57	1.3	35	59.3	92	2.0	262	0.4	289	59.0	551	0.8	319	0.4	19	324	59.016	643 0.839
All births	4 542	100.0	59	100	4 601	100	71 586	100	490	100.0	72 076	100.0	76 128	100	549	100	76 677	100
Mean birthweight (g)	3 122	..	1 273	..	3 098	..	3 372	..	1 304	..	3 358	..	3 357	..	1 301	..	3 342	..
SA																		
Number of low birthweight (<2500g)	203	15.7	17	77.3	220	16.7	3 349	6.3	252	78.8	3 601	6.7	3 552	6.5	269	78.7	3 821	6.9
Number of very low birthweight (<1500g)	46	3.6	13	59.1	59	4.5	589	1.1	207	64.7	796	1.5	635	1.2	220	64.3	855	1.6
Number of extremely low birthweight (<1000g)	20	1.5	10	45.5	30	2.3	254	0.5	188	58.8	442	0.8	274	0.5	198	57.9	472	0.9
All births	1 293	100.0	22	100.0	1 315	100.0	53 489	100.0	320	100.0	53 809	100.0	54 782	100.0	342	100.0	55 124	100.0
Mean birthweight (g)	3 123	..	1 580	..	3 097	..	3 374	..	1 317	..	3 361	..	3 368	..	1 334	..	3 355	..

Table 5A.3.5

Baby's birthweight, by live births and fetal deaths, 1998–2000 (a), (b), (c)

Table 5A.3.5

	Indigenous						Non-Indigenous						All births						
	Live births			Fetal deaths			Live births			Fetal deaths			Live births			Fetal deaths			Total births
	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.	%	no.
Tasmania	12	4.9	5	83.3	17	6.7	1 133	6.4	84	51.9	1 217	6.8	1 145	6.4	89	53.0	1 234	6.8	
Number of low birthweight (<2500g)	n.p.	0.8	n.p.	50.0	5	2.0	212	1.2	62	38.3	274	1.5	214	1.2	65	38.7	279	1.5	
Number of very low birthweight (<1500g)	n.p.	0.4	n.p.	50.0	4	1.6	83	0.5	51	31.5	134	0.7	84	0.5	54	32.1	138	0.8	
Number of extremely low birthweight (<1000g)	All births	247	100.0	6	100.0	253	100.0	17 714	100.0	162	100.0	17 876	100.0	17 961	100.0	168	100.0	18 129	100.0
Mean birthweight (g)	3 417	..	1 476	..	3 370	..	3 389	..	1 949	..	3 377	..	3 389	..	1 930	..	3 377
ACT	n.p.	16.5	n.p.	100.0	30	17.4	980	7.0	58	70.7	1 038	7.4	1 008	7.1	60	71.4	1 068	7.5	
Number of low birthweight (<2500g)	n.p.	6.5	n.p.	50.0	12	7.0	228	1.6	50	61.0	278	2.0	239	1.7	51	60.7	290	2.0	
Number of very low birthweight (<1500g)	n.p.	1.2	n.p.	..	n.p.	1.2	121	0.9	45	54.9	166	1.2	123	0.9	45	53.6	168	1.2	
Number of extremely low birthweight (<1000g)	All births	n.p.	100.0	n.p.	100.0	172	100.0	13 945	100.0	82	100.0	14 027	100.0	14 115	100.0	84	100.0	14 199	100.0
Mean birthweight (g)	3 119	..	1 748	..	3 103	..	3 388	..	1 533	..	3 372	..	3 385	..	1 538	..	3 369
NT	n.p.	12.7	43	70.5	533	13.6	491	7.2	32	66.7	523	7.6	981	9.2	75	68.8	1 056	9.8	
Number of low birthweight (<2500g)	490
Number of very low birthweight (<1500g)	76	2.0	38	62.3	114	2.9	76	1.1	25	52.1	101	1.5	152	1.4	63	57.8	215	2.0	
Number of extremely low birthweight (<1000g)	All births	40	1.0	32	52.5	72	1.8	31	0.5	22	45.8	53	0.8	71	0.7	54	49.5	125	1.2
Mean birthweight (g)	3 868	100.0	61	100.0	3 929	100.0	6 819	100.0	48	100.0	6 867	100.0	10 687	100.0	109	100.0	10 796	100.0	
Mean birthweight (g)	3 120	..	1 319	..	3 093	..	3 341	..	1 585	..	3 329	..	3 261	..	1 437	..	3 244

(a) Data are presented in a three year grouping due to small numbers from year to year.

(b) Data on Indigenous births relate to babies born to Indigenous mothers only, and excludes babies born to non-Indigenous mothers and Indigenous fathers. Thus, the information is not based on the total count of Indigenous babies.

(c) Birthweight data on babies born to Indigenous mothers residing in the ACT or Tasmania should be viewed with caution as they are based on small numbers of births.

.. Not applicable. n.p. Not published.
Source: AIHW National Perinatal Statistics Unit (unpublished).

Table 5A.4.1 Persons with a long-term hearing problem: type of ear/hearing problem, by age and Indigenous status, 2004-05

**Table 5A.4.1
Persons with a long-term hearing problem: type of ear/hearing problem, by age and Indigenous status, 2004-05¹**

	Deafness						Other hearing problems (a) (b)						Total with hearing problems (b) (c) (d)			
	Indigenous		Non-Indigenous		Indigenous		Non-Indigenous		Indigenous		Non-Indigenous		Indigenous		Non-Indigenous	
	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Proportion (%)	RSE ratio (e)	Rate ratio (e)	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)	Rate ratio (e)	
0-14	4.5	10.8	1.2	15.1	3.8	5.4	11.3	1.9	14.1	2.8	9.5	8.0	3.0	10.5	3.1	
15-24	5.4	15.6	2.5	15.2	2.1	2.6	22.9	1.4	22.4	1.9	7.7	12.8	3.8	13.6	2.0	
25-34	10.3	9.9	5.1	10.5	2.0	(f) 2.6	21.4	(f) 2.3	13.1	1.1	12.1	8.8	7.1	8.0	1.7	
35-44	10.7	11.6	8.4	6.3	1.3	(f) 4.7	21.9	(f) 2.7	11.6	1.7	14.5	9.5	10.7	5.4	1.4	
45-54	16.7	12.9	10.8	5.3	1.5	(f) 3.6	31.4	(f) 3.4	11.1	1.1	19.6	12.1	13.6	4.8	1.4	
55 and over	(f) 25.3	8.7	(f) 26.3	2.7	1.0	(f) 3.2	25.7	(f) 5.3	5.9	0.6	(f) 27.0	8.3	(f) 29.8	2.5	0.9	
Total (d)	8.8	4.6	10.2	1.9	0.9	3.8	10.0	3.0	3.8	1.3	12.2	4.1	12.7	1.7	1.0	

- (a) Totals for other hearing problems have been age standardised.
- (b) Includes Tinnitus, Meniere's disease/Vertiginous syndrome.
- (c) Includes 'Type of ear/hearing problem' not known.
- (d) Components may not add to total as persons may have reported more than one type of ear/hearing problem.
- (e) The ratio is equal to the rate for Indigenous persons divided by the rate for non-Indigenous persons.
- (f) The row or individual figure should be viewed with caution as the differences in proportions were found to be statistically insignificant.
– Nil or rounded to zero.

Source : Australian Bureau of Statistics (ABS) 2006, National Aboriginal and Torres Strait Islander Health Survey 2004-05 (NATSIHS 2004-05), Cat. no. 4715.0 (unpublished); National Health Survey 2004-05 (NHS 2004-05), Cat. no. 4362.0 (unpublished).

Table 5A.4.2

Persons with a long-term hearing problem: type of ear/hearing problem, by age and Indigenous status, 2001 (per cent of population)

	Deafness										Other hearing problems (a) (b)										Total with hearing problems (b) (c) (d)			
	Indigenous		Non-Indigenous		Indigenous		Non-Indigenous		Indigenous		Non-Indigenous		Indigenous		Non-Indigenous		Indigenous		Non-Indigenous		Indigenous		Non-Indigenous	
	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Rate ratio (e)	Proportion (%)	RSE (%)	Rate ratio (e)	
0–14	6.5	14.8	1.9	9.3	3.4	5.8	16.9	2.9	8.9	2.0	11.2	10.3	4.7	5.6	2.4									
15–24	9.0	21.7	3	13.3	3.0	(f) 4.9	45.3	(f) 2.2	16.6	3.5	13.5	18.8	5.2	11.0	2.6									
25–34	9.8	17.4	5.3	11.9	1.8	(f) 3.8	25.6	(f) 2.2	14.3	1.7	13.0	16.0	7.2	10.6	1.8									
35–44	(f) 10.6	23.6	(f) 8.5	6.3	1.3	(f) 6.1	32.5	(f) 3.1	8.1	2.3	(f) 16	17.0	(f) 11.2	4.9	1.4									
45–54	(f) 16.1	16.2	(f) 13.1	5.3	1.2	(f) 8	26.7	(f) 5.1	8.2	1.6	(f) 22.8	13.7	(f) 17.2	4.3	1.7									
55 and over	(f) 26	17.9	(f) 27.4	2.5	1.0	(f) 5	42.1	(f) 6.6	6.7	0.9	(f) 29.1	14.6	(f) 31.9	2.5	1.0									
Total (d)	(f) 10	8.1	(f) 10.7	2.2	1.0	5.6	17.1	(f) 3.9	3.9	1.4	(f) 14.6	6.4	(f) 13.8	1.9	1.1									

- (a) Totals for other hearing problems have been age standardised.
- (b) Includes Titis media, Tinnitus and Meniere's disease/Vertiginous syndrome.
- (c) Includes 'Type of ear/hearing problem' not known.
- (d) Components may not add to total as persons may have reported more than one type of ear/hearing problem.
- (e) The ratio is equal to the rate for Indigenous persons divided by the rate for non-Indigenous persons.
- (f) The row or individual figure should be viewed with caution as the differences in proportions were found to be statistically insignificant.
 - Nil or rounded to zero.

Source: ABS 2002, National Health Survey 2001 (NHS 2001), Cat. no. 4364.0 (unpublished).

Table 5A.4.3

Persons with a long-term hearing problem: type of ear/hearing problem, by remoteness areas and Indigenous status, 2004-05 (per cent of population)

Type of ear/hearing problem	Indigenous						Non-Indigenous			Rate ratio (b)	
	Remote (g)		Non-remote (g)		Total		AS Total (a)		AS Total (a)		
	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)	Proportion (%)		
Deafness	9.2	7.8	8.6	5.7	8.8	4.6	np	np	np	np	
Otitis media	3.6	15.3	1.6	15.9	2.2	11.2	1.4	11.2	0.5	9.6	
Other (c)	1.6	22.3	2.1	14.1	1.9	12.2	(f) 2.4	12.9	(f) 2.4	4.5	
Total with a hearing condition (d)	13.4	6.8	11.8	5.1	12	4	np	np	np	np	

- (a) Statistical testing indicates that deafness and total hearing problems are not suitable for age standardisation.
- (b) The ratio is equal to the rate for Indigenous persons divided by the rate for non-Indigenous persons.
- (c) Includes Tinnitus and Meniere's disease/Vertiginous syndrome.
- (d) Includes 'Type of ear/hearing problem' not known.
- (e) Components may not add to total as persons may have reported more than one type of ear/hearing problem.
- (f) The row or individual figure should be viewed with caution as the differences in proportions were found to be not statistically significant.
- (g) Non-remote includes major cities and inner and outer regional areas. Remote includes remote and very remote areas. See glossary for definitions of remoteness areas.
 - Nil or rounded to zero. np Not published.

Source: ABS 2006, NATSIHS 2004-05 (unpublished); NHS 2004-05 (unpublished).

Table 5A.4.4 Persons with a long-term hearing problem: type of ear/hearing problem, by remoteness areas and Indigenous status, 2001 (per cent of population)

Type of ear/hearing problem	Indigenous						Non-Indigenous						Rate ratio (b)	
	Remote (g)		Non-remote (g)		Total	RSE (%)	AS Total (a)		RSE (%)	AS Total (a)		RSE (%)		
	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)			Proportion (%)	RSE (%)		Proportion (%)	RSE (%)			
Deafness	9.6	9.6	10.2	11.1	10	8.1	np	np	np	np	np	np	np	
Otitis media	4.5	18	1.3	29.4	2.2	18.3	(f) 1.2	22.4	(f) 0.8	7.4	7.4	1.5	1.5	
Other (c)	3.4	23	3.5	21.2	3.4	17.8	(f) 4.1	21.8	(f) 3.1	4.4	4.4	1.0	1.0	
Total with a hear	15.9	8.7	14.1	8.6	14.6	6.4	np	np	np	np	np	np	np	

(a) Statistical testing indicates that deafness and total hearing problems are not suitable for age standardisation.

(b) The ratio is equal to the rate for Indigenous persons divided by the rate for non-Indigenous persons.

(c) Includes Tinnitus and Meniere's disease/Vertiginous syndrome.

(d) Includes 'Type of ear/hearing problem' not known.

(e) Components may not add to total as persons may have reported more than one type of ear/hearing problem.

(f) The row or individual figure should be viewed with caution as the differences in proportions were found to be not statistically significant.

(g) Non-remote includes major cities and inner and outer regional areas. Remote includes remote and very remote areas. See glossary for definitions of remoteness areas.

– Nil or rounded to zero. np Not published.

Source: ABS 2002, NHS 2001 (unpublished).

Table 5A.4.5 **Persons with otitis media, by age and Indigenous status, 2004-05 (per cent of population) (a)**

	<i>Indigenous</i>		<i>Non-Indigenous</i>	
	Proportion (%)	RSE (%)	Proportion (%)	RSE (%)
0-4	3.6	19.8	1.8	25.4
5-9	6.5	20.4	2.3	23.9
10-14	2.7	20.8	0.7	32.9
Total	2.2	11.2	0.5	9.5

(a) Estimate has a relative standard error of between 25 to 50 per cent and should be used with caution. Data are subject to sampling variability too high for most practical purposes.

Source: ABS 2006, NATSIHS 2004-05 (unpublished); NHS 2004-05 (unpublished).

Table 5A.4.6

Table 5A.4.6 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA, and public hospitals in NT, 2004-05 (a) (b) (c)

Principal diagnosis	Indigenous			Non-Indigenous			Non-Indigenous (d)		
	no.	no.	no.	no.	no.	Total	Indigenous	no.	per 1000
People aged 0-3 years									
Diseases of external ear (H60-H62)	18	94	np	np	5 697	5 697	0.6	0.3	0.3
Diseases of middle ear and mastoid (H65-H75)	275	4 956	466	466	1 870	1 870	9.5	15.3	15.3
suppurative and unspecified otitis media (H66)	164	1 596	110	110	7	7	5.7	4.8	4.8
Diseases of inner ear (H80-H83)	-	7	-	-	-	-	-	-	-
Other disorders of ear (H90-H95)	10	277	9	296	9	296	0.3	0.8	0.8
People aged 4-14 years									
Diseases of external ear (H60-H62)	27	316	12	12	355	355	0.3	0.3	0.3
Diseases of middle ear and mastoid (H65-H75)	558	5 419	485	485	6 462	6 462	7.0	5.5	5.5
suppurative and unspecified otitis media (H66)	148	1 140	74	74	1 362	1 362	1.9	1.1	1.1
Diseases of inner ear (H80-H83)	-	13	np	np	-	-	-	-	-
Other disorders of ear (H90-H95)	19	261	7	287	7	287	0.2	0.2	0.2

(a) Hospital separation is the discharge, transfer, death or change of episode of care of an admitted patient (see glossary for a detailed definition). Separations are based on ICD-10-AM classification.

(b) Data are based on state of usual residence.

(c) Identification of Indigenous patients is incomplete and completeness varies across jurisdictions. The AIHW has advised that only data for Queensland, WA, SA and the NT are considered to be acceptable for analytical purposes. Data for NSW, Vic, Tasmania and the ACT were withheld by AIHW due to high rates of Indigenous under-identification (see chapter 2 and appendix 4 for more information).

(d) Includes separations where Indigenous status was not reported.

- Nil or rounded to zero np Not published.

Source: AIHW National hospital morbidity database (unpublished).

Table 5A.4.7

Table 5A.4.7 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA, and public hospitals in NT, 2003-04 (a) (b) (c)

Principal diagnosis	Indigenous			Non-Indigenous			Non-Indigenous (d)		
	no.	no.	no.	no.	Total	Indigenous	per 1000	per 1000	
People aged 0-3 years									
Diseases of external ear (H60-H62)	10	80	9	99	99	0.3	0.3	0.3	
Diseases of middle ear and mastoid (H65-H75)	299	5 054	492	5 845	10 4	15.7	15.7	15.7	
suppurative and unspecified otitis media (H66)	187	1 774	92	2 053	6.5	5.3	5.3	5.3	
Diseases of inner ear (H80-H83)	-	np	-	np	-	np	np	np	
Other disorders of ear (H90-H95)	17	251	np	np	0.6	0.7	0.7	0.7	
People aged 4-14 years									
Diseases of external ear (H60-H62)	14	278	20	312	0.2	0.3	0.3	0.3	
Diseases of middle ear and mastoid (H65-H75)	608	5 420	605	6 633	7.7	5.6	5.6	5.6	
suppurative and unspecified otitis media (H66)	156	1 225	90	1 471	2.0	1.2	1.2	1.2	
Diseases of inner ear (H80-H83)	np	22	np	26	np	np	np	np	
Other disorders of ear (H90-H95)	16	222	6	244	0.2	0.2	0.2	0.2	

(a) Hospital separation is the discharge, transfer, death or change of episode of care of an admitted patient (see glossary for a detailed definition). Separations are based on ICD-10-AM classification.

(b) Data are based on state of usual residence.

(c) Identification of Indigenous patients is incomplete and completeness varies across jurisdictions. The AIHW has advised that only data for Queensland, WA, SA and the NT are considered to be acceptable for analytical purposes. Data for NSW, Vic, Tasmania and the ACT were withheld by AIHW due to high rates of Indigenous under-identification (see chapter 2 and appendix 4 for more information).

(d) Includes separations where Indigenous status was not reported.

- Nil or rounded to zero np Not published.

Source: AIHW National hospital morbidity database (unpublished).

Table 5A.4.8

Table 5A.4.8 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA and public hospitals in NT, 2002-03 (a) (b) (c)

Principal diagnosis	Principal diagnosis			Non-Indigenous			Non-Indigenous		
	Indigenous	Indigenous	Not stated	Total	Indigenous	Total	Indigenous	Total	(d)
	no.	no.	no.	no.	no.	no.	per 1000	no.	per 1000
People aged 0-3 years									
Diseases of external ear (H60-H62)	9	75	9	93	0.3	0.2			
Diseases of middle ear and mastoid (H65-H75)	308	5 149	594	6 051	10.7	14.5			
suppurative and unspecified otitis media (H66)	160	1 596	150	1 906	5.6	4.5			
Diseases of inner ear (H80-H83)	-	12	-	12	-	-			
Other disorders of ear (H90-H95)	16	228	7	251	0.6	0.6			
People aged 4-14 years									
Diseases of external ear (H60-H62)	24	254	26	304	0.3	0.2			
Diseases of middle ear and mastoid (H65-H75)	505	5 563	723	6 791	6.4	5.2			
suppurative and unspecified otitis media (H66)	134	1 136	104	1 374	1.7	1.1			
Diseases of inner ear (H80-H83)	np	9	np	11	np	-			
Other disorders of ear (H90-H95)	17	187	11	215	0.2	0.2			

(a) Hospital separation is the discharge, transfer, death or change of episode of care of an admitted patient (see glossary for a detailed definition). Separations are based on ICD-10-AM classification.

- (b) Data are based on state of usual residence.
- (c) Identification of Indigenous patients is incomplete and completeness varies across jurisdictions. The AIHW has advised that only data for Queensland, WA, SA and the NT are considered to be acceptable for analytical purposes. Data for NSW, Vic, Tasmania and the ACT were withheld by AIHW due to high rates of Indigenous under-identification (see chapter 2 and appendix 4 for more information).
- (d) Includes separations where Indigenous status was not reported.

- Nil or rounded to zero np Not published.

Source: AIHW National hospital morbidity database (unpublished).

Table 5A.4.9

Table 5A.4.9 Age specific separations where the principal diagnosis was diseases of the ear and mastoid process, Qld, WA, SA and public hospitals in NT, 2001-02 (a) (b) (c)

Principal diagnosis	Indigenous			Non-Indigenous			Non-Indigenous (d)		
	no.	no.	no.	no.	Total	Indigenous	no.	per 1000	per 1000
People aged 0-3 years									
Diseases of external ear (H60-H62)	8	80	12	100	0.3	0.3	0.3	0.3	0.3
Diseases of middle ear and mastoid (H65-H75)	369	5 192	571	6 132	12.9	12.9	16.2	16.2	16.2
suppurative and unspecified otitis media (H66)	200	1 550	143	1 893	7.0	7.0	4.8	4.8	4.8
Diseases of inner ear (H80-H83)	-	10	-	10	-	-	-	-	-
Other disorders of ear (H90-H95)	19	246	11	276	0.7	0.7	0.7	0.7	0.7
People aged 4-14 years									
Diseases of external ear (H60-H62)	15	252	36	303	0.2	0.2	0.3	0.3	0.3
Diseases of middle ear and mastoid (H65-H75)	565	5 831	682	7 078	7.3	7.3	6.2	6.2	6.2
suppurative and unspecified otitis media (H66)	159	1 102	145	1 406	2.0	2.0	1.2	1.2	1.2
Diseases of inner ear (H80-H83)	-	19	-	19	-	-	-	-	-
Other disorders of ear (H90-H95)	28	155	9	192	0.4	0.4	0.2	0.2	0.2

(a) Hospital separation is the discharge, transfer, death or change of episode of care of an admitted patient (see glossary for a detailed definition). Separations are based on ICD-10-AM classification.

(b) Data are based on state of usual residence.

(c) Identification of Indigenous patients is incomplete and completeness varies across jurisdictions. The AIHW has advised that only data for Queensland, WA, SA and the NT are considered to be acceptable for analytical purposes. Data for NSW, Vic, Tasmania and the ACT were withheld by AIHW due to high rates of Indigenous under-identification (see chapter 2 and appendix 4 for more information).

(d) Includes separations where Indigenous status was not reported.

- Nil or rounded to zero np Not published.

Source: AIHW National hospital morbidity database (unpublished).

Table 5A.5.1

Table 5A.5.1 Decayed and decay-free infant teeth in 4- to 10-year-old children enrolled in school dental services in NSW, SA and the NT, by Indigenous status and location (a) (b)

Age	Indigenous				Non-Indigenous			
	Metropolitan		Rural		Metropolitan		Rural	
dmft=0	%	SE	%	SE	%	SE	%	SE
4 years	41.6	4.4	31.0	2.3	69.7	0.6	64.3	0.5
5 years	47.7	2.5	31.8	1.6	70.9	0.2	62.2	0.5
6 years	36.6	2.8	24.2	1.7	64.5	0.4	56.8	0.3
7 years	51.8	2.2	28.8	1.5	62.9	0.3	54.2	0.5
8 years	43.6	2.9	28.6	1.6	58.9	0.4	51.6	0.6
9 years	48.9	2.3	38.1	1.6	62.2	0.3	54.2	0.5
10 years	57.0	2.9	44.9	1.8	66.1	0.4	59.7	0.6
dmft	Mean	SE	Mean	SE	Mean	SE	Mean	SE
4 years	2.89	0.33	3.56	0.19	1.31	0.03	1.39	0.05
5 years	2.63	0.20	4.13	0.15	1.20	0.01	1.68	0.03
6 years	2.74	0.18	4.09	0.15	1.42	0.02	1.82	0.03
7 years	2.09	0.13	3.43	0.12	1.44	0.01	1.86	0.02
8 years	2.20	0.15	3.16	0.11	1.48	0.02	1.87	0.03
9 years	1.98	0.12	2.28	0.09	1.26	0.01	1.60	0.02
10 years	1.39	0.13	1.66	0.08	1.00	0.01	1.27	0.02
d/dmft (c)	%	SE	%	SE	%	SE	%	SE
4 years	86.8	3.0	82.1	1.9	64.2	0.6	68.7	0.5
5 years	59.1	2.4	70.8	1.5	55.5	0.3	52.5	0.5
6 years	54.0	2.9	68.3	1.9	48.5	0.4	43.9	0.3
7 years	44.6	2.2	65.7	1.6	43.4	0.3	38.2	0.5
8 years	43.4	2.9	61.5	1.7	39.3	0.4	33.8	0.5
9 years	40.4	2.2	55.6	1.7	37.0	0.3	30.9	0.4
10 years	41.1	2.9	55.5	1.8	35.1	0.4	29.7	0.5

SE Standard error.

- (a) d=number of untreated decayed teeth; m=number of missing teeth; f=number of filled teeth; dmft=d+m+f (for deciduous infant teeth). The indicator dmft=0 refers to decay-free teeth. Per cent d/dmft is the proportion of decayed teeth and is calculated to indicate levels of unmet treatment need.
- (b) Values have been calculated using data collected over three 12-month periods; 2000 for NSW, 2002 for NT, and 2003 for SA. Of the 326 099 children examined, 10 743 (3.2 per cent) were Indigenous.
- (c) Proportion of untreated decayed infant teeth.

Source: Jamieson, L., Armfield, J.A., Roberts-Thomson, K.F. 2006a, 'The role of location in Indigenous and non-Indigenous child oral health', *Journal of Public Health Dentistry*, vol. 66, pp. 123–130.

Table 5A.5.2

Table 5A.5.2 Decayed and decay-free permanent teeth in 6- to 14-year-old children enrolled in school dental services in NSW, SA and the NT, by Indigenous status and location (a) (b)

Age	Indigenous				Non-Indigenous			
	Metropolitan		Rural		Metropolitan		Rural	
DMFT=0	%	SE	%	SE	%	SE	%	SE
6 years	93.8	1.4	90.4	1.2	96.7	0.1	97.1	0.1
7 years	89.5	1.3	83.5	1.3	90.4	0.2	90.1	0.3
8 years	78.7	2.4	74.6	1.5	86.0	0.3	86.7	0.4
9 years	74.7	2.0	68.9	1.6	83.1	0.2	83.3	0.4
10 years	66.6	2.8	60.6	1.8	78.9	0.3	79.1	0.5
11 years	67.8	2.2	60.5	1.7	75.4	0.3	75.3	0.4
12 years	59.3	3.2	53.5	2.2	70.6	0.4	70.9	0.6
13 years	57.4	3.2	52.3	3.1	70.0	0.3	65.6	0.6
14 years	41.3	5.1	40.0	4.1	62.1	0.6	59.1	0.9
DMFT	Mean	SE	Mean	SE	Mean	SE	Mean	SE
6 years	0.13	0.04	0.18	0.03	0.08	0.00	0.05	0.00
7 years	0.22	0.03	0.36	0.03	0.22	0.00	0.22	0.01
8 years	0.44	0.06	0.53	0.04	0.30	0.01	0.28	0.01
9 years	0.56	0.05	0.68	0.04	0.39	0.01	0.36	0.01
10 years	0.77	0.08	1.00	0.06	0.47	0.01	0.44	0.01
11 years	0.80	0.07	1.05	0.06	0.59	0.01	0.59	0.01
12 years	1.08	0.11	1.33	0.09	0.77	0.01	0.70	0.02
13 years	1.44	0.16	1.77	0.17	0.88	0.01	0.90	0.01
14 years	1.83	0.31	2.27	0.24	1.18	0.03	1.19	0.04
D/DMFT	%	SE	%	SE	%	SE	%	SE
6 years	78.8	2.4	73.0	1.8	65.4	0.4	66.5	0.31
7 years	55.6	2.2	71.5	1.5	55.3	0.3	54.7	0.46
8 years	59.8	2.9	71.5	1.6	50.6	0.4	48.7	0.57
9 years	50.2	2.3	62.8	1.6	42.6	0.3	39.7	0.45
10 years	45.0	2.9	64.2	1.7	39.0	0.4	37.4	0.56
11 years	44.5	2.3	59.3	1.7	37.3	0.3	35.0	0.46
12 years	38.5	3.2	63.2	2.1	37.3	0.4	35.7	0.64
13 years	50.3	3.3	59.4	3.0	37.1	0.3	33.1	0.6
14 years	44.5	5.2	52.8	4.2	37.1	0.6	34.7	0.88

SE Standard error.

(a) D=number of untreated decayed teeth; M=number of missing teeth; F=number of filled teeth; DMFT=D+M+F (for deciduous permanent adult teeth). The indicator DMFT=0 refers to decay-free teeth. Per cent D/DMFT is the proportion of decayed teeth and is calculated to indicate levels of unmet treatment need.

(b) Values have been calculated using data collected over three 12-month periods; 2000 for NSW, 2002 for NT, and 2003 for SA. Of the 326 099 children examined, 10 743 (3.2 per cent) were Indigenous.

Table 5A.5.2

Table 5A.5.2 Decayed and decay-free permanent teeth in 6- to 14-year-old children enrolled in school dental services in NSW, SA and the NT, by Indigenous status and location (a) (b)

(c) Proportion of untreated decayed permanent teeth.

Source: Jamieson, L., Armfield, J.A., Roberts-Thomson, K.F. 2006a, 'The role of location in Indigenous and non-Indigenous child oral health', *Journal of Public Health Dentistry*, vol. 66, pp. 123–130.

Table 5A.5.3

Table 5A.5.3 Mean number of teeth with decay for children enrolled in the NT School Dental Service, by age and Indigenous status, 2002-03 (a) (b)

Age	Indigenous		Non-Indigenous	
	Mean	SE	Mean	SE
dmft (infant teeth)				
4 years	3.40	0.19	1.01	0.07
5 years	4.34	0.18	1.47	0.08
6 years	4.14	0.18	1.76	0.09
7 years	3.73	0.15	1.80	0.08
8 years	3.17	0.13	1.89	0.08
9 years	2.44	0.12	1.71	0.07
10 years	1.74	0.09	1.21	0.06
DMFT (permanent teeth)				
6 years	0.15	0.03	0.07	0.01
7 years	0.27	0.03	0.15	0.02
8 years	0.47	0.04	0.21	0.02
9 years	0.60	0.05	0.30	0.03
10 years	0.93	0.06	0.36	0.03
11 years	1.02	0.07	0.52	0.04
12 years	1.24	0.10	0.71	0.05
13 years	1.81	0.27	0.80	0.10

SE Standard error.

(a) D=number of untreated decayed teeth; M=number of missing teeth; F=number of filled teeth;
 DMFT=D+M+F (for permanent adult teeth); dmft=refers to deciduous infant teeth and is derived in the same way as DMFT. The indicator DMFT=0 (dmft=0 for infant teeth) refers to decay-free teeth.

(b) Of the 12 584 children examined, 4 417 (35.1 per cent) were Indigenous.

Source: Jamieson, L., Armfield, J.A., Roberts-Thomson, K.F. 2006b, 'Oral health inequalities among Indigenous and non-Indigenous children in the Northern Territory of Australia', *Community Dentistry and Oral Epidemiology*, vol. 34, pp. 267–76.

Table 5A.5.4

Table 5A.5.4 Mean tooth decay (dmft) for 4- to 10-year-old children enrolled in the NT School Dental Service, by SEIFA index of relative socio-economic disadvantage, age and Indigenous status, 2002-03 (a) (b) (c) (d)

Age SEIFA	Indigenous		Non-Indigenous	
	Mean	SE	Mean	SE
dmft				
4 years				
1	4.26	0.28	0.69	0.16
2	3.58	0.61	1.45	0.22
3	2.48	0.26	0.89	0.08
4	na	na	1.94	0.51
5 years				
1	5.31	0.25	1.36	0.25
2	3.32	0.38	1.85	0.17
3	2.99	0.28	1.28	0.10
4	na	na	1.23	0.37
6 years				
1	4.78	0.26	1.84	0.30
2	3.91	0.41	1.73	0.16
3	2.87	0.30	1.78	0.20
4	na	na	1.57	0.36
7 years				
1	4.28	0.21	1.57	0.28
2	3.57	0.37	1.82	0.14
3	2.65	0.24	1.81	0.11
4	na	na	1.96	0.40
8 years				
1	3.47	0.18	1.89	0.24
2	3.13	0.29	2.12	0.15
3	2.45	0.21	1.77	0.10
4	na	na	1.48	0.33
9 years				
1	2.54	0.17	1.91	0.23
2	2.29	0.33	1.82	0.14
3	2.38	0.20	1.62	0.09
4	na	na	1.80	0.33
10 years				
1	1.93	0.13	1.36	0.19
2	1.72	0.28	1.48	0.14
3	1.31	0.16	1.05	0.08
4	na	na	1.00	0.20

Table 5A.5.4

Table 5A.5.4 Mean tooth decay (dmft) for 4- to 10-year-old children enrolled in the NT School Dental Service, by SEIFA index of relative socio-economic disadvantage, age and Indigenous status, 2002-03 (a) (b) (c) (d)

SEIFA Socio-Economic Indices for Areas. SE Standard error.

- (a) d=number of untreated decayed teeth; m=number of missing teeth; f=number of filled teeth;
 $dmft=d+m+f$ (for deciduous infant teeth).
- (b) The SEIFA was developed by the ABS using data derived from the 2001 Census of Population and Housing, and uses a range of measures to rank areas based on their relative social and economic well-being. It takes into account variables relating to income, educational attainment, unemployment, and dwellings without motor vehicles. In the table above, '1' denotes the most disadvantaged areas and '4' denotes the least disadvantaged areas.
- (c) Of the 12 584 children examined, 4 417 (35.1 per cent) were Indigenous.
- (d) For Indigenous children, no cases fell into the least disadvantaged category (4).

na Not applicable.

Source: Jamieson, L., Armfield, J.A., Roberts-Thomson, K.F. 2006b, 'Oral health inequalities among Indigenous and non-Indigenous children in the Northern Territory of Australia', *Community Dentistry and Oral Epidemiology*, vol. 34, pp. 267-76.

Table 5A.5.5

Table 5A.5.5 Mean tooth decay (DMFT) for 6- to 13-year-old children enrolled in the NT School Dental Service, by SEIFA index of relative socio-economic disadvantage, age and Indigenous status, 2002-03 (a) (b) (c) (d)

Age SEIFA	Indigenous		Non-Indigenous	
	Mean	SE	Mean	SE
DMFT				
6 years				
1	0.15	0.03	0.00	0.00
2	0.28	0.11	0.08	0.02
3	0.09	0.04	0.06	0.01
4	na	na	0.10	0.08
7 years				
1	0.33	0.05	0.14	0.07
2	0.29	0.08	0.15	0.03
3	0.15	0.05	0.12	0.02
4	na	na	0.36	0.14
8 years				
1	0.51	0.06	0.14	0.06
2	0.43	0.08	0.22	0.03
3	0.36	0.07	0.20	0.03
4	na	na	0.42	0.18
9 years				
1	0.74	0.08	0.24	0.07
2	0.52	0.10	0.29	0.05
3	0.40	0.06	0.31	0.03
4	na	na	0.42	0.13
10 years				
1	1.10	0.09	0.24	0.07
2	0.77	0.11	0.43	0.06
3	0.67	0.10	0.36	0.04
4	na	na	0.23	0.09
11 years				
1	1.19	0.11	0.65	0.15
2	0.83	0.12	0.60	0.07
3	0.70	0.10	0.45	0.05
4	na	na	0.61	0.16
12 years				
1	1.35	0.13	0.43	0.13
2	1.27	0.21	0.82	0.09
3	0.81	0.17	0.69	0.07
4	na	na	0.67	0.22

Table 5A.5.5

Table 5A.5.5 Mean tooth decay (DMFT) for 6- to 13-year-old children enrolled in the NT School Dental Service, by SEIFA index of relative socio-economic disadvantage, age and Indigenous status, 2002-03 (a) (b) (c) (d)

Age SEIFA	Indigenous		Non-Indigenous	
	Mean	SE	Mean	SE
13 years				
1	1.93	0.13	0.85	0.36
2	1.62	0.21	0.89	0.18
3	1.33	0.17	0.83	0.16
4	na	na	0.29	0.16

SEIFA Socio-Economic Indices for Areas. SE Standard error.

- (a) D=number of untreated decayed teeth; M=number of missing teeth; F=number of filled teeth; DMFT=D+M+F (for deciduous permanent adult teeth).
- (b) The SEIFA was developed by the ABS using data derived from the 2001 Census of Population and Housing, and uses a range of measures to rank areas based on their relative social and economic well-being. It takes into account variables relating to income, educational attainment, unemployment, and dwellings without motor vehicles. In the table above, '1' denotes the most disadvantaged areas and '4' denotes the least disadvantaged areas.
- (c) Of the 12 584 children examined, 4 417 (35.1 per cent) were Indigenous.
- (d) For Indigenous children, no cases fell into the least disadvantaged category (4).

na Not applicable.

Source: Jamieson, L., Armfield, J.A., Roberts-Thomson, K.F. 2006b, 'Oral health inequalities among Indigenous and non-Indigenous children in the Northern Territory of Australia', *Community Dentistry and Oral Epidemiology*, vol. 34, pp. 267-76.

Table 5A.5.6

Table 5A.5.6 SA, infant and permanent tooth decay, mean dmft and DMFT by Indigenous status and age, 2002 Child Dental Health Survey (a)

Age	Indigenous	Non-Indigenous
<i>Mean dmft (infant teeth)</i>		
4 years	3.91	1.31
5 years	3.40	1.50
6 years	2.71	1.77
7 years	3.04	1.88
8 years	2.89	1.93
9 years	2.74	1.81
10 years	1.81	1.50
<i>Mean DMFT (permanent teeth)</i>		
7 years	0.44	0.20
8 years	0.62	0.36
9 years	0.79	0.45
10 years	0.75	0.53
11 years	1.24	0.61
12 years	1.31	0.81
13 years	1.95	1.02
14 years	2.28	1.40
15 years	2.27	1.65

- (a) D=number of untreated decayed teeth; M=number of missing teeth; F=number of filled teeth; DMFT=D+M+F (for permanent adult teeth); dmft=refers to deciduous infant teeth and is derived in the same way as DMFT. The indicator DMFT=0 (dmft=0 for infant teeth) refers to decay-free teeth.

Source: Ellershaw, A., Spencer, A.J., Slade, G.D. 2005. *Oral Health in South Australia 2004*. Australian Research Centre for Population Oral Health, University of Adelaide.

Table 5A.5.7

Table 5A.5.7 SA, proportion of children with calculus and gingival bleeding by Indigenous status and age, children attending school dental service, 2002 Child Dental Health Survey

	<i>Unit</i>	<i>Indigenous</i>	<i>Non-Indigenous</i>
6 to 7 years			
Healthy	%	63.8	80.2
Bleeding (a)	%	33.4	15.0
Calculus (b)	%	2.8	4.8
8 to 9 years			
Healthy	%	51.8	66.7
Bleeding (a)	%	43.5	23.5
Calculus (b)	%	4.7	9.8
10 to 11 years			
Healthy	%	50.4	62.4
Bleeding (a)	%	40.1	24.2
Calculus (b)	%	9.5	13.4
12 to 13 years			
Healthy	%	47.8	59.7
Bleeding (a)	%	36.7	22.6
Calculus (b)	%	15.5	17.7
14 to 15 years			
Healthy	%	42.7	57.8
Bleeding (a)	%	39.4	19.4
Calculus (b)	%	17.9	22.8

(a) Gingival bleeding is bleeding of the gums associated with gingivitis, which is an inflammation of the gums surrounding the teeth caused by a buildup of plaque or food particles.

(b) Calculus is a hard deposit of mineralized plaque which is attached to crowns and/or roots of teeth.

Source: Ellershaw, A., Spencer, A.J., Slade, G.D. 2005. *Oral Health in South Australia 2004*. Australian Research Centre for Population Oral Health, University of Adelaide.

Table 5A.5.8

Table 5A.5.8 Dental hospitalisations for children aged 2 to 14 years, per 100 000 children, by sociodemographic characteristics and Indigenous status, 2002-03 (a) (b)

	<i>Indigenous</i>		<i>Non-Indigenous</i>	
	Mean	SE	Mean	SE
Sex				
Males	641.8	26.4	587.5	5.3
Females	544.2	25.0	607.5	5.6
Age-group				
less than 5 years	906.9	38.6	625.6	7.0
5 to 9 years	667.4	32.7	687.7	7.1
10 to 14 years	180.7	17.9	479.8	5.9
Location				
Major city	539.9	31.8	577.4	4.7
Regional	529.0	25.5	636.9	6.9
Remote	778.5	41.7	571.5	20.9

SE Standard error.

- (a) Admission rates are based on hospitalisations data from 1 297 public and private hospitals obtained from the Australian Institute of Health and Welfare National Hospital Morbidity Database for 2002-03. Rates were calculated by dividing the number of hospital dental general anaesthetic admissions for each age by the estimated resident population (ERP) of the same age and multiplying by 100 000 (Jamieson and Roberts-Thomson 2006). ERP counts for all ages were provided by the ABS (unpublished).
- (b) Of the 24 874 children examined, 1 062 (4.3 per cent) were Indigenous.

Source: Jamieson, L., Roberts-Thomson, K.F. 2006, 'Indigenous children and receipt of hospital dental care in Australia', *International Journal of Paediatric Dentistry*, vol. 16, pp. 327-34.

Table 5A.5.9

Table 5A.5.9 Hospital dental procedure rates for children aged 2 to 14 years, per 100 000 children, by sex and Indigenous status, 2002-03 (a) (b)

	<i>Indigenous</i>		<i>Non-Indigenous</i>	
	<i>Mean</i>	<i>SE</i>	<i>Mean</i>	<i>SE</i>
<i>Male</i>				
Extraction	601.5	25.5	450.5	4.7
Pulpal	59.9	8.1	88.2	2.1
Restoration	396.7	20.7	310.9	3.9
Other	91.5	10.0	129.9	2.5
<i>Female</i>				
Extraction	520.0	24.4	470.5	4.9
Pulpal	52.9	7.8	83.0	2.1
Restoration	317.5	19.1	291.3	3.9
Other	58.7	8.2	117.3	2.5

SE Standard error.

- (a) Procedure rates are based on separation data from 1 297 public and private hospitals obtained from the Australian Institute of Health and Welfare National Hospital Morbidity Database for 2002-03. Data were collected for administrative purposes by hospital-employed dentists and recorded in standardized International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian modification (ICD-10-AM) codes, which are patient record codes used throughout Australian hospitals. There were over 80 ICD-10-AM dental procedure codes which were grouped into 'extraction', 'pulpal care', 'restorative care', or 'other' (included the provision of splints, crowns or bridges).
- (b) Of the 24 874 children examined, 1 062 (4.3 per cent) were Indigenous.

Source: Jamieson, L., Roberts-Thomson, K.F. 2006, 'Indigenous children and receipt of hospital dental care in Australia', *International Journal of Paediatric Dentistry*, vol. 16, pp. 327-34.

Table 5A.5.10

Table 5A.5.10 Hospital dental procedure rates for children aged 2 to 14 years, per 100 000 children, by age group and Indigenous status, 2002-03 (a) (b)

	<i>Indigenous</i>		<i>Non-Indigenous</i>	
	Mean	SE	Mean	SE
Less than 5 years old				
Extraction	865.4	37.7	390.1	5.5
Pulpal	99.7	12.9	136.1	3.3
Restoration	579.7	30.9	426.3	5.8
Other	129.6	14.7	199.0	3.9
5 to 9 years old				
Extraction	628.7	31.7	530.5	6.2
Pulpal	62.9	10.1	115.1	2.9
Restoration	409.5	25.6	418.5	5.6
Other	85.4	11.7	165.6	3.5
10 to 14 years old				
Extraction	164.7	17.1	456.5	5.8
Pulpal	3.5	2.5	8.6	0.8
Restoration	65.5	10.8	65.9	2.2
Other	7.1	3.5	10.5	0.9

SE Standard error.

- (a) Procedure rates are based on separation data from 1 297 public and private hospitals obtained from the Australian Institute of Health and Welfare National Hospital Morbidity Database for 2002-03. Data were collected for administrative purposes by hospital-employed dentists and recorded in standardized International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian modification (ICD-10-AM) codes, which are patient record codes used throughout Australian hospitals. There were over 80 ICD-10-AM dental procedure codes which were grouped into 'extraction', 'pulpal care', 'restorative care', or 'other' (included the provision of splints, crowns or bridges).
- (b) Of the 24 874 children examined, 1 062 (4.3 per cent) were Indigenous.

Source: Jamieson, L., Roberts-Thomson, K.F. 2006, 'Indigenous children and receipt of hospital dental care in Australia', *International Journal of Paediatric Dentistry*, vol. 16, pp. 327-34.

Table 5A.5.11

Table 5A.5.11 Hospital dental procedure rates for children aged 2 to 14 years, per 100 000 children, by remoteness area and Indigenous status, 2002-03 (a) (b) (c)

	<i>Indigenous</i>		<i>Non-Indigenous</i>	
	Mean	SE	Mean	SE
Major city				
Extraction	498.6	30.5	436.6	4.2
Pulpal	58.3	10.5	98.6	0.2
Restoration	280.3	22.9	283.5	3.3
Other	122.3	15.2	147.3	2.4
Regional (d)				
Extraction	506.8	24.9	508.9	6.2
Pulpal	32.1	0.6	59.6	0.2
Restoration	318.1	19.8	331.1	5.0
Other	34.5	6.5	76.5	2.4
Remote (e)				
Extraction	738.0	40.6	418.0	17.9
Pulpal	99.0	14.9	104.1	0.9
Restoration	524.3	34.3	342.4	16.2
Other	94.5	14.6	151.9	10.8

SE Standard error.

- (a) Procedure rates are based on separation data from 1 297 public and private hospitals obtained from the Australian Institute of Health and Welfare National Hospital Morbidity Database for 2002-03. Data were collected for administrative purposes by hospital-employed dentists and recorded in standardized International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian modification (ICD-10-AM) codes, which are patient record codes used throughout Australian hospitals. There were over 80 ICD-10-AM dental procedure codes which were grouped into 'extraction', 'pulpal care', 'restorative care', or 'other' (included the provision of splints, crowns or bridges).
- (b) Of the 24 874 children examined, 1 062 (4.3 per cent) were Indigenous.
- (c) Remoteness areas are derived from the Australian Standard Geographical Classification (ASGC) developed by the ABS. The ASGC Remoteness classification identifies a region in Australia as having a particular degree of remoteness. Remoteness areas comprise five categories: major cities, inner regional, outer regional, remote, and very remote. The degree of remoteness of each place is determined using the Accessibility/Remoteness Index of Australia (ARIA). Places with similar ARIA scores are aggregated to form the remoteness areas in each State and Territory. For more information on how ARIA is defined see chapter 2 of the Report.
- (d) Includes 'inner regional' and 'outer regional' classifications of the ASGC.
- (e) Includes 'remote' and 'very remote' classifications of the ASGC.

Source: Jamieson, L., Roberts-Thomson, K.F. 2006, 'Indigenous children and receipt of hospital dental care in Australia', *International Journal of Paediatric Dentistry*, vol. 16, pp. 327-34.